

NATIONAL INSTITUTE FOR FUSION SCIENCE

L-Mode Confinement Model Based on Transport-MHD Theory in Tokamaks

K. Itoh, M. Yagi, S.-I. Itoh, A. Fukuyama and M. Azumi

(Received – Sep. 22, 1992)

NIFS-192

Oct. 1992

RESEARCH REPORT
NIFS Series

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

L-Mode Confinement Model

Based on Transport-MHD Theory in Tokamaks

K. Itoh*, M. Yagi[†], S.-I. Itoh**, A. Fukuyama^{††}, and M. Azumi[†]

* National Institute for Fusion Science, Nagoya 464-01, Japan

† Japan Atomic Energy Research Institute, Naka,

Ibaraki 311-01, Japan

** Institute for Applied Mechanics, Kyushu University 87,

Kasuga 816, Japan

†† Faculty of Engineering, Okayama University, Okayama 700, Japan

Abstract

Theory of the L-mode confinement in tokamaks is developed based on the microscopic ballooning instability which is destabilized by the plasma transport below the critical pressure gradient against the ideal MHD instability. The destabilization by the current-diffusivity and the stabilization by the thermal transport and ion viscosity, are analyzed. The least stable mode determines the anomalous transport coefficients. The formula of the thermal transport coefficient is derived, which explains major experimental observations on the L-mode confinement.

Keywords: L-Mode, Anomalous Transport, Current Diffusivity,
Tokamak, Ballooning Instability, Prandtl Number

The transport of the plasma across the magnetic field in tokamaks is much faster than that by the Coulomb collision. It has been known as the anomalous transport¹⁾. The L-mode confinement, in which the energy confinement time τ_E decreases as the heating power P is increased, is observed in all tokamaks^{2,3)}. The L-mode confinement is a generic nature of tokamak plasmas. The database has been made on how τ_E depends on the externally controllable parameters, such as P and the plasma current I_p ³⁾. The radial profile of the effective thermal conductivity χ (the ratio of the energy flux per particle to the temperature gradient ∇T) has been studied⁴⁾. It has been confirmed that the microscopic fluctuations play important roles for the anomalous transport⁴⁾. The ion viscosity is also enhanced and of the same order of the electron and ion thermal conductivities⁵⁾. Ohkawa's model, based on the fluctuations of the scale length of the collisionless skin depth δ ⁶⁾, is one of the few which could explain the large χ -value near edge and at high temperature, but does not fully explain the dependences of τ_E . Theoretical development has been made based on the mixing length estimate⁷⁾, scale invariance method⁸⁾ or one/two point renormalization methods⁹⁾. These gave identical results from the view point of the physics argument¹⁰⁾. No theory has succeeded in explaining the radial shape of χ and the scaling $\tau_E[P, I_p, \dots]$ simultaneously¹¹⁾. The understanding of the L-mode confinement is far from satisfactory.

Microscopic modes grow in tokamaks extracting the free energy source of the pressure gradient. The shear-stabilized plasma can become unstable due to the fluctuation-driven trans-

port, and the analysis on the self-sustaining turbulence has been developed¹²⁻¹⁵). A model of the subcritical turbulence of the interchange mode was proposed to explain the anomalous transport in helical plasmas¹⁵). We apply this method to tokamaks. Below the beta-limit of the ideal magnetohydrodynamic (MHD) mode, the microscopic ballooning mode can be unstable if there is the plasma transport such as the current-diffusivity λ . We find that λ destabilizes the mode while other transport coefficients, χ and the ion viscosity μ , stabilize it. The unstable microscopic ballooning mode enhances the transport coefficients λ , χ and μ . If the anomalous transport is enhanced much, then χ and μ stabilize the mode. The transport coefficients are determined by the marginal stability condition for the least stable mode. The result on χ is compared to the experimental observations and shows agreements.

We study the circular tokamak with the toroidal coordinates (r, θ, ζ) . The reduced set of equations¹⁶) is employed. The current diffusivity is due to the electron viscosity and is kept in the Ohm's law as¹⁷) $E + v \times B = J / \sigma - \nabla_{\perp}^2 \lambda J$ (σ is the conductivity). The equation of motion is given as $n_i m_i \{d(\nabla_{\perp}^2 \phi) / dt - \mu \nabla_{\perp}^4 \phi\} = B \nabla_{\parallel} J_{\parallel} + \nabla p \times \nabla (2r \cos \theta / R)$ and the energy balance equation $dp / dt = \chi \nabla_{\perp}^2 p$ is employed, where m_i is the ion mass, n_i is the ion density, ϕ is the stream function, B is the main magnetic field, p is the plasma pressure, and J is the current.

The ballooning transformation is employed as¹⁸) $\phi(r, \theta, \zeta) = \sum_m \exp(-im\theta + in\zeta) f(\eta) \exp\{im\eta - inq\eta\} d\eta$, (q is the safety factor) since we are interested in the microscopic modes. The linearized

equation is reduced to the ordinary differential equation as¹⁹⁾

$$\frac{d}{d\eta} \frac{F}{\tau + \hat{\Sigma}F + \Lambda F^2} \frac{d\phi}{d\eta} + \frac{\alpha[\kappa + \cos\eta + (s\eta - \alpha \sin\eta)\sin\eta]\phi}{\tau + \chi F} - (\tau + MF)F\phi = 0 \quad (1)$$

We use the normalizations $r/a \rightarrow r$, $t/\tau_{Ap} \rightarrow t$, $\chi\tau_{Ap}/a^2 \rightarrow \chi$, $\mu\tau_{Ap}/a^2 \rightarrow \mu$, $\tau_{Ap}/\mu_0 \sigma a^2 \rightarrow 1/\delta$, $\lambda\tau_{Ap}/\mu_0 a^4 \rightarrow \lambda$, $\tau_{Ap} \equiv a\sqrt{\mu_0 m_i n_i}/B_p$, $\tau\tau_{Ap} \rightarrow \tau$. Notations: $\hat{\Sigma} = n^2 q^2/\delta$, $\Lambda = \lambda n^4 q^4$, $\chi = \chi n^2 q^2$, $M = \mu n^2 q^2$, τ is the growth rate, $s = r(dq/dr)/q$, $F = 1 + (s\eta - \alpha \sin\eta)^2$, $\kappa = -(r/R)(1 - 1/q^2)$ (average well), $B_p = Br/qR$, $\alpha = q^2 \beta'/\epsilon$, $\epsilon = r/R$, a and R are the major and minor radii, β is the pressure divided by the magnetic pressure, and $\beta' \equiv d\beta/d(r/a)$. If we neglect λ , χ and μ , Eq.(1) is reduced to the resistive ballooning equation. The ideal MHD mode equation is recovered by further taking $1/\delta = 0$.

Since κ is small but negative, the interchange mode is stable and the ballooning mode is the most unstable. Equation (1) predicts that the current-diffusive ballooning mode has a large growth rate. We take $1/\delta = 0$ for simplicity. (This is appropriate as is shown at the end of this article.) The growth rate of the short wave length mode, driven by the λ term, is first estimated by the Wentzel-Kramers-Brillouin (WKB) method by neglecting χ and μ terms. We have

$$\frac{\pi}{4} = \int_0^{\eta_c} d\eta \sqrt{\frac{1 + \Lambda F^2/\tau}{F}} \sqrt{\alpha(\cos\eta - (s\eta - \alpha \sin\eta)\sin\eta) - \tau^2 F} \quad (2)$$

where the kernel of the integral vanishes at $\eta = \eta_c$ and the well

term κ is neglected. For the analytic insight, we take the short wave length limit, $\Lambda/\tau \gg 1$, which yields $\sqrt{1/F + \Lambda F/\tau} \approx \sqrt{\Lambda F/\tau}$. By approximating $dF/d\eta \approx 2s\sqrt{F}$, Eq.(2) is reduced to $\pi/4 = [\sqrt{\Lambda/\tau}/2s]\sqrt{\alpha}F_c$, where $F_c = F(\eta_c)$. We also consider the case that the inertia term determines η_c , having the estimate $F_c = \alpha/\tau^2$. Using these limiting approximations, the dispersion relation (2) is written as $\pi/4 = [\sqrt{\Lambda/\tau}/2s]\alpha^{3/2}/\tau^2$, or

$$\tau \approx \lambda^{1/5} (nq)^{4/5} \alpha^{3/5} s^{-2/5}. \quad (3)$$

Since the exponent to η is 1/5, even the very small current diffusivity gives rise to the ballooning instability. The condition $\Lambda/\tau \gg 1$ requires $\lambda n^4 q^4 > \alpha^{3/4}/\sqrt{s}$.

This large growth rate is confirmed for a wide range of parameters by the numerical calculation¹⁹⁾. Figure 1 illustrates α vs τ and λ vs τ , keeping λ/τ and τ/μ constant. The analytic estimation for small λ is confirmed. As the transport coefficients are increased much, the stabilizing effects by τ and μ overcome the destabilizing effect of λ .

The stability boundary is derived. Setting $\tau=0$ in Eq.(1), we have the eigenvalue equation, which determines the relation between τ , λ and μ . We here study the case that the ballooning mode is destabilized by the normal curvature, not by the geodesic curvature, i.e., $1/2 + \alpha > s$. For the strongly localized mode, $s^2 \eta^2 < 1$ and $\eta^2 < 1$, this eigenvalue equation is approximated by the Weber type equation as

$$d^2\phi/d\eta^2 + (\alpha\lambda n^2 q^2/\tau)\{1-(1/2+\alpha-s)\eta^2\} - \mu\lambda n^2 q^2(1+3s^2\eta^2) = 0, \quad (4)$$

and we have the stability boundary as

$$\alpha^{3/2} \lambda \tau^{-3/2} \mu^{-1/2} = N^{-2} (1-N^4)^{-2} \{(1/2+\alpha-s)+3s^2 N^4\} \quad (5)$$

where N is the normalized mode number $N=nq(\lambda\mu/\alpha)^{1/4}$. This result is confirmed by the numerical computation as is shown in Fig.2.

It is shown that, when (τ, λ, μ) increase, the mode is stabilized by the enhanced transport coefficient. The stabilization is possible when both τ and μ are finite. The lower boundary of α for the stability is calculated by obtaining the minimum of the RHS of Eq.(5). When N^{-2} and the term in $\{ \}$ determines the minimum, it is estimated as $\sqrt{6}s$ (neglecting $\alpha-s$ for the simplicity). In the limit $s=0$, the minimum is given as $25\sqrt{5}/32$. We have the stability limit of α for the least stable mode as

$$\alpha^{3/2} = f(s)\sqrt{\mu}\lambda^{3/2}\tau^{-1}, \quad (6)$$

where $f(s) \approx \sqrt{6}s$ or 1.7 ($s \rightarrow 0$).

Based on the stability analysis, we can derive the formula for the anomalous transport coefficient. When the mode amplitude and the associated transport coefficients are small, Eq.(1) gives the instability. In the self-sustained state, the enhanced transport coefficients satisfy Eq.(6) for the given pressure gradient α . This state is thermodynamically stable: The excess

growth of the mode and enhanced transport coefficients lead to the damping of the mode. When the mode amplitude and transport coefficients are small, the mode continues to grow until Eq.(6) is satisfied.

From Eq.(6) χ is expressed in terms of the Prandtl numbers μ/χ and λ/χ . (Note that the current-diffusivity λ is originated from the electron viscosity¹⁷.) We have

$$\chi = \alpha^{3/2} (\lambda/\chi) \sqrt{\chi/\mu} / f(s). \quad (7)$$

In enhancing (λ, μ, χ) by fluctuations, the ratio λ/χ and μ/χ are given to be constant. The relations $\lambda/\chi \approx \delta^2/a^2$ and $\mu/\chi \approx 1$ hold for electrostatic perturbations^{17, 20}). The formula of χ is obtained in an explicit form as

$$\chi = f(s)^{-1} q^2 (R\beta'/r)^{3/2} \delta^2 v_A / R. \quad (8)$$

We here note that the usual method for the estimation of χ by τ/k^2 for the most unstable mode gives the same results as $\chi \sim \alpha^{3/2} (\delta/a)^2$.

This form of χ is consistent with experimental results known for the L-mode. In the following, we compare the theoretical prediction to observations by choosing $n_i = n_e$ and $T_i = T_e$.

(i) The dimensional dependence of χ is $[T]^{1.5} / [a][B]^2$. (ii) Not the local beta value but the *gradient* of β generates χ so that the density and q profiles governs the radial profile of χ . Equation (8) indicates that χ increases towards the edge for the

usual plasma profiles in the L-mode. Experiments on χ is reported that $\chi \sim B_p^{-y} s^{-z}$ with $1 < y < 2$ and $0 < z < 1$ (21), which is consistent with Eq.(6). (iii) The point model argument of the energy balance, $\tau_E = a^2/\chi$ and $2\pi^2 a^2 R n_e T = \tau_E P$, provides the scaling law

$$\tau_E = C a^{0.4} R^{1.2} I_p^{0.8} P^{-0.6} A^{0.5} f^{-0.4} \{n_e \ell_p / \sqrt{A}\}^{0.6}, \quad (9)$$

where C is a numerical coefficient, A is the ion mass number, and ℓ_p is the gradient scale length $(nT)/|\nabla(nT)|$. This result is consistent with the L-mode scaling law, including the dependences on a, R, I_p , P and favourable dependences on the ion mass and magnetic shear^{3, 22}). Slight difference is seen in the the final term in the parenthesis { }, which is discussed later. (iv) $T_e(r)$ profile is predicted from Eq.(6). The peaking parameter, $T(\text{at } q=1)/\langle T \rangle$, only weakly depends on the location of the peak of the power deposition, but depends strongly on $q(a)$ ¹⁹). This explains the 'profile resilience'. (v) Since $\chi \propto \{ \nabla(nT)/n \}^{1.5}$, the thermal diffusion coefficient deduced from the pulse propagation, χ_{HP} , is larger than that evaluated by the power balance χ . If $|\nabla n_i/n_i| \ll |\nabla T/T|$ holds, for the simplicity, we have $\chi_{HP} = 2.5\chi$. (vi) The typical perpendicular wave number of the most unstable mode satisfies

$$k_{\perp 6} \approx 1/\sqrt{\alpha}. \quad (10)$$

It should be noticed that though the dimensional relation $k_{\perp} \propto [B]/[\sqrt{T}]$ holds, k_{\perp} does not scale with the local gyroradius. The

collisionless skin depth is the more relevant length. The correlation time τ_c is estimated as $1/\tau \approx \sqrt{6s}/\alpha\tau_{Ap}$. (vii) The estimation $\bar{p}/p \sim 1/k_{\perp} \ell_p$ shows that the mode amplitude is larger near edge and larger for the high heating power.

In summary, we developed the stability theory of the microscopic ballooning mode in tokamaks under the influence of the anomalous transport coefficients χ , λ and μ . It is found that this mode is unstable when the transport is small and finite, but can be stabilized by the thermal transport and ion viscosity. The stability boundary is obtained, and the transport coefficients are derived. The important role of the collisionless skin depth, pointed out in Ohkawa's model, is confirmed: the consistent calculation for stability gives the further explicit dependence on β' and the geometrical factor. The form of χ is compared with experiments. Major part of the observations on L-mode can be explained by this model simultaneously.

Since χ is dimensionally independent of the density, τ_E derived in Eq.(9) includes the density dependence (see, e.g., [23]). However, the density dependence is offset by the gradient scale length. In L-mode plasmas, the density gradient profile is often steeper than the temperature near edge, and ℓ_p in Eq.(9) would be replaced by $\ell_n = |n_i/\nabla n_i|$. The high density plasma has more steeper edge density profile; Tsuji found that $n_e(0)/\bar{n}_e - 1$ is a decreasing function of n_e and $\ell_n n_e$ is a weak function of the density²⁴⁾. For some dataset of JT-60, Takizuka reported²⁵⁾ the dependence as $\tau_E(\text{thermal}) \propto n_e^{-0.5}$, suggesting that the classification of the dataset by the profile is necessary.

We here take various simplifications for the analytic insight. The resistivity is neglected. The similar analysis can be done for the resistive plasma to have the instability boundary $\alpha = \tau \delta / 2$. The resistive modes²⁶⁾ give higher stability limit of α than the current diffusive modes if $\delta \tau^{1/3} > (\tau / \lambda)^{2/3}$ holds. This condition is usually satisfied and supports the simplification. Other extension is necessary for the case that the mode is driven by the geodesic curvature. For such a case, the results in this article are also confirmed, and will be reported elsewhere¹⁹⁾.

The present result (9) is obtained except for the numerical factor. Nonlinear simulation would give this coefficient and examine the validity of the ansatz of χ that the transport coefficients affecting the microscopic mode is equated to that for the global quantity. Also necessary is the study of the effects such as the diamagnetic drift for kinetic corrections. These research are open for future study.

Authors would like to acknowledge useful discussions with Drs. T. Takizuka, S. Tsuji and K. Ida. This work is partly supported by the Grant-in-Aid for Scientific Research of Ministry of Education Japan and collaboration program between universities and JAERI on fusion.

References

- [1] See, for instance, P. Liewer: Nucl. Fusion **25** 543 (1985).
- [2] R. J. Goldston: Plasma Phys. Control. Fusion **26** 87 (1984).
- [3] P. Yushmanov, et al.: Nucl. Fusion **30** 1999 (1990).
- [4] For a review of present status, see W. A. Houlberg, et al., Phys. Fluids **B2** 2913 (1990); A. Wootton, et al.: *ibid.* **B2** 2879 (1990); D. W. Ross, et al.: *ibid.* **B3** 2251 (1991).
- [5] K. H. Burrell, et al.: Nucl. Fusion **28** 3 (1998);
K. Ida, et al.: in *Plasma Physics and Controlled Nuclear Fusion Research 1990* (IAEA, Vienna, 1991) Vol.2, p577.
- [6] T. Ohkawa: Phys. Letters **67A** 35 (1978).
- [7] B. B. Kadomtsev: in Plasma Turbulence (Academic Press, New York, 1965).
- [8] J. W. Connor and J. B. Taylor: Nucl. Fusion **17** 1047 (1977).
- [9] T. H. Dupree: Phys. Fluids **9** 1773 (1966), **15** 334 (1972).
- [10] M. Yagi: PhD Thesis, Kyoto University (1989).
- [11] For a review, see J. D. Callen: Phys. Fluids **B4** 2142 (1992).
- [12] S. P. Hirshman, K. Molvig: Phys. Rev. Lett. **42** 648 (1979).
- [13] B. D. Scott: Phys. Rev. Lett. **65** 3289 (1990).
- [14] M. Wakatani, et al.: Phys. Fluids **B4** 1754 (1992).
- [15] K. Itoh, S.-I. Itoh, A. Fukuyama, Phys. Rev. Lett. **69** 1050 (1992).
- [16] H. Strauss: Phys. Fluids **20** 1354 (1977).
- [17] J. Schmidt, S. Yoshikawa: Phys. Rev. Lett. **26** 753 (1971).
See also P. J. Kaw, E. J. Valeo, P. H. Rutherford: Phys. Rev. Lett. **43** 1398 (1979), and A. J. Lichtenberg, K. Itoh, S.-I. Itoh, A. Fukuyama: Nucl. Fusion **31** 495 (1992).

- [18] J. W. Connor, R. J. Hastie and J. B. Taylor: Proc. R. Soc. London, **A365** 1 (1979).
- [19] M. Yagi, et al.: paper in preparation.
- [20] S.-I. Itoh: Phys. Fluids **B4** 796 (1992).
- [21] S. Christiansen, et al: in *Controlled Fusion and Plasma Heating* (Proc. 19th Eur. Conf. Innsbruck, 1992) Vol.16C, European Physical Society, (1992) Part 1, p15.
- [22] M. C. Zarnstorff, et al.: in *Plasma Physics and Controlled Nuclear Fusion Research 1990* (IAEA, Vienna, 1991) Vol.1, p109.
- [23] K. Lackner and N. Gottardi: Nucl. Fusion **30** 767 (1990).
- [24] S. Tsuji: Fusion Engineering and Design **15** 311 (1992).
- [25] T. Takizuka: in *Controlled Fusion and Plasma Heating* (Proc. 19th Eur. Conf. Innsbruck, 1992) Vol.16C, European Physical Society, (1992) Part 1, p51.
- [26] B. A. Carreras, P. H. Diamond: Phys. Fluids **B1** 1011 (1989).

Figure Captions

Fig.1 (a) Growth rate of the current-diffusive ballooning mode as a function of α for various values of λ . Dashed line shows the ideal MHD limit. (b) Growth rate as a function of λ for $\beta=0.8\%$. Parameters are: $s=0.4$, $\tau/\lambda=1000$, $\mu=\tau$, $q=3$, $r/l_p=0.6$, $\epsilon=1/8$, $n=30$, and $1/\delta=0$.

Fig.2 Stability boundary as a function of the mode number n . Solid line indicates the analytic formula (5), and dashed line is obtained by the numerical calculation. Parameters are $s=0.4$, $\tau=1.7 \times 10^{-5}$, $\tau/\lambda=1000$, $\mu=\tau$, $q=3$, $r/l_p=0.6$, $\epsilon=1/8$, and $1/\delta=0$.

Fig. 1






Fig. 2



Recent Issues of NIFS Series

- NIFS-151 Y. Takeiri, A. Ando, O. Kaneko, Y. Oka and T. Kuroda, *Negative Ion Extraction Characteristics of a Large Negative Ion Source with Double-Magnetic Filter Configuration*; May 1992
- NIFS-152 T. Tanabe, N. Noda and H. Nakamura, *Review of High Z Materials for PSI Applications*; Jun. 1992
- NIFS-153 Sergey V. Bazdenkov and T. Sato, *On a Ballistic Method for Double Layer Regeneration in a Vlasov-Poisson Plasma*; Jun. 1992
- NIFS-154 J. Todoroki, *On the Lagrangian of the Linearized MHD Equations*; Jun. 1992
- NIFS-155 K. Sato, H. Katayama and F. Miyawaki, *Electrostatic Potential in a Collisionless Plasma Flow Along Open Magnetic Field Lines*; Jun. 1992
- NIFS-156 O.J.W.F.Kardaun, J.W.P.F.Kardaun, S.-I. Itoh and K. Itoh, *Discriminant Analysis of Plasma Fusion Data*; Jun. 1992
- NIFS-157 K. Itoh, S.-I. Itoh, A. Fukuyama and S. Tsuji, *Critical Issues and Experimental Examination on Sawtooth and Disruption Physics*; Jun. 1992
- NIFS-158 K. Itoh and S.-I. Itoh, *Transition to H-Mode by Energetic Electrons*; July 1992
- NIFS-159 K. Itoh, S.-I. Itoh and A. Fukuyama, *Steady State Tokamak Sustained by Bootstrap Current Without Seed Current*; July 1992
- NIFS-160 H. Sanuki, K. Itoh and S.-I. Itoh, *Effects of Nonclassical Ion Losses on Radial Electric Field in CHS Torsatron/Heliotron*; July 1992
- NIFS-161 O. Motojima, K. Akaishi, K. Fujii, S. Fujiwaka, S. Imagawa, H. Ji, H. Kaneko, S. Kitagawa, Y. Kubota, K. Matsuoka, T. Mito, S. Morimoto, A. Nishimura, K. Nishimura, N. Noda, I. Ohtake, N. Ohyabu, S. Okamura, A. Sagara, M. Sakamoto, S. Satoh, T. Satow, K. Takahata, H. Tamura, S. Tanahashi, T. Tsuzuki, S. Yamada, H. Yamada, K. Yamazaki, N. Yanagi, H. Yonezu, J. Yamamoto, M. Fujiwara and A. Iiyoshi, *Physics and Engineering Design Studies on Large Helical Device*; Aug. 1992
- NIFS-162 V. D. Pustovitov, *Refined Theory of Diamagnetic Effect in Stellarators*; Aug. 1992
- NIFS-163 K. Itoh, *A Review on Application of MHD Theory to Plasma*

Boundary Problems in Tokamaks; Aug. 1992

- NIFS-164 Y.Kondoh and T.Sato, *Thought Analysis on Self-Organization Theories of MHD Plasma; Aug. 1992*
- NIFS-165 T. Seki, R. Kumazawa, T. Watari, M. Ono, Y. Yasaka, F. Shimpo, A. Ando, O. Kaneko, Y. Oka, K. Adati, R. Akiyama, Y. Hamada, S. Hidekuma, S. Hirokura, K. Ida, A. Karita, K. Kawahata, Y. Kawasumi, Y. Kitoh, T. Kohmoto, M. Kojima, K. Masai, S. Morita, K. Narihara, Y. Ogawa, K. Ohkubo, S. Okajima, T. Ozaki, M. Sakamoto, M. Sasao, K. Sato, K. N. Sato, H. Takahashi, Y. Taniguchi, K. Toi and T. Tsuzuki, *High Frequency Ion Bernstein Wave Heating Experiment on JIPP T-IIU Tokamak; Aug. 1992*
- NIFS-166 Vo Hong Anh and Nguyen Tien Dung, *A Synergetic Treatment of the Vortices Behaviour of a Plasma with Viscosity; Sep. 1992*
- NIFS-167 K. Watanabe and T. Sato, *A Triggering Mechanism of Fast Crash in Sawtooth Oscillation; Sep. 1992*
- NIFS-168 T. Hayashi, T. Sato, W. Lotz, P. Merkel, J. Nührenberg, U. Schwenn and E. Strumberger, *3D MHD Study of Helias and Heliotron; Sep. 1992*
- NIFS-169 N. Nakajima, K. Ichiguchi, K. Watanabe, H. Sugama, M. Okamoto, M. Wakatani, Y. Nakamura and C. Z. Cheng, *Neoclassical Current and Related MHD Stability, Gap Modes, and Radial Electric Field Effects in Heliotron and Torsatron Plasmas; Sep. 1992*
- NIFS-170 H. Sugama, M. Okamoto and M. Wakatani, *K- ϵ Model of Anomalous Transport in Resistive Interchange Turbulence ; Sep, 1992*
- NIFS-171 H. Sugama, M. Okamoto and M. Wakatani, *Vlasov Equation in the Stochastic Magnetic Field ; Sep. 1992*
- NIFS-172 N. Nakajima, M. Okamoto and M. Fujiwara, *Physical Mechanism of E_{ψ} -Driven Current in Asymmetric Toroidal Systems ; Sep.1992*
- NIFS-173 N. Nakajima, J. Todoroki and M. Okamoto, *On Relation between Hamada and Boozer Magnetic Coordinate System ; Sep. 1992*
- NIFS-174 K. Ichiguchi, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani, *Effects of Net Toroidal Current on Mercier Criterion in the Large Helical Device ; Sep. 1992*
- NIFS-175 S. -I. Itoh, K. Itoh and A. Fukuyama, *Modelling of ELMs and Dynamic Responses of the H-Mode ; Sep. 1992*

- NIFS-176 K. Itoh, S.-I. Itoh, A. Fukuyama, H. Sanuki, K. Ichiguchi and J. Todoroki, *Improved Models of β -Limit, Anomalous Transport and Radial Electric Field with Loss Cone Loss in Heliotron / Torsatron* ; Sep. 1992
- NIFS-177 N. Ohyabu, K. Yamazaki, I. Katanuma, H. Ji, T. Watanabe, K. Watanabe, H. Akao, K. Akaishi, T. Ono, H. Kaneko, T. Kawamura, Y. Kubota, N. Noda, A. Sagara, O. Motojima, M. Fujiwara and A. Iiyoshi, *Design Study of LHD Helical Divertor and High Temperature Divertor Plasma Operation* ; Sep. 1992
- NIFS-178 H. Sanuki, K. Itoh and S.-I. Itoh, *Selfconsistent Analysis of Radial Electric Field and Fast Ion Losses in CHS Torsatron / Heliotron* ; Sep. 1992
- NIFS-179 K. Toi, S. Morita, K. Kawahata, K. Ida, T. Watari, R. Kumazawa, A. Ando, Y. Oka, K. Ohkubo, Y. Hamada, K. Adati, R. Akiyama, S. Hidekuma, S. Hirokura, O. Kaneko, T. Kawamoto, Y. Kawasumi, M. Kojima, T. Kuroda, K. Masai, K. Narihara, Y. Ogawa, S. Okajima, M. Sakamoto, M. Sasao, K. Sato, K. N. Sato, T. Seki, F. Shimpō, S. Tanahashi, Y. Taniguchi, T. Tsuzuki, *New Features of L-H Transition in Limiter H-Modes of JIPP T-IIU* ; Sep. 1992
- NIFS-180 H. Momota, Y. Tomita, A. Ishida, Y. Kohzaki, M. Ohnishi, S. Ohi, Y. Nakao and M. Nishikawa, *D-³He Fueled FRC Reactor "Artemis-L"* ; Sep. 1992
- NIFS-181 T. Watari, R. Kumazawa, T. Seki, Y. Yasaka, A. Ando, Y. Oka, O. Kaneko, K. Adati, R. Akiyama, Y. Hamada, S. Hidekuma, S. Hirokura, K. Ida, K. Kawahata, T. Kawamoto, Y. Kawasumi, S. Kitagawa, M. Kojima, T. Kuroda, K. Masai, S. Morita, K. Narihara, Y. Ogawa, K. Ohkubo, S. Okajima, T. Ozaki, M. Sakamoto, M. Sasao, K. Sato, K. N. Sato, F. Shimpō, H. Takahashi, S. Tanahasi, Y. Taniguchi, K. Toi, T. Tsuzuki and M. Ono, *The New Features of Ion Bernstein Wave Heating in JIPP T-IIU Tokamak* ; Sep. 1992
- NIFS-182 K. Itoh, H. Sanuki and S.-I. Itoh, *Effect of Alpha Particles on Radial Electric Field Structure in Torsatron / Heliotron Reactor*; Sep. 1992
- NIFS-183 S. Morimoto, M. Sato, H. Yamada, H. Ji, S. Okamura, S. Kubo, O. Motojima, M. Murakami, T. C. Jernigan, T. S. Bigelow, A. C. England, R. S. Isler, J. F. Lyon, C. H. Ma, D. A. Rasmussen, C. R. Schaich, J. B. Wilgen and J. L. Yarber, *Long Pulse Discharges Sustained by Second Harmonic Electron Cyclotron Heating Using a 35GHz Gyrotron in the Advanced Toroidal Facility*; Sep. 1992
- NIFS-184 S. Okamura, K. Hanatani, K. Nishimura, R. Akiyama, T. Amano, H. Arimoto, M. Fujiwara, M. Hosokawa, K. Ida, H. Idei, H. Iguchi,

O. Kaneko, T. Kawamoto, S. Kubo, R. Kumazawa, K. Matsuoka, S. Morita, O. Motojima, T. Mutoh, N. Nakajima, N. Noda, M. Okamoto, T. Ozaki, A. Sagara, S. Sakakibara, H. Sanuki, T. Saki, T. Shoji, F. Shimbo, C. Takahashi, Y. Takeiri, Y. Takita, K. Toi, K. Tsumori, M. Ueda, T. Watari, H. Yamada and I. Yamada, *Heating Experiments Using Neutral Beams with Variable Injection Angle and ICRF Waves in CHS* ; Sep. 1992

NIFS-185 H. Yamada, S. Morita, K. Ida, S. Okamura, H. Iguchi, S. Sakakibara, K. Nishimura, R. Akiyama, H. Arimoto, M. Fujiwara, K. Hanatani, S. P. Hirshman, K. Ichiguchi, H. Idei, O. Kaneko, T. Kawamoto, S. Kubo, D. K. Lee, K. Matsuoka, O. Motojima, T. Ozaki, V. D. Pustovitov, A. Sagara, H. Sanuki, T. Shoji, C. Takahashi, Y. Takeiri, Y. Takita, S. Tanahashi, J. Todoroki, K. Toi, K. Tsumori, M. Ueda and I. Yamada, *MHD and Confinement Characteristics in the High- β Regime on the CHS Low-Aspect-Ratio Heliotron / Torsatron* ; Sep. 1992

NIFS-186 S. Morita, H. Yamada, H. Iguchi, K. Adati, R. Akiyama, H. Arimoto, M. Fujiwara, Y. Hamada, K. Ida, H. Idei, O. Kaneko, K. Kawahata, T. Kawamoto, S. Kubo, R. Kumazawa, K. Matsuoka, T. Morisaki, K. Nishimura, S. Okamura, T. Ozaki, T. Seki, M. Sakurai, S. Sakakibara, A. Sagara, C. Takahashi, Y. Takeiri, H. Takenaga, Y. Takita, K. Toi, K. Tsumori, K. Uchino, M. Ueda, T. Watari, I. Yamada, *A Role of Neutral Hydrogen in CHS Plasmas with Reheat and Collapse and Comparison with JIPP T-IIU Tokamak Plasmas* ; Sep. 1992

NIFS-187 K. Itoh, S.-I. Itoh, A. Fukuyama, M. Yagi and M. Azumi, *Model of the L-Mode Confinement in Tokamaks* ; Sep. 1992

NIFS-188 K. Itoh, A. Fukuyama and S.-I. Itoh, *Beta-Limiting Phenomena in High-Aspect-Ratio Toroidal Helical Plasmas*; Oct. 1992

NIFS-189 K. Itoh, S. -I. Itoh and A. Fukuyama, *Cross Field Ion Motion at Sawtooth Crash* ; Oct. 1992

NIFS-190 N. Noda, Y. Kubota, A. Sagara, N. Ohyabu, K. Akaishi, H. Ji, O. Motojima, M. Hashiba, I. Fujita, T. Hino, T. Yamashina, T. Matsuda, T. Sogabe, T. Matsumoto, K. Kuroda, S. Yamazaki, H. Ise, J. Adachi and T. Suzuki, *Design Study on Divertor Plates of Large Helical Device (LHD)* ; Oct. 1992

NIFS-191 Y. Kondoh, Y. Hosaka and K. Ishii, *Kernel Optimum Nearly-Analytical Discretization (KOND) Algorithm Applied to Parabolic and Hyperbolic Equations* : Oct. 1992