

# NATIONAL INSTITUTE FOR FUSION SCIENCE

## Entropy Production and Onsager Symmetry in Neoclassical Transport Processes of Toroidal Plasmas

H. Sugama and W. Horton

(Received - June 21, 1995)

NIFS-364

July 1995

### RESEARCH REPORT NIFS Series

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

# Entropy production and Onsager symmetry in neoclassical transport processes of toroidal plasmas

H. Sugama

*National Institute for Fusion Science, Nagoya 464-01, Japan*

and

W. Horton

*Institute for Fusion Studies, The University of Texas at Austin*

*Austin, Texas 78712*

Entropy production and Onsager symmetry in neoclassical transport processes of magnetically confined plasmas are studied in detail for general toroidal systems including nonaxisymmetric configurations. We find that the flux surface average of the entropy production defined from the linearized collision operator and the gyroangle-averaged distribution function coincides with the sum of the inner products of the thermodynamic forces and the conjugate fluxes consisting of the Pfirsch-Schlüter, banana-plateau, nonaxisymmetric parts of the neoclassical radial fluxes and the parallel current. We prove from the self-adjointness of the linearized collision operator that the Onsager symmetry is robustly valid for the neoclassical transport equations in the cases of general toroidal plasmas consisting of electrons and multi-species ions with arbitrary collision frequencies. It is shown that the Onsager symmetry holds whether or not the ambipolarity condition is used to reduce the number of the conjugate pairs of the transport fluxes and the thermodynamic forces. We also derive the full transport coefficients for the banana-plateau and nonaxisymmetric parts, separately, and investigate their symmetry properties. The nonaxisymmetric transport equations are obtained for arbitrary collision frequencies in the Pfirsch-Schlüter and plateau regimes, and it is directly confirmed that the total banana-plateau and nonaxisymmetric transport equations satisfy the Onsager symmetry.

Keywords: neoclassical transport, entropy production, Onsager symmetry

# I. INTRODUCTION

Transport processes caused by binary Coulomb collisions between charged particles in toroidal magnetic configurations are described by the neoclassical transport theory.<sup>1-3</sup> Particle and heat transport fluxes observed in most fusion devices exceed the predictions of the neoclassical transport theory and thus are called anomalous transport.<sup>4</sup> The anomalous transport is considered to result from the turbulent fluctuations driven by various plasma instabilities, which are not taken into account by the neoclassical theory. However, the neoclassical theory is regarded as a standard model with a well-established framework, which gives lower limits of the transport fluxes in quiescent plasmas close to thermal equilibria, and it also has practical use for predicting the parallel transport such as a bootstrap current with more accuracy than the predictions for the radial transport. The characteristics of the neoclassical theory lie in the inclusion of the effects of the global magnetic field geometry due to the long mean free path of the hot particles. In less collisional regimes such as plateau and banana regimes, the processes in deriving the neoclassical transport equations, which relate the neoclassical transport fluxes to the thermodynamic forces, are more complicated than the derivation of the classical transport equations in more collisional regimes. Since the neoclassical transport coefficients contain the parameters relating to both the collisionality and the magnetic geometry, it is less trivial than in the case of the classical transport to show the validity of the Onsager symmetry<sup>5</sup> for the neoclassical transport. In the cases of axisymmetric systems, the analytical expressions of the full neoclassical transport coefficients for all collisionality regimes have been obtained, and it is well-known that the Onsager symmetry holds for the axisymmetric neoclassical transport matrix.

The neoclassical transport equations for nonaxisymmetric systems<sup>6-11</sup> are more complicated due to the geometrical complexities, and they involve the nonambipolar parts. The absence of symmetry may also cause the breakup of magnetic surfaces into islands and ergodic regions,<sup>12</sup> although such problems are beyond the scope of this work. Here we assume the existence of toroidal nested magnetic surfaces as in other previous works.<sup>6-11</sup> Balescu and Fantechi derive the full neoclassical transport coefficients for the nonaxisymmetric plasma in the plateau regime, and claim that the Onsager symmetry partly breaks down in that case.<sup>11</sup> Contrary to Ref.11, we show in the present work that the Onsager symmetry is robustly valid for the neoclassical transport equations in the cases of general toroidal plasmas consisting of electrons and multi-species ions with arbitrary collision frequencies.

Concerning the Onsager relation, it is important to discuss the entropy production resulting from the transport processes since the Onsager relation, if it holds, is satisfied only by the transport matrix connecting the conjugate pairs of transport fluxes and thermodynamic forces, which should be specified through the entropy production.<sup>5</sup> According to the terminology in Ref.3, we expect that the kinetic form of the entropy production defined from the collision operator should coincide with its thermodynamic form in which the entropy production is expressed as the sum of the products of the conjugate pairs of the fluxes and forces.

In Chap.17 of Ref.3, Balescu presented detailed analyses on the kinetic and thermodynamic forms of the entropy production in the classical and neoclassical transport processes for the axisymmetric case. Using the Hermitian moment representation, he confirmed that the kinetic form of the entropy production includes the thermodynamic form given by the products of the thermodynamic forces and their conjugate classical and neoclassical Pfirsch-Schlüter fluxes, although he did not identify the thermodynamic form corresponding to the the neoclassical banana-plateau transport. In this work, before the proof of the Onsager symmetry, we show for general toroidal geometry the complete coincidence between the kinetic form of the entropy production and its full thermodynamic form including all contributions from the classical and neoclassical (Pfirsch-Schlüter, banana-plateau, nonaxisymmetric) fluxes. The main difference between our treatment and that in Ref.3 is that we use directly the distribution function and the drift kinetic equations instead of the Hermitian moment expansion.

The proof of the Onsager symmetry for the neoclassical transport equations in general toroidal configurations is given in the similar manner to that in the Appendix of Ref.2. The proof uses the self-adjointness of the linearized collision operator and the formal solution of the linearized drift kinetic equation although neither axisymmetry of the magnetic configuration nor any condition for collisionality is not required. We also derive the full neoclassical transport coefficients in the nonaxisymmetric system for collision frequencies in the Pfirsch-Schlüter and plateau regimes, from which the Onsager symmetry of the full neoclassical transport matrix is directly confirmed.

In the axisymmetric configurations, the ambipolarity is automatically satisfied by the neoclassical transport and the radial electric field does not affect the transport fluxes. On the other hand, in the nonaxisymmetric configurations, the radial electric field is determined through the particle transport equations if the ambipolarity condition is imposed. In both cases with and without the ambipolarity condition, we give the neoclassical transport equations and check their Onsager symmetry.

This work is organized as follows. In Sec.II, the entropy production defined from the collision operator is divided into the two parts, which are derived from the gyroangle-averaged and gyroangle-dependent parts of the distribution function. The entropy production from the gyroangle-dependent distribution function is given by the sum of the inner products of the classical radial particle and heat fluxes and the radial gradient thermodynamic forces. The transport matrix relating these classical fluxes and forces is shown to satisfy the Onsager symmetry. We find that the entropy production from the gyroangle-averaged distribution function is written as the sum of the inner products of the thermodynamic forces and the corresponding conjugate fluxes which consist of the Pfirsch-Schlüter, banana-plateau, non-axisymmetric parts of the neoclassical radial fluxes and the parallel current. In Sec.III, using the formal solution of the linearized drift kinetic equation and the self-adjointness of the linearized collision operator, we prove that the Onsager symmetry is satisfied by the neoclassical transport equations for arbitrary collision frequencies in general toroidal systems including nonaxisymmetric cases. The effects of the ambipolarity on the neoclassical transport coefficients are examined for both axisymmetric and nonaxisymmetric cases to show the robust validity of the Onsager symmetry independent of the use of the ambipolar-

ity condition. In Sec.IV, the full transport coefficients are derived for the banana-plateau and nonaxisymmetric parts, separately, and their symmetry properties are investigated. We derive the nonaxisymmetric transport coefficients for arbitrary collision frequencies in the Pfirsch-Schlüter and plateau regimes, and directly confirm that the total banana-plateau and nonaxisymmetric transport equations satisfy the Onsager symmetry. Finally, conclusions and discussions are given in Sec.V. There, we discuss the reason why our two main results, i.e., the complete correspondence between the kinetic and thermodynamic forms of the entropy production, and the Onsager symmetry in the nonaxisymmetric case, were not confirmed in Ref.3 and in Ref.11, respectively.

## II. ENTROPY PRODUCTION IN CLASSICAL AND NEO-CLASSICAL TRANSPORT PROCESSES

Here, we show that the thermodynamic form of the entropy production is equivalent to its kinetic form defined from the collision operator for the classical and neoclassical transport in the general case of magnetically confined plasma with arbitrary toroidal geometry. For that purpose, we first describe several properties of the collision operator which is denoted for species  $a$  by

$$C_a = \sum_b C_{ab}(f_a, f_b) \quad (1)$$

where  $f_a$  ( $f_b$ ) is the distribution function of the species  $a$  ( $b$ ) and  $C_{ab}$  represents the contribution from the collision between the particles  $a$  and  $b$ .

The collision operator conserves the particles' number, momentum, and kinetic energy, which is written as

$$\begin{aligned} \int d^3v C_{ab} &= 0 \\ \int d^3v m_a \mathbf{v} C_{ab} + \int d^3v m_b \mathbf{v} C_{ba} &= 0 \\ \int d^3v \frac{1}{2} m_a v^2 C_{ab} + \int d^3v \frac{1}{2} m_b v^2 C_{ba} &= 0. \end{aligned} \quad (2)$$

Furthermore, the collision operator is invariant under arbitrary translational and rotational transform of the velocity variable  $\mathbf{v}$  of distribution functions, which is expressed by

$$\mathcal{T}C_{ab}(f_a, f_b) = C_{ab}(\mathcal{T}f_a, \mathcal{T}f_b), \quad \mathcal{R}C_{ab}(f_a, f_b) = C_{ab}(\mathcal{R}f_a, \mathcal{R}f_b) \quad (3)$$

where  $\mathcal{T}f$  and  $\mathcal{R}f$  denote functions  $f$  with arbitrary translational and rotational transform of the velocity variable  $\mathbf{v}$ , respectively.

The entropy production for the species  $a$  is defined from the collision operator by

$$\dot{S}_a \equiv \sum_b \dot{S}_{ab} \equiv - \sum_b \int d^3v (\ln f_a) C_{ab}(f_a, f_b). \quad (4)$$

The second law of the thermodynamics or the positive definiteness of the entropy production is given by

$$\dot{S}_{ab} + \dot{S}_{ba} \geq 0, \quad \sum_a \dot{S}_a \geq 0 \quad (5)$$

where the total entropy production  $\sum_a \dot{S}_a$  vanishes if and only if the distribution functions for all species are the Maxwellian with the same temperature and the same mean velocity. However, even if two particle species with much different masses (such as for electrons and ions) have the Maxwellian distributions with different temperatures, the entropy production due to their collisions is so small that we hereafter neglect it based on the small mass ratio ordering.

For magnetically confined plasmas, the lowest-order distribution function for the species  $a$  with respect to the drift ordering is given by the Maxwellian with no mean velocity as

$$f_{a0} = \pi^{-3/2} n_a v_{T_a}^{-3} \exp(-x_a^2) \equiv f_{aM} \quad (6)$$

where  $v_{T_a}^2 \equiv 2T_a/m_a$  and  $x_a \equiv v/v_{T_a}$  are defined from the temperature  $T_a$ . The distribution functions and the collision operator are perturbatively expanded in the drift ordering parameter  $\delta \equiv \rho/L$  ( $\rho$ : the thermal gyroradius,  $L$ : the equilibrium scale length) as

$$\begin{aligned} f_a &= f_{a0} + f_{a1} + \mathcal{O}(\delta^2) \\ C_{ab} &= C_{ab}(f_{a0}, f_{b0}) + C_{ab}^L(f_{a1}, f_{b1}) + \mathcal{O}(\delta^2) \end{aligned} \quad (7)$$

where the linearized collision operator  $C_{ab}^L$  is defined by

$$C_{ab}^L(f_{a1}, f_{b1}) \equiv C_{ab}(f_{a1}, f_{b0}) + C_{ab}(f_{a0}, f_{b1}). \quad (8)$$

The linearized collision operator also has the conservation and symmetry properties, which are expressed by Eqs.(2) and (3) with  $C_{ab}$  replaced by  $C_{ab}^L$ . The self-adjointness of the linearized collision operator<sup>13</sup> is written as

$$\begin{aligned} & T_a \int d^3v \frac{g_{a1}}{f_{a0}} C_{ab}^L(h_{a1}, h_{b1}) + T_b \int d^3v \frac{g_{b1}}{f_{b0}} C_{ba}^L(h_{b1}, h_{a1}) \\ &= T_a \int d^3v \frac{h_{a1}}{f_{a0}} C_{ab}^L(g_{a1}, g_{b1}) + T_b \int d^3v \frac{h_{b1}}{f_{b0}} C_{ba}^L(g_{b1}, g_{a1}) \end{aligned} \quad (9)$$

which is exactly valid for  $T_a = T_b$  and is approximately satisfied for  $T_a \neq T_b$  when  $m_a/m_b \ll 1$  or  $m_b/m_a \ll 1$ . Concerning the positive definiteness of the entropy production described in Eq.(5), we find the positive definiteness of the quadratic form associated with the linearized collision operator as

$$-T_a \int d^3v \frac{g_{a1}}{f_{a0}} C_{ab}^L(g_{a1}, g_{b1}) - T_b \int d^3v \frac{g_{b1}}{f_{b0}} C_{ba}^L(g_{b1}, g_{a1}) \geq 0 \quad (10)$$

which is valid for  $T_a \neq T_b$  to the lowest order of the small mass ratio  $m_a/m_b \ll 1$  (or  $m_b/m_a \ll 1$ ) as Eq.(9).

Using in Eq.(4)

$$\ln f_a = \ln f_{a0} + \frac{f_{a1}}{f_{a0}} + \mathcal{O}(\delta^2)$$

the entropy production  $\dot{S}_a$  up to  $\mathcal{O}(\delta^2)$  is given by

$$\sigma_a \equiv \sum_b \sigma_{ab} \equiv - \sum_b \int d^3v \frac{f_{a1}}{f_{a0}} C_{ab}^L(f_{a1}, f_{b1}). \quad (11)$$

Instead of Eq.(5), we have from Eq.(10)

$$T_a \sigma_{ab} + T_b \sigma_{ba} \geq 0, \quad \sum_a T_a \sigma_a \geq 0. \quad (12)$$

Now, let us divide the first-order distribution function  $f_{a1}$  into the gyroangle-averaged part  $\bar{f}_{a1}$  and the gyroangle-dependent part  $\tilde{f}_{a1}$  as

$$f_{a1} = \bar{f}_{a1} + \tilde{f}_{a1}. \quad (13)$$

Due to the rotational symmetry of the collision operator in the velocity space shown in Eq.(3), the entropy production  $\sigma_a$  separates into into the corresponding parts as

$$\sigma_a = \bar{\sigma}_a + \tilde{\sigma}_a \quad (14)$$

where

$$\bar{\sigma}_a = - \sum_b \int d^3v \frac{\bar{f}_{a1}}{f_{a0}} C_{ab}^L(\bar{f}_{a1}, \bar{f}_{b1}) \quad (15)$$

$$\tilde{\sigma}_a = - \sum_b \int d^3v \frac{\tilde{f}_{a1}}{f_{a0}} C_{ab}^L(\tilde{f}_{a1}, \tilde{f}_{b1}). \quad (16)$$

First, we consider the entropy production  $\tilde{\sigma}_a$  due to the gyroangle-dependent part of the distribution function. The gyroangle-dependent part  $\tilde{f}_{a1}$  is given from the lowest-order distribution function  $f_{a0}$  by

$$\begin{aligned} \tilde{f}_{a1} &= \frac{\mathbf{v} \times \mathbf{n}}{\Omega_a} \cdot \nabla f_{a0} = -f_{a0} \frac{\mathbf{v} \times \mathbf{n} \cdot \nabla V}{\Omega_a T_a} \left[ X_{a1} + X_{a2} \left( x_a^2 - \frac{5}{2} \right) \right] \\ &= \frac{m_a}{T_a} \mathbf{v} \cdot \left[ \mathbf{u}_{\perp a} + \frac{2}{5} \frac{\mathbf{q}_{\perp a}}{p_a} \left( x_a^2 - \frac{5}{2} \right) \right] f_{a0} \end{aligned} \quad (17)$$

where  $\mathbf{n} = \mathbf{B}/B$  is the unit vector along the magnetic field  $\mathbf{B}$  and  $\Omega_a = e_a B/m_a c$  is the gyrofrequency of the particle with the mass  $m_a$  and the charge  $e_a$ . In Eq.(17),  $f_{a0}$  is regarded as a function of  $(V, E, \mu)$  ( $V$ : the volume inside the flux surface,  $E \equiv \frac{1}{2} m_a v^2 + e_a \Phi$ : the particle's energy, and  $\mu \equiv m_a v_{\perp}^2 / 2B$ : the magnetic moment) and we have used

$$\frac{\partial \ln f_{a0}}{\partial V} = -\frac{1}{T_a} \left[ X_{a1} + X_{a2} \left( x_a^2 - \frac{5}{2} \right) \right] \quad (18)$$

where the thermodynamic forces  $X_{a1}$  and  $X_{a2}$  are defined from the radial gradients of the pressure  $p_a$ , the electrostatic potential  $\Phi$ , and the temperature  $T_a$  as

$$X_{a1} \equiv -\frac{p'_a}{n_a} - e_a \Phi', \quad X_{a2} \equiv -T'_a \quad (' \equiv \partial/\partial V) \quad (19)$$

in terms of which the perpendicular components of the fluid velocity and the heat flow are

$$\begin{aligned} \mathbf{u}_{\perp a} &\equiv \frac{1}{n_a} \int d^3v \tilde{f}_{a1} \mathbf{v}_{\perp} = \frac{cX_{a1}}{e_a B} \nabla V \times \mathbf{n} \\ \frac{\mathbf{q}_{\perp a}}{p_a} &\equiv \frac{1}{n_a} \int d^3v \tilde{f}_{a1} \mathbf{v}_{\perp} \left(x_a^2 - \frac{5}{2}\right) = \frac{5}{2} \frac{cX_{a2}}{e_a B} \nabla V \times \mathbf{n}. \end{aligned} \quad (20)$$

Substituting Eq.(17) into Eq.(16), we obtain

$$T_a \tilde{\sigma}_a = J_{a1}^{\text{cl}} X_{a1} + J_{a2}^{\text{cl}} X_{a2} \quad (21)$$

where

$$J_{a1}^{\text{cl}} \equiv \mathbf{\Gamma}_a^{\text{cl}} \cdot \nabla V, \quad J_{a2}^{\text{cl}} \equiv \frac{1}{T_a} \mathbf{q}_a^{\text{cl}} \cdot \nabla V \quad (22)$$

are the radial components of the classical particle and heat fluxes defined by

$$\begin{aligned} \mathbf{\Gamma}_a^{\text{cl}} &= \frac{c}{e_a B} \mathbf{F}_{a1} \times \mathbf{n} \\ \frac{1}{T_a} \mathbf{q}_a^{\text{cl}} &= \frac{c}{e_a B} \mathbf{F}_{a2} \times \mathbf{n}. \end{aligned} \quad (23)$$

Here the friction forces  $\mathbf{F}_{a1}$  and  $\mathbf{F}_{a2}$  are given by

$$\begin{aligned} \mathbf{F}_{a1} &= \sum_b \int d^3v m_a \mathbf{v} C_{ab}^L(f_{a1}, f_{b1}) \\ \mathbf{F}_{a2} &= \sum_b \int d^3v m_a \mathbf{v} \left(x_a^2 - \frac{5}{2}\right) C_{ab}^L(f_{a1}, f_{b1}). \end{aligned} \quad (24)$$

It should be noted that, from the rotational symmetry of the collision operator, that  $\bar{f}_{a1}$  does not contribute to the perpendicular friction forces  $\mathbf{F}_{\perp aj}$  while  $\tilde{f}_{a1}$  does not contribute to the parallel friction forces  $\mathbf{F}_{\parallel aj}$ . Equation (21) shows that the entropy production  $\tilde{\sigma}_a$  defined from the gyroangle dependent part of the distribution function is caused by the classical particle and heat transport, and that the classical fluxes  $J_{a1}^{\text{cl}}$  and  $J_{a2}^{\text{cl}}$  are conjugate to the thermodynamic forces  $X_{a1}$  and  $X_{a2}$ , respectively. The momentum conservation due to the collision operator gives

$$\sum_a \mathbf{F}_{a1} = 0 \quad (25)$$



which in turn causes the classical particle fluxes to satisfy the ambipolarity as

$$\sum_a e_a \Gamma_a^{\text{cl}} = 0. \quad (26)$$

We have the relations between the perpendicular friction forces and flows from Eqs.(17) and (24) as

$$\begin{bmatrix} \mathbf{F}_{\perp a1} \\ -\mathbf{F}_{\perp a2} \end{bmatrix} = \sum_b \begin{bmatrix} l_{11}^{ab} & l_{12}^{ab} \\ l_{21}^{ab} & l_{22}^{ab} \end{bmatrix} \begin{bmatrix} \mathbf{u}_{\perp b} \\ -\frac{2}{5p_b} \mathbf{q}_{\perp b} \end{bmatrix} \quad (27)$$

where the coefficients  $l_{jk}^{ab}$  are the same ones defined in Ref.2 and are given by

$$\begin{aligned} l_{jk}^{ab} = & \delta_{ab} \frac{m_a^2}{T_a} \sum_{a'} \int d^3v v_{\parallel} L_{j-1}^{(3/2)}(x_a^2) C_{aa'} [v_{\parallel} L_{k-1}^{(3/2)}(x_a^2) f_{a0}, f_{a'0}] \\ & + \frac{m_a m_b}{T_b} \int d^3v v_{\parallel} L_{j-1}^{(3/2)}(x_a^2) C_{ab} [f_{a0}, v_{\parallel} L_{k-1}^{(3/2)}(x_b^2) f_{b0}]. \end{aligned} \quad (28)$$

Here,  $L_0^{(3/2)}(x^2) = 1$ ,  $L_1^{(3/2)}(x^2) = \frac{5}{2} - x^2$ ,  $\dots$  are the Laguerre polynomials of order  $\frac{3}{2}$ . In Eqs.(27) and (28), the rotational symmetry of the linearized collision operator is used. From the self-adjointness of the linearized collision operator shown in Eq.(9), the coefficients  $l_{jk}^{ab}$  have the following symmetry :

$$l_{jk}^{ab} = l_{kj}^{ba}. \quad (29)$$

The momentum conservation property described in Eq.(25) imposes another constraint on the coefficients  $l_{jk}^{ab}$  :

$$\sum_a l_{1k}^{ab} = 0. \quad (30)$$

From Eq.(21) and the ambipolarity condition  $\sum_a e_a J_{a1}^{\text{cl}} = 0$  given by Eq.(26), we obtain

$$\begin{aligned} \sum_a T_a \tilde{\sigma}_a &= \sum_a [J_{a1}^{\text{cl}} X_{a1} + J_{a2}^{\text{cl}} X_{a2}] \\ &= \sum_{a \neq I} J_{a1}^{\text{cl}} X_{a1}^* + \sum_a J_{a2}^{\text{cl}} X_{a2} \end{aligned} \quad (31)$$

where  $X_{a1}^*$  ( $a \neq I$ ) is defined by

$$X_{a1}^* = X_{a1} - \frac{e_a}{e_I} X_{I1} = -\frac{p'_a}{n_a} + \frac{e_a p'_I}{e_I n_I} \quad (a \neq I). \quad (32)$$

Here we have chosen a certain particle species denoted by  $I$ . We hereafter regard  $I$  as the ion species with the smallest particle number density. If a plasma consists of electrons and a single ion species  $i$ , we take  $I = i$ .

Equation (31) shows that the number of the conjugated pairs of the classical fluxes and thermodynamic forces is reduced by employing the new pairs  $(J_{a1}, X_{a1}^*)_{a \neq I}$ ,  $(J_{a2}, X_{a2})$  instead of  $(J_{a1}, X_{a1})$ ,  $(J_{a2}, X_{a2})$ . We also find that the radial electric field does not appear in the

new set of the thermodynamic forces. The transport equations which relate the classical fluxes ( $J_{a1}, J_{a2}$ ) to the thermodynamic forces ( $X_{a1}, X_{a2}$ ) are obtained from Eqs.(20), (22), (23), and (27) as

$$\begin{bmatrix} J_{a1}^{\text{cl}} \\ J_{a2}^{\text{cl}} \end{bmatrix} = \sum_b \begin{bmatrix} (L^{\text{cl}})_{11}^{ab} & (L^{\text{cl}})_{12}^{ab} \\ (L^{\text{cl}})_{21}^{ab} & (L^{\text{cl}})_{22}^{ab} \end{bmatrix} \begin{bmatrix} X_{b1} \\ X_{b2} \end{bmatrix} \quad (33)$$

where the classical transport coefficients  $(L^{\text{cl}})_{jk}^{ab}$  are given by

$$(L^{\text{cl}})_{jk}^{ab} = (-1)^{j+k-1} \frac{c^2 |\nabla V|^2}{e_a e_b B^2} l_{jk}^{ab}, \quad (j, k = 1, 2). \quad (34)$$

From Eq.(29), we have the Onsager symmetry for the classical transport as

$$(L^{\text{cl}})_{jk}^{ab} = (L^{\text{cl}})_{kj}^{ba}, \quad (j, k = 1, 2). \quad (35)$$

Equation (30) yields

$$\sum_a e_a (L^{\text{cl}})_{1k}^{ab} = 0, \quad (k = 1, 2). \quad (36)$$

We easily find from Eqs.(32), (33), (35), and (36) that the classical transport equations for the pairs  $(J_{a1}^{\text{cl}}, X_{a1}^*)_{a \neq I}$ ,  $(J_{a2}^{\text{cl}}, X_{a2})$  are given by

$$\begin{aligned} J_{a1(a \neq I)}^{\text{cl}} &= \sum_{b \neq I} (L^{\text{cl}})_{11}^{ab} X_{b1}^* + \sum_b (L^{\text{cl}})_{12}^{ab} X_{b2} \\ J_{a2}^{\text{cl}} &= \sum_{b \neq I} (L^{\text{cl}})_{21}^{ab} X_{b1}^* + \sum_b (L^{\text{cl}})_{22}^{ab} X_{b2} \end{aligned} \quad (37)$$

which shows that the transport coefficients  $(L^{\text{cl}})_{jk}^{ab}$  are the same as in Eq.(33) except for the limitation  $(a, j), (b, k) \neq (I, 1)$  and that the Onsager symmetry is valid for both of the conjugate pairs. We should note that the ambipolarity condition (36) reduces the number of the thermodynamic forces required for determining the classical fluxes by one, and that the radial electric field does not enter the reduced set of the thermodynamic forces  $(X_{a1(a \neq I)}^*, X_{a2})$ .

Next, let us consider the entropy production  $\bar{\sigma}_a$  due to the first-order gyroangle-averaged distribution function  $\bar{f}_{a1}$ , which satisfies the drift kinetic equation<sup>14,15</sup> :

$$v_{\parallel} \mathbf{n} \cdot \nabla \bar{f}_{a1} + \mathbf{v}_{da} \cdot \nabla f_{a0} - \frac{e_a}{T_a} v_{\parallel} E_{\parallel} f_{a0} = C_a^L(\bar{f}_{a1}) \quad (38)$$

where  $\mathbf{v}_{da}$  is the sum of the  $\mathbf{E} \times \mathbf{B}$ ,  $\nabla B$  and curvature drift velocities, and  $C_a^L(\bar{f}_{a1}) \equiv \sum_b C_{ab}^L(\bar{f}_{a1}, \bar{f}_{b1})$ . We obtain from Eqs.(15) and (38)

$$\begin{aligned} \bar{\sigma}_a &\equiv - \int d^3 v \frac{\bar{f}_{a1}}{f_{a0}} C_a^L(\bar{f}_{a1}) \\ &= - \nabla \cdot \left[ \mathbf{n} \int d^3 v v_{\parallel} \frac{(\bar{f}_{a1})^2}{2 f_{a0}} \right] - \int d^3 v \bar{f}_{a1} \mathbf{v}_{da} \cdot \nabla V \frac{\partial \ln f_{a0}}{\partial V} + \frac{1}{T_a} n_a e_a u_{\parallel a} E_{\parallel} \end{aligned} \quad (39)$$

where the parallel flow velocity  $u_{\parallel a}$  is defined by  $n_a u_{\parallel a} \equiv \int d^3 v \bar{f}_{a1} v_{\parallel}$ . Here, the flux surface average of the first term in the right-hand side vanishes. In taking the flux surface average of the second term, we use the following two equations for the radial particle and heat fluxes :

$$\begin{aligned}
& \left\langle \int d^3 v \bar{f}_{a1} \mathbf{v}_{da} \cdot \nabla V \right\rangle = \left\langle \frac{\nabla V}{m_a \Omega_a} \cdot \mathbf{n} \times (\nabla p_{a1} + \nabla \cdot \boldsymbol{\pi}_a) \right\rangle \\
& = -\frac{c}{e_a B^\theta} \left\langle \frac{B_\zeta}{B} \mathbf{n} \cdot (\nabla p_{a1} + \nabla \cdot \boldsymbol{\pi}_a) \right\rangle + \frac{c}{e_a B^\theta B_\zeta} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\
& = -\frac{c}{e_a B^\theta} \frac{\langle B_\zeta \rangle}{\langle B^2 \rangle} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle + \frac{c}{e_a B^\theta B_\zeta} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\
& \quad - \frac{c}{e_a B^\theta} \left\langle \frac{1}{B} \underbrace{\mathbf{n} \cdot (\nabla p_{a1} + \nabla \cdot \boldsymbol{\pi}_a)}_{(F_{\parallel a1} + n_a e_a E_{\parallel})} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \\
& = J_{a1}^{\text{ncl}} - \frac{n_a c}{B^\theta} \left\langle \frac{E_{\parallel}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \tag{40}
\end{aligned}$$

$$\begin{aligned}
& \left\langle \int d^3 v \bar{f}_{a1} \left( x_a^2 - \frac{5}{2} \right) \mathbf{v}_{da} \cdot \nabla V \right\rangle = \left\langle \frac{\nabla V}{m_a \Omega_a} \cdot \mathbf{n} \times (\nabla \theta_{a1} + \nabla \cdot \boldsymbol{\Theta}_a) \right\rangle \\
& = -\frac{c}{e_a B^\theta} \frac{\langle B_\zeta \rangle}{\langle B^2 \rangle} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle + \frac{c}{e_a B^\theta B_\zeta} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \\
& \quad - \frac{c}{e_a B^\theta} \left\langle \frac{1}{B} \underbrace{\mathbf{n} \cdot (\nabla \theta_{a1} + \nabla \cdot \boldsymbol{\Theta}_a)}_{F_{\parallel a2}} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \\
& = J_{a2}^{\text{ncl}} \tag{41}
\end{aligned}$$

where the Hamada coordinates  $(V, \theta, \zeta)$  are employed to represent the poloidal and toroidal components of the magnetic fields (see Appendix A) and Eq.(A2) is used. Here, we have used the definitions :  $p_{a1} \equiv \int d^3 v \frac{1}{3} m_a v^2 \bar{f}_{a1}$ ,  $\theta_{a1} \equiv \int d^3 v \frac{1}{3} m_a v^2 (x_a^2 - \frac{5}{2}) \bar{f}_{a1}$ ,  $\boldsymbol{\pi}_a \equiv (p_{\parallel a} - p_{\perp a})(\mathbf{nn} - \frac{1}{3}\mathbf{l}) \equiv \int d^3 v m_a (v_{\parallel}^2 - \frac{1}{2}v_{\perp}^2) \bar{f}_{a1} (\mathbf{nn} - \frac{1}{3}\mathbf{l})$ ,  $\boldsymbol{\Theta}_a \equiv (\Theta_{\parallel a} - \Theta_{\perp a})(\mathbf{nn} - \frac{1}{3}\mathbf{l}) \equiv \int d^3 v m_a (v_{\parallel}^2 - \frac{1}{2}v_{\perp}^2) (x_a^2 - \frac{5}{2}) \bar{f}_{a1} (\mathbf{nn} - \frac{1}{3}\mathbf{l})$ . The neoclassical particle and heat fluxes are given by

$$\begin{aligned}
J_{a1}^{\text{ncl}} & \equiv \langle \boldsymbol{\Gamma}_a \cdot \nabla V \rangle^{\text{ncl}} \equiv J_{a1}^{\text{PS}} + J_{a1}^{\text{bp}} + J_{a1}^{\text{na}} \\
J_{a2}^{\text{ncl}} & \equiv \frac{1}{T_a} \langle \mathbf{q}_a \cdot \nabla V \rangle^{\text{ncl}} \equiv J_{a2}^{\text{PS}} + J_{a2}^{\text{bp}} + J_{a2}^{\text{na}} \tag{42}
\end{aligned}$$

which consist of the Pfirsch-Schlüter ( $J_{aj}^{\text{PS}}$ ), the banana-plateau ( $J_{aj}^{\text{bp}}$ ), and nonaxisymmetric

( $J_{aj}^{\text{na}}$ ) parts defined by

$$\begin{aligned}
J_{a1}^{\text{PS}} &\equiv \langle \mathbf{\Gamma}_a \cdot \nabla V \rangle^{\text{PS}} \equiv -\frac{c}{e_a B^\theta} \left\langle \frac{F_{\parallel a1}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \\
J_{a1}^{\text{bp}} &\equiv \langle \mathbf{\Gamma}_a \cdot \nabla V \rangle^{\text{bp}} \equiv -\frac{c}{e_a B^\theta} \frac{\langle B_\zeta \rangle}{\langle B^2 \rangle} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\
J_{a1}^{\text{na}} &\equiv \langle \mathbf{\Gamma}_a \cdot \nabla V \rangle^{\text{na}} \equiv \frac{c}{e_a B^\theta B_\zeta} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\
J_{a2}^{\text{PS}} &\equiv \frac{1}{T_a} \langle \mathbf{q}_a \cdot \nabla V \rangle^{\text{PS}} \equiv -\frac{c}{e_a B^\theta} \left\langle \frac{F_{\parallel a2}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \\
J_{a2}^{\text{bp}} &\equiv \frac{1}{T_a} \langle \mathbf{q}_a \cdot \nabla V \rangle^{\text{bp}} \equiv -\frac{c}{e_a B^\theta} \frac{\langle B_\zeta \rangle}{\langle B^2 \rangle} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \\
J_{a2}^{\text{na}} &\equiv \frac{1}{T_a} \langle \mathbf{q}_a \cdot \nabla V \rangle^{\text{na}} \equiv \frac{c}{e_a B^\theta B_\zeta} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle
\end{aligned} \tag{43}$$

Then, we find that the flux surface average of the second term in the right-hand side of Eq.(39) is given by the products of the neoclassical radial fluxes and the thermodynamic forces as

$$-\left\langle \int d^3v \bar{f}_{a1} \mathbf{v}_{da} \cdot \nabla V \frac{\partial \ln f_{a0}}{\partial V} \right\rangle = \frac{1}{T_a} \left[ J_{a1}^{\text{ncl}} - \frac{n_a c}{B^\theta} \left\langle \frac{E_{\parallel}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \right] X_{a1} + \frac{1}{T_a} J_{a2}^{\text{ncl}} X_{a2}. \tag{44}$$

The flux surface average of the third term in the right-hand side of Eq.(39) is given by

$$\frac{n_a e_a \langle u_{\parallel a} E_{\parallel} \rangle}{T_a} = \frac{n_a e_a \langle B u_{\parallel a} \rangle \langle B E_{\parallel} \rangle}{T_a \langle B^2 \rangle} + \frac{n_a c}{T_a B^\theta} \left\langle \frac{E_{\parallel}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle X_{a1} \tag{45}$$

where Eq.(A7) is used. Then, we finally obtain from Eqs.(39), (44), and (45) the thermodynamic form of the flux-surface-averaged entropy production  $\langle \bar{\sigma}_a \rangle$  as

$$T_a \langle \bar{\sigma}_a \rangle = J_{a1}^{\text{ncl}} X_{a1} + J_{a2}^{\text{ncl}} X_{a2} + J_{a3} X_{a3} \tag{46}$$

where the parallel flux  $J_{a3}$  and the parallel force  $X_{a3}$  are defined by

$$J_{a3} \equiv \frac{n_a \langle B u_{\parallel a} \rangle}{\langle B^2 \rangle^{1/2}}, \quad X_{a3} \equiv e_a \frac{\langle B E_{\parallel} \rangle}{\langle B^2 \rangle^{1/2}}. \tag{47}$$

Taking the species summation of Eq.(46), we have

$$\sum_a T_a \langle \bar{\sigma}_a \rangle = \sum_a \left( J_{a1}^{\text{ncl}} X_{a1} + J_{a2}^{\text{ncl}} X_{a2} \right) + J_E X_E \tag{48}$$

where  $J_E$  and  $X_E$  are defined from the total parallel current  $J_{\parallel}$  and the parallel electric field  $E_{\parallel}$  as

$$J_E \equiv \frac{\langle BJ_{\parallel} \rangle}{\langle B^2 \rangle^{1/2}} \equiv \sum_a e_a J_{a3}, \quad X_E \equiv \frac{\langle BE_{\parallel} \rangle}{\langle B^2 \rangle^{1/2}}. \quad (49)$$

Thus the flux surface average of the entropy production due to the gyroangle-averaged distribution functions is given in the thermodynamic form, in which the neoclassical radial fluxes  $J_{a1}^{\text{nc1}}$ ,  $J_{a2}^{\text{nc1}}$  and the parallel current  $J_E$  are conjugate to the radial gradient forces  $X_{a1}$ ,  $X_{a2}$  and the parallel electric field  $X_E$ , respectively. It should be noted that the neoclassical thermodynamic form of the entropy production can be obtained only through the magnetic surface average as in Eq.(48), which is a remarkable contrast to the classical thermodynamic form (21) defined locally in the configuration space.

Now, let us consider the ambipolarity condition for the neoclassical particle fluxes. From the momentum conservation (25) by collisions and the charge neutrality condition  $\sum_a n_a e_a = 0$ , the intrinsic ambipolarity holds for both of the Pfirsch-Schlüter and banana-plateau particle fluxes in the same way as for the classical particle fluxes, which implies that the ambipolar conditions

$$\sum_a e_a J_{a1}^{\text{PS}} = \sum_a e_a J_{a1}^{\text{bp}} = 0. \quad (50)$$

are valid for arbitrary values of the thermodynamic forces ( $X_{a1}, X_{a2}, X_E$ ). On the other hand, the nonaxisymmetric particle fluxes  $J_{a1}^{\text{na}}$  and accordingly the total neoclassical particle fluxes  $J_{a1}^{\text{nc1}}$  are not ambipolar generally. If the ambipolarity condition for the total neoclassical particle fluxes

$$\sum_a e_a J_{a1}^{\text{nc1}} = \sum_a e_a J_{a1}^{\text{na}} = 0 \quad (51)$$

is imposed, we find in the similar way as in Eq.(31) that Eq.(48) is rewritten as

$$\sum_a T_a \langle \bar{\sigma}_a \rangle = \sum_{a \neq I} J_{a1}^{\text{nc1}} X_{a1}^* + \sum_a J_{a2}^{\text{nc1}} X_{a2} + J_E X_E. \quad (52)$$

In axisymmetric toroidal systems, the nonaxisymmetric fluxes  $J_{aj}^{\text{na}}$  ( $j = 1, 2$ ) vanish and therefore the neoclassical particle fluxes  $J_{a1}^{\text{nc1}}$  are intrinsically ambipolar. As discussed in the next section, in nonaxisymmetric systems, the ambipolarity condition (51) combined with the neoclassical transport equations gives a constraint on the thermodynamic forces ( $X_{a1}, X_{a2}, X_E$ ) from which the radial electric field  $-\Phi'$  is expressed by a linear form in the pressure and temperature gradients and the parallel electric field. Then, independent thermodynamic forces for nonaxisymmetric systems are given not by the set ( $X_{a1}, X_{a2}, X_E$ ) but by the reduced one ( $X_{a1(a \neq I)}^*, X_{a2}, X_E$ ). In the present work, we show the neoclassical transport equations for both cases with ( $X_{a1}, X_{a2}, X_E$ ) and with ( $X_{a1(a \neq I)}^*, X_{a2}, X_E$ ) used as the forces, in order to elucidate the relation of the ambipolarity to the axisymmetry and to the Onsager symmetry.

### III. ONSAGER SYMMETRY OF NEOCLASSICAL TRANSPORT EQUATIONS FOR GENERAL TOROIDAL SYSTEMS

In this section, it is proved that the Onsager symmetry is satisfied by the neoclassical transport equations for general toroidal systems including nonaxisymmetric cases. For that, it is convenient to define the distribution function  $\bar{g}_a$  by

$$\bar{g}_a = \bar{f}_{a1} - \frac{e_a}{T_a} f_{a0} \int^l \frac{dl}{B} \left[ BE_{\parallel} - \frac{B^2}{\langle B^2 \rangle} \langle BE_{\parallel} \rangle \right] \quad (53)$$

where  $\int^l dl$  denotes the integral along the magnetic field line. Then, Eq.(38) is rewritten as

$$v_{\parallel} \mathbf{n} \cdot \nabla \bar{g}_a - C_a^L(\bar{g}_a) = \frac{1}{T_a} f_{a0} (S_{a1} X_{a1} + S_{a2} X_{a2} + S_{a3} X_{a3}) \quad (54)$$

where the functions  $S_{aj}$  ( $j = 1, 2, 3$ ) are defined by

$$S_{aj} = \left( x_a^2 - \frac{5}{2} \right)^{j-1} \mathbf{v}_{da} \cdot \nabla V \quad (j = 1, 2), \quad S_{a3} = v_{\parallel} B / \langle B^2 \rangle^{1/2}. \quad (55)$$

The neoclassical radial fluxes  $J_{aj}^{\text{nc1}}$  ( $j = 1, 2$ ) and the parallel flux  $J_{a3}$  are expressed in terms of  $\bar{g}_a$  and  $S_{aj}$  ( $j = 1, 2, 3$ ) as

$$\begin{aligned} J_{aj}^{\text{nc1}} &\equiv \left\langle \int d^3v \bar{g}_a S_{aj} \right\rangle \quad (j = 1, 2) \\ J_{a3} &\equiv \left\langle \int d^3v \bar{g}_a S_{a3} \right\rangle \end{aligned} \quad (56)$$

where we have used the following formula for an arbitrary function  $F(\mathbf{x})$  :

$$\left\langle F \int d^3v \left( x_a^2 - \frac{5}{2} \right)^{j-1} (\mathbf{v}_{da} \cdot \nabla V) f_{a0} \right\rangle = -\delta_{j1} \frac{cp_a}{e_a B \theta} \left\langle \frac{B_{\zeta}}{B} \mathbf{n} \cdot \nabla F \right\rangle \quad (j = 1, 2) \quad (57)$$

Noting in Eq.(54) that the left-hand side is linear with respect to  $\bar{g}_a$  and that  $X_{aj}$  ( $j = 1, 2, 3$ ) occur in the right-hand side as parameters, we find that the solution  $\bar{g}_a$  of Eq.(54) is given by

$$\bar{g}_a = \sum_b (G_{ab1} X_{b1} + G_{ab2} X_{b2} + G_{ab3} X_{b3}) \quad (58)$$

where  $G_{abj}$  ( $j = 1, 2, 3$ ) are defined as the solutions of

$$v_{\parallel} \mathbf{n} \cdot \nabla G_{abj} - \sum_{a'} C_{aa'}^L(G_{abj}, G_{a'bj}) = \delta_{ab} \frac{1}{T_b} f_{b0} S_{bj} \quad (j = 1, 2, 3). \quad (59)$$

Then, using Eqs.(56) and (58), we obtain the transport equations relating  $J_{aj}^{\text{ncI}}$  ( $j = 1, 2$ ) and  $J_{a3}$  to  $X_{aj}$  ( $j = 1, 2, 3$ ) as

$$\begin{bmatrix} J_{a1} \\ J_{a2} \\ J_{a3} \end{bmatrix} = \sum_b \begin{bmatrix} L_{11}^{ab} & L_{12}^{ab} & L_{13}^{ab} \\ L_{21}^{ab} & L_{22}^{ab} & L_{23}^{ab} \\ L_{31}^{ab} & L_{32}^{ab} & L_{33}^{ab} \end{bmatrix} \begin{bmatrix} X_{b1} \\ X_{b2} \\ X_{b3} \end{bmatrix} \quad (60)$$

where the transport coefficients  $L_{jk}^{ab}$  are given by

$$L_{jk}^{ab} = \left\langle \int d^3v S_{aj} G_{abk} \right\rangle \quad (j, k = 1, 2, 3). \quad (61)$$

Equation (60) is rewritten as the transport equations relating  $J_{aj}^{\text{ncI}}$  ( $j = 1, 2$ ) and  $J_E$  to  $X_{aj}$  ( $j = 1, 2$ ) and  $X_E$  :

$$\begin{aligned} J_{aj}^{\text{ncI}} &= \sum_b \sum_{k=1,2} L_{jk}^{ab} X_{bk} + L_{jE}^a X_E \quad (j = 1, 2) \\ J_E &= \sum_b \sum_{k=1,2} L_{Ek}^b X_{bk} + L_{EE} X_E \end{aligned} \quad (62)$$

where the coefficients  $L_{jE}^a$ ,  $L_{Ej}^a$  ( $j = 1, 2$ ) and  $L_{EE}$  are given by

$$L_{jE}^a \equiv \sum_b e_b L_{j3}^{ab} \quad L_{Ej}^a \equiv \sum_b e_b L_{3j}^{ba} \quad (j = 1, 2) \quad L_{EE} \equiv \sum_{a,b} e_a e_b L_{33}^{ab}. \quad (63)$$

In order to show the Onsager symmetry of the transport coefficients, it is useful to separate Eq.(59) into even and odd parts with respect to  $v_{\parallel}$  as

$$\begin{aligned} v_{\parallel} \mathbf{n} \cdot \nabla G_{abj}^- - \sum_{a'} C_{aa'}^L(G_{abj}^+, G_{a'bj}^+) &= \delta_{ab} \frac{1}{T_b} f_{b0} S_{bj}^+ \\ v_{\parallel} \mathbf{n} \cdot \nabla G_{abj}^+ - \sum_{a'} C_{aa'}^L(G_{abj}^-, G_{a'bj}^-) &= \delta_{ab} \frac{1}{T_b} f_{b0} S_{bj}^- \end{aligned} \quad (64)$$

where the superscripts + and - denote the even and odd parts, respectively. Noting that  $S_{aj}$  are even for  $j = 1, 2$  and odd for  $j = 3$ , and using Eqs.(61) and (64), we obtain

$$\begin{aligned} \sum_A T_A \left\langle \int d^3v \frac{1}{f_{A0}} G_{Abk}^+ v_{\parallel} \mathbf{n} \cdot \nabla G_{Aaj}^- \right\rangle - \sum_{A,B} T_A \left\langle \int d^3v \frac{1}{f_{A0}} G_{Abk}^+ C_{AB}^L(G_{Aaj}^+, G_{Baj}^+) \right\rangle &= (\delta_{j1} + \delta_{j2}) L_{jk}^{ab} \\ \sum_A T_A \left\langle \int d^3v \frac{1}{f_{A0}} G_{Abk}^- v_{\parallel} \mathbf{n} \cdot \nabla G_{Aaj}^+ \right\rangle - \sum_{A,B} T_A \left\langle \int d^3v \frac{1}{f_{A0}} G_{Abk}^- C_{AB}^L(G_{Aaj}^-, G_{Baj}^-) \right\rangle &= \delta_{j3} L_{jk}^{ab} \end{aligned} \quad (65)$$

from which we have

$$(\delta_{j1} + \delta_{j2}) L_{jk}^{ab} + \delta_{k3} L_{kj}^{ba} = - \sum_{A,B} T_A \left\langle \int d^3v \frac{1}{f_{A0}} \left[ G_{Abk}^+ C_{AB}^L(G_{Aaj}^+, G_{Baj}^+) + G_{Aaj}^- C_{AB}^L(G_{Abk}^-, G_{Bbk}^-) \right] \right\rangle. \quad (66)$$

We find from the self-adjointness of the linearized collision operator given by Eq.(9) that the right-hand side of Eq.(66) is invariant under the permutation of the subscripts  $(a, j) \leftrightarrow (b, k)$  and that

$$(\delta_{j1} + \delta_{j2})L_{jk}^{ab} + \delta_{k3}L_{kj}^{ba} = (\delta_{k1} + \delta_{k2})L_{kj}^{ba} + \delta_{j3}L_{jk}^{ab}.$$

Thus, we obtain

$$(\delta_{j1} + \delta_{j2} - \delta_{j3})L_{jk}^{ab} = (\delta_{k1} + \delta_{k2} - \delta_{k3})L_{kj}^{ba}$$

which is rewritten as the well-known Onsager relations :

$$\begin{aligned} L_{jk}^{ab} &= L_{kj}^{ba} & (j, k = 1, 2) \\ L_{j3}^{ab} &= -L_{3j}^{ba} & (j = 1, 2) \\ L_{33}^{ab} &= L_{33}^{ba}. \end{aligned} \tag{67}$$

We see from Eqs.(63) and (67) that

$$L_{jE}^a = -L_{Ej}^a \quad (j = 1, 2). \tag{68}$$

Equations (67) and (68) show that the Onsager symmetry is satisfied by the transport matrix which combines the conjugate pairs of the fluxes  $(J_{a1}^{\text{nc1}}, J_{a2}^{\text{nc1}}, J_E)$  and the forces  $(X_{a1}, X_{a3}, X_E)$  for general toroidal systems.

Here, let us discuss the relation between the transport equations and the ambipolarity condition. In axisymmetric systems, intrinsic ambipolarity holds for the neoclassical particle fluxes and is expressed in terms of the relation between the transport coefficients as

$$\sum_a e_a L_{1j}^{ab} = \sum_a e_a L_{1E}^a = 0 \quad (j = 1, 2, 3). \tag{69}$$

Then, the number of the conjugate pairs of fluxes and forces is reduced by one as shown in Eq.(52) using  $X_{a1(a \neq I)}^*$  instead of  $X_{a1}$ . We find from Eqs.(67), (68), and (69) that the transport equations relating the fluxes  $(J_{a1(a \neq I)}^{\text{nc1}}, J_{a2}^{\text{nc1}}, J_E)$  to  $(X_{a1(a \neq I)}^*, X_{a2}, X_E)$  in the axisymmetric case are given by

$$\begin{aligned} J_{a1(a \neq I)}^{\text{nc1}} &= \sum_{b \neq I} L_{11}^{ab} X_{b1}^* + \sum_b L_{12}^{ab} X_{b2} + L_{1E}^a X_E \\ J_{a2}^{\text{nc1}} &= \sum_{b \neq I} L_{21}^{ab} X_{b1}^* + \sum_b L_{22}^{ab} X_{b2} + L_{2E}^a X_E \\ J_E &= \sum_{b \neq I} L_{E1}^b X_{b1}^* + \sum_b L_{E2}^b X_{b2} + L_{EE} X_E. \end{aligned} \tag{70}$$

In the transport equations (70), the transport coefficients  $L_{jk}^{ab}$  are the same as in Eq.(62) except for the limitation  $(a, j), (b, k) \neq (I, 1)$  and the Onsager symmetry is still valid.



In nonaxisymmetric systems, the ambipolarity condition (51) gives a relation between the thermodynamic forces which is used to express one thermodynamic force  $X_I$  in terms of the other thermodynamic forces ( $X_{a1(a \neq I)}^*$ ,  $X_{a2}$ ,  $X_E$ ) as

$$X_{I1} = -e_I \left( \sum_{a,b} e_a e_b L_{11}^{ab} \right)^{-1} \sum_a e_a \left( \sum_{b \neq I} L_{11}^{ab} X_{b1}^* + \sum_b L_{12}^{ab} X_{b2} + L_{1E}^a X_E \right). \quad (71)$$

Equation (71) determines the radial electric field  $-\Phi' = X_{I1}/e_I + p'_I/(n_I e_I)$  as a linear combination of the pressure and temperature gradients and the parallel electric field.

Then, in the nonaxisymmetric case with the ambipolarity condition (71), the transport equations relating ( $J_{a1(a \neq I)}^{\text{nc1}}$ ,  $J_{a2}^{\text{nc1}}$ ,  $J_E$ ) to ( $X_{a1(a \neq I)}^*$ ,  $X_{a2}$ ,  $X_E$ ) are given by

$$\begin{aligned} J_{a1(a \neq I)} &= \sum_{b \neq I} \mathcal{L}_{11}^{ab} X_{b1}^* + \sum_b \mathcal{L}_{12}^{ab} X_{b2} + \mathcal{L}_{1E}^a X_E \\ J_{a2} &= \sum_{b \neq I} \mathcal{L}_{21}^{ab} X_{b1}^* + \sum_b \mathcal{L}_{22}^{ab} X_{b2} + \mathcal{L}_{2E}^a X_E \\ J_E &= \sum_{b \neq I} \mathcal{L}_{E1}^b X_{b1}^* + \sum_b \mathcal{L}_{E2}^b X_{b2} + \mathcal{L}_{EE} X_E \end{aligned} \quad (72)$$

where the transport coefficients  $\mathcal{L}_{jk}^{ab}$ ,  $\mathcal{L}_{jE}^a$ ,  $\mathcal{L}_{Ej}^a$ , and  $\mathcal{L}_{EE}$  are

$$\begin{aligned} \mathcal{L}_{jk}^{ab} &= L_{jk}^{ab} - \left( \sum_A e_A L_{j1}^{aA} \right) \left( \sum_A e_A L_{1k}^{Ab} \right) \left( \sum_{A,B} e_A e_B L_{11}^{AB} \right)^{-1} \\ \mathcal{L}_{jE}^a &= L_{jE}^a - \left( \sum_A e_A L_{j1}^{aA} \right) \left( \sum_A e_A L_{1E}^A \right) \left( \sum_{A,B} e_A e_B L_{11}^{AB} \right)^{-1} \\ \mathcal{L}_{Ej}^a &= L_{Ej}^a - \left( \sum_A e_A L_{E1}^A \right) \left( \sum_A e_A L_{1j}^{Aa} \right) \left( \sum_{A,B} e_A e_B L_{11}^{AB} \right)^{-1} \\ \mathcal{L}_{EE} &= L_{EE} - \left( \sum_A e_A L_{E1}^A \right) \left( \sum_A e_A L_{1E}^A \right) \left( \sum_{A,B} e_A e_B L_{11}^{AB} \right)^{-1}. \end{aligned} \quad (73)$$

We see from Eqs.(67) and (68) that the Onsager symmetry still holds for the transport coefficients given by Eq.(73) :

$$\mathcal{L}_{jk}^{ab} = \mathcal{L}_{kj}^{ba} \quad \mathcal{L}_{jE}^a = -\mathcal{L}_{Ej}^a. \quad (74)$$

Thus, we have established for nonaxisymmetric systems with no net radial current that the radial electric field is determined by the pressure, temperature gradients, and the parallel electric field, and that transport satisfies the Onsager symmetry.

## IV. BANANA-PLATEAU AND NONAXISYMMETRIC TRANSPORT COEFFICIENTS

In the previous section, we have shown the Onsager symmetry for the transport coefficients relating the neoclassical radial fluxes and the parallel current ( $J_{a1}^{\text{nc1}}, J_{a2}^{\text{nc1}}, J_E$ ) to the radial gradient forces and the parallel electric field ( $X_{a1}, X_{a2}, X_E$ ) in general toroidal systems. The neoclassical radial particle and heat fluxes ( $J_{a1}^{\text{nc1}}, J_{a2}^{\text{nc1}}$ ) consist of the Pfirsch-Schlüter, banana-plateau, and nonaxisymmetric parts while, as shown in Appendix B, it is well-known that the transport equations for the Pfirsch-Schlüter fluxes ( $J_{a1}^{\text{PS}}, J_{a2}^{\text{PS}}$ ) and the radial gradient forces ( $X_{a1}, X_{a2}$ ) satisfy the Onsager symmetry. Thus, it is clear that the sum of the banana-plateau and nonaxisymmetric radial fluxes, and the parallel current ( $J_{a1}^{\text{bp}} + J_{a1}^{\text{na}}, J_{a2}^{\text{bp}} + J_{a2}^{\text{na}}, J_E$ ) are related to the forces ( $X_{a1}, X_{a2}, X_E$ ) by the transport coefficients with the Onsager symmetry. In this section, using the thirteen moment (13M) approximation,<sup>3</sup> we derive the transport equations for the banana-plateau fluxes and those for the nonaxisymmetric fluxes separately, in the case of general toroidal plasma consisting of electrons and one ion species. Then, the symmetry properties are investigated for each of the transport equations, and the Onsager symmetry for the total transport is directly confirmed.

The parallel momentum balance equations combined with the friction-flow relations are given in the 13M approximation by

$$\left[ \begin{array}{c} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle + n_e e \langle E_{\parallel} B \rangle \\ \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_e \rangle \end{array} \right] = \left[ \begin{array}{c} \langle B F_{\parallel e1} \rangle \\ \langle B F_{\parallel e2} \rangle \end{array} \right] = -\frac{n_e m_e}{\tau_{ee}} \left[ \begin{array}{cc} \hat{l}_{11}^e & -\hat{l}_{12}^e \\ -\hat{l}_{12}^e & \hat{l}_{22}^e \end{array} \right] \left[ \begin{array}{c} \langle B(u_{\parallel e} - u_{\parallel i}) \rangle \\ \frac{2}{5p_a} \langle B q_{\parallel e} \rangle \end{array} \right] \quad (75)$$

$$\langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_i \rangle = \langle B F_{\parallel i2} \rangle = -\frac{n_i m_i}{\tau_{ii}} \hat{l}_{22}^i \frac{2}{5p_i} \langle B q_{\parallel i} \rangle \quad (76)$$

where the dimensionless friction coefficients  $\hat{l}_{ij}^a \equiv -(\tau_{aa}/n_a m_a) l_{ij}^{aa}$  are given by  $\hat{l}_{11}^e = Z_i$ ,  $\hat{l}_{12}^e = \frac{3}{2} Z_i$ ,  $\hat{l}_{22}^e = \sqrt{2} + \frac{13}{4} Z_i$ , and  $\hat{l}_{22}^i = \sqrt{2}$  with the ion charge number  $Z_i$ . The species summation of the parallel momentum balances reduces to

$$\langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle + \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_i \rangle = 0 \quad (77)$$

where the momentum conservation (25) in collisions and the charge neutrality condition  $\sum_a e_a n_a = 0$  are used.

Solving the linearized drift kinetic equation gives the equations for the parallel viscosities, which have the following form for all collision frequencies in the Pfirsch-Schlüter, plateau, and banana regimes :

$$\left[ \begin{array}{c} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{array} \right] = \frac{n_a m_a}{\tau_{aa}} c_a \left[ \begin{array}{cc} \hat{\mu}_{a1} & \hat{\mu}_{a2} \\ \hat{\mu}_{a2} & \hat{\mu}_{a3} \end{array} \right] \left( \left[ \begin{array}{c} \langle u_{\parallel a} B \rangle \\ \frac{2}{5p_a} \langle q_{\parallel a} B \rangle \end{array} \right] - \langle G_a \rangle \frac{c}{e_a} \left[ \begin{array}{c} X_{a1} \\ X_{a2} \end{array} \right] \right) \quad (78)$$

where  $c_a$  and  $\hat{\mu}_{aj}$  ( $j = 1, 2, 3$ ) are the dimensionless parameters for the viscosity coefficients, and  $\langle G_a \rangle$  represents the geometrical factor which measures the deviation from the

axisymmetric configuration. These parameters  $c_a$ ,  $\hat{\mu}_{aj}$  ( $j = 1, 2, 3$ ) and  $\langle G_a \rangle$  are given in Appendix C and all of them are generally dependent on the collision frequencies although, in the axisymmetric case, the geometrical factor is given by  $\langle G_a \rangle = \langle B_\zeta \rangle / B^\theta$  for both species and for all the collision frequencies.

Using Eqs.(76)–(78), the ion parallel flows and viscosity are written as

$$\langle Bu_{\parallel i} \rangle = \langle G_i \rangle \frac{c}{e_i} \left( X_{i1} + \frac{\hat{\mu}'_{i2}}{\hat{\mu}'_{i1}} X_{i2} \right) - \frac{1}{c_i \hat{\mu}'_{i1}} \frac{\tau_{ii}}{n_i m_i} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle \quad (79)$$

$$\frac{2}{5p_i} \langle Bq_{\parallel i} \rangle = \frac{c_i (\hat{\mu}_{i1} \hat{\mu}_{i3} - (\hat{\mu}_{i2})^2)}{\hat{\mu}'_{i1} (c_i \hat{\mu}_{i3} + \hat{l}_{22}^i)} \langle G_i \rangle \frac{c}{e_i} X_{i2} + \frac{\hat{\mu}'_{i2}}{\hat{\mu}'_{i1} \hat{l}_{22}^i} \frac{\tau_{ii}}{n_i m_i} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle \quad (80)$$

$$\langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_i \rangle = - \frac{n_i m_i c_i \hat{l}_{22}^i (\hat{\mu}_{i1} \hat{\mu}_{i3} - (\hat{\mu}_{i2})^2)}{\tau_{ii} \hat{\mu}'_{i1} (c_i \hat{\mu}_{i3} + \hat{l}_{22}^i)} \langle G_i \rangle \frac{c}{e_i} X_{i2} - \frac{\hat{\mu}'_{i2}}{\hat{\mu}'_{i1}} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle \quad (81)$$

where we have defined  $\hat{\mu}'_{i1} \equiv \hat{\mu}_{i1} - \gamma \hat{\mu}_{i2}$  and  $\hat{\mu}'_{i2} \equiv \hat{\mu}_{i2} - \gamma \hat{\mu}_{i3}$  with  $\gamma \equiv c_i \hat{\mu}_{i2} / (c_i \hat{\mu}_{i3} + \hat{l}_{22}^i)$ . Equations relating the electron parallel viscosities  $\langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle$  and  $\langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_e \rangle$  to the thermodynamic forces ( $X_{e1}, X_{e2}, X_{i1}, X_{i2}, X_E$ ) are obtained from Eqs.(75), (78), and (79) as

$$\begin{aligned} & \left( \left[ \begin{array}{cc} \hat{l}_{11}^e & -\hat{l}_{12}^e \\ -\hat{l}_{12}^e & \hat{l}_{22}^e \end{array} \right]^{-1} + \frac{1}{c_e} \left[ \begin{array}{cc} \hat{\mu}_{e1} & \hat{\mu}_{e2} \\ \hat{\mu}_{e2} & \hat{\mu}_{e3} \end{array} \right]^{-1} + \frac{1}{c_i \hat{\mu}'_{i1}} \frac{n_e m_e \tau_{ii}}{n_i m_i \tau_{ee}} \left[ \begin{array}{cc} 1 & 0 \\ 0 & 0 \end{array} \right] \right) \left[ \begin{array}{c} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle \\ \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_e \rangle \end{array} \right] \\ & = - \left[ \begin{array}{cc} \hat{l}_{11}^e & -\hat{l}_{12}^e \\ -\hat{l}_{12}^e & \hat{l}_{22}^e \end{array} \right]^{-1} \left[ \begin{array}{c} n_e e \langle B^2 \rangle^{1/2} X_E \\ 0 \end{array} \right] + \frac{n_e m_e}{\tau_{ee}} \left[ \begin{array}{c} \langle G_e \rangle \frac{c}{e} X_{e1} + \langle G_i \rangle \frac{c}{e_i} \left( X_{i1} + \frac{\hat{\mu}'_{i2}}{\hat{\mu}'_{i1}} X_{i2} \right) \\ \langle G_e \rangle \frac{c}{e} X_{e2} \end{array} \right] \end{aligned} \quad (82)$$

We also find from Eq.(75) that the parallel current is divided into the classical part  $J_E^{\text{cl}}$  and the neoclassical part  $J_E^{\text{bp}}$  due to the electron parallel viscosities as

$$\begin{aligned} J_E & \equiv \langle B J_{\parallel} \rangle / \langle B^2 \rangle^{1/2} = J_E^{\text{cl}} + J_E^{\text{bp}} \\ J_E^{\text{cl}} & \equiv \sigma_s X_E \\ J_E^{\text{bp}} & \equiv \frac{\sigma_s}{n_e e} \langle B^2 \rangle^{-1/2} \left( \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_e \rangle + \frac{\hat{l}_{12}^e}{\hat{l}_{22}^e} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_e \rangle \right) \end{aligned} \quad (83)$$

where  $\sigma_s \equiv (n_e e^2 \tau_{ee} / m_e) \hat{l}_{22}^e / [\hat{l}_{11}^e \hat{l}_{22}^e - (\hat{l}_{12}^e)^2]$  is the classical Spitzer conductivity.

Finally, we obtain from Eqs.(77) and (81)–(83) the banana-plateau transport equations which relate  $(J_{e1}^{\text{bp}}, J_{e2}^{\text{bp}}, J_{i1}^{\text{bp}}, J_{i2}^{\text{bp}}, J_E^{\text{bp}})$  to  $(X_{e1}, X_{e2}, X_{i1}, X_{i2}, X_E)$  as

$$\begin{bmatrix} J_{e1}^{\text{bp}} \\ J_{e2}^{\text{bp}} \\ J_{i1}^{\text{bp}} \\ J_{i2}^{\text{bp}} \\ J_E^{\text{bp}} \end{bmatrix} = \begin{bmatrix} (L^{\text{bp}})_{11}^{ee} & (L^{\text{bp}})_{12}^{ee} & (L^{\text{bp}})_{11}^{ei} & (L^{\text{bp}})_{12}^{ei} & (L^{\text{bp}})_{1E}^e \\ (L^{\text{bp}})_{21}^{ee} & (L^{\text{bp}})_{22}^{ee} & (L^{\text{bp}})_{21}^{ei} & (L^{\text{bp}})_{22}^{ei} & (L^{\text{bp}})_{2E}^e \\ (L^{\text{bp}})_{11}^{ie} & (L^{\text{bp}})_{12}^{ie} & (L^{\text{bp}})_{11}^{ii} & (L^{\text{bp}})_{12}^{ii} & (L^{\text{bp}})_{1E}^i \\ (L^{\text{bp}})_{21}^{ie} & (L^{\text{bp}})_{22}^{ie} & (L^{\text{bp}})_{21}^{ii} & (L^{\text{bp}})_{22}^{ii} & (L^{\text{bp}})_{2E}^i \\ (L^{\text{bp}})_{E1}^e & (L^{\text{bp}})_{E2}^e & (L^{\text{bp}})_{E1}^i & (L^{\text{bp}})_{E2}^i & (L^{\text{bp}})_{EE} \end{bmatrix} \begin{bmatrix} X_{e1} \\ X_{e2} \\ X_{i1} \\ X_{i2} \\ X_E \end{bmatrix} \quad (84)$$

where the banana-plateau transport coefficients are given by

$$\begin{bmatrix} (L^{\text{bp}})_{11}^{ee} & (L^{\text{bp}})_{12}^{ee} \\ (L^{\text{bp}})_{21}^{ee} & (L^{\text{bp}})_{22}^{ee} \end{bmatrix} = \frac{n_e m_e c^2 \langle B_\zeta \rangle \langle G_e \rangle}{\tau_{ee} e^2 B^\theta \langle B^2 \rangle} \begin{bmatrix} M_1 & M_2 \\ M_2 & M_3 \end{bmatrix} \quad (85)$$

$$\begin{bmatrix} M_1 & M_2 \\ M_2 & M_3 \end{bmatrix} \equiv \left( \left[ \begin{array}{cc} \hat{l}_{11}^e & -\hat{l}_{12}^e \\ -\hat{l}_{12}^e & \hat{l}_{22}^e \end{array} \right]^{-1} + \frac{1}{c_e} \left[ \begin{array}{cc} \hat{\mu}_{e1} & \hat{\mu}_{e2} \\ \hat{\mu}_{e2} & \hat{\mu}_{e3} \end{array} \right]^{-1} + \frac{1}{c_i \hat{\mu}'_{i1}} \frac{n_e m_e \tau_{ii}}{n_i m_i \tau_{ee}} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \right)^{-1} \quad (86)$$

$$\begin{bmatrix} (L^{\text{bp}})_{11}^{ei} \\ (L^{\text{bp}})_{21}^{ei} \end{bmatrix} = \frac{1}{Z_i \langle G_e \rangle} \begin{bmatrix} (L^{\text{bp}})_{11}^{ee} \\ (L^{\text{bp}})_{21}^{ee} \end{bmatrix}, \quad \begin{bmatrix} (L^{\text{bp}})_{12}^{ei} \\ (L^{\text{bp}})_{22}^{ei} \end{bmatrix} = \frac{1}{Z_i \hat{\mu}'_{i1} \langle G_e \rangle} \begin{bmatrix} (L^{\text{bp}})_{11}^{ee} \\ (L^{\text{bp}})_{21}^{ee} \end{bmatrix} \quad (87)$$

$$\begin{bmatrix} (L^{\text{bp}})_{1E}^e \\ (L^{\text{bp}})_{2E}^e \end{bmatrix} = -\frac{n_e c}{[\hat{l}_{11}^e \hat{l}_{22}^e - (\hat{l}_{12}^e)^2] B^\theta \langle B^2 \rangle^{1/2}} \begin{bmatrix} M_1 & M_2 \\ M_2 & M_3 \end{bmatrix} \begin{bmatrix} \hat{l}_{22}^e \\ \hat{l}_{12}^e \end{bmatrix} \quad (88)$$

$$\begin{aligned} & \left[ (L^{\text{bp}})_{11}^{ie}, (L^{\text{bp}})_{12}^{ie}, (L^{\text{bp}})_{11}^{ii}, (L^{\text{bp}})_{12}^{ii}, (L^{\text{bp}})_{1E}^i \right] \\ &= \frac{1}{Z_i} \left[ (L^{\text{bp}})_{11}^{ee}, (L^{\text{bp}})_{12}^{ee}, (L^{\text{bp}})_{11}^{ei}, (L^{\text{bp}})_{12}^{ei}, (L^{\text{bp}})_{1E}^e \right] \end{aligned} \quad (89)$$

$$\begin{aligned} & \left[ (L^{\text{bp}})_{21}^{ie}, (L^{\text{bp}})_{22}^{ie}, (L^{\text{bp}})_{21}^{ii}, (L^{\text{bp}})_{22}^{ii}, (L^{\text{bp}})_{2E}^i \right] \\ &= \frac{\hat{\mu}'_{i2}}{\hat{\mu}'_{i1} Z_i} \left[ (L^{\text{bp}})_{11}^{ee}, (L^{\text{bp}})_{12}^{ee}, (L^{\text{bp}})_{11}^{ei}, (L^{\text{bp}})_{12}^{ei}, (L^{\text{bp}})_{1E}^e \right] \\ &+ \frac{n_i m_i c^2 c_i \hat{l}_{22}^i (\hat{\mu}_{i1} \hat{\mu}_{i3} - (\hat{\mu}_{i2})^2) \langle B_\zeta \rangle \langle G_i \rangle}{\tau_{ii} e_i^2 \hat{\mu}'_{i1} (c_i \hat{\mu}_{i3} + \hat{l}_{22}^i) B^\theta \langle B^2 \rangle} [0, 0, 0, 1, 0] \end{aligned} \quad (90)$$

$$\left[ (L^{\text{bp}})_{E1}^e, (L^{\text{bp}})_{E2}^e \right] = -\frac{B^\theta \langle G_e \rangle}{\langle B_\zeta \rangle} \left[ (L^{\text{bp}})_{1E}^e, (L^{\text{bp}})_{2E}^e \right] \quad (91)$$

$$\left[ (L^{\text{bp}})_{E1}^i, (L^{\text{bp}})_{E2}^i \right] = -\frac{B^\theta \langle G_i \rangle}{\langle B_\zeta \rangle} \left[ (L^{\text{bp}})_{1E}^i, (L^{\text{bp}})_{2E}^i \right] \quad (92)$$

$$(L^{\text{bp}})_{EE} = -\frac{\sigma_s}{\hat{l}_{22}^e [\hat{l}_{11}^e \hat{l}_{22}^e - (\hat{l}_{12}^e)^2]} \left[ \hat{l}_{22}^e \hat{l}_{12}^e \right] \begin{bmatrix} M_1 & M_2 \\ M_2 & M_3 \end{bmatrix} \begin{bmatrix} \hat{l}_{22}^e \\ \hat{l}_{12}^e \end{bmatrix}. \quad (93)$$

The banana-plateau transport coefficients given in Eqs.(85)–(93) satisfy the following symmetry properties :

$$\begin{aligned} (L^{\text{bp}})_{21}^{ee} &= (L^{\text{bp}})_{12}^{ee}, & (L^{\text{bp}})_{21}^{ii} &= (L^{\text{bp}})_{12}^{ii} \\ (L^{\text{bp}})_{jk}^{ie} &= \frac{\langle G_e \rangle}{\langle G_i \rangle} (L^{\text{bp}})_{kj}^{ei}, & (j, k &= 1, 2) \\ (L^{\text{bp}})_{Ej}^a &= -\frac{B^\theta \langle G_a \rangle}{\langle B_\zeta \rangle} (L^{\text{bp}})_{jE}^a & (a = e, i; j &= 1, 2). \end{aligned} \quad (94)$$

Thus, we confirm that the banana-plateau transport equations have the complete Onsager symmetry in the axisymmetric case where  $\langle G_e \rangle = \langle G_i \rangle = \langle B_\zeta \rangle / B^\theta$ . The intrinsic ambipolarity of the banana-plateau particle fluxes is expressed by

$$(L^{\text{bp}})_{1j}^{ia} = \frac{1}{Z_i} (L^{\text{bp}})_{1j}^{ea} \quad (a = e, i; j = 1, 2), \quad (L^{\text{bp}})_{1E}^i = \frac{1}{Z_i} (L^{\text{bp}})_{1E}^e \quad (95)$$

which are rewritten by Eq.(94) as

$$(L^{\text{bp}})_{j1}^{ai} = \frac{\langle G_i \rangle}{Z_i \langle G_e \rangle} (L^{\text{bp}})_{j1}^{ae} \quad (a = e, i; j = 1, 2), \quad (L^{\text{bp}})_{E1}^i = \frac{\langle G_i \rangle}{Z_i \langle G_e \rangle} (L^{\text{bp}})_{E1}^e. \quad (96)$$

Using the above relations, the fluxes ( $J_{e1}^{\text{bp}} (= Z_i J_{i1}^{\text{bp}})$ ,  $J_{e2}^{\text{bp}}$ ,  $J_{i2}^{\text{bp}}$ ,  $J_E$ ) are related to the new reduced set of forces ( $X_{e1}^{**}$ ,  $X_{e2}$ ,  $X_{i2}$ ,  $X_E$ ) by the transport equations

$$\begin{bmatrix} J_{e1}^{\text{bp}} \\ J_{e2}^{\text{bp}} \\ J_{i2}^{\text{bp}} \\ J_E^{\text{bp}} \end{bmatrix} = \begin{bmatrix} (L^{\text{bp}})_{11}^{ee} & (L^{\text{bp}})_{12}^{ee} & (L^{\text{bp}})_{12}^{ei} & (L^{\text{bp}})_{1E}^e \\ (L^{\text{bp}})_{21}^{ee} & (L^{\text{bp}})_{22}^{ee} & (L^{\text{bp}})_{22}^{ei} & (L^{\text{bp}})_{2E}^e \\ (L^{\text{bp}})_{21}^{ie} & (L^{\text{bp}})_{22}^{ie} & (L^{\text{bp}})_{22}^{ii} & (L^{\text{bp}})_{2E}^i \\ (L^{\text{bp}})_{E1}^e & (L^{\text{bp}})_{E2}^e & (L^{\text{bp}})_{E2}^i & (L^{\text{bp}})_{EE} \end{bmatrix} \begin{bmatrix} X_{e1}^{**} \\ X_{e2} \\ X_{i2} \\ X_E \end{bmatrix}. \quad (97)$$

Here, the force  $X_{e1}^{**}$  is defined by

$$X_{e1}^{**} \equiv X_{e1} + \frac{\langle G_i \rangle}{Z_i \langle G_e \rangle} X_{i1} \equiv -\frac{1}{n_e} \left( p'_e + \frac{\langle G_i \rangle}{\langle G_e \rangle} p'_i \right) - e\Phi' \left( \frac{\langle G_i \rangle}{\langle G_e \rangle} - 1 \right). \quad (98)$$

When  $\langle G_e \rangle = \langle G_i \rangle$ ,  $X_{e1}^{**}$  coincides with  $X_{e1}^*$ , which is proportional to the total pressure gradient  $-p' \equiv -(p'_e + p'_i)$ , and the radial electric field  $-\Phi'$  never affects the banana-plateau transport. However, if the electrons and the ions belong to different collisionality regimes in the nonaxisymmetric case, the banana-plateau radial particle and heat fluxes, and the bootstrap current depend on the radial electric field  $-\Phi'$  through  $X_{e1}^{**}$  since  $\langle G_e \rangle \neq \langle G_i \rangle$ .

Next, let us derive the nonaxisymmetric transport equations. For the derivation, it is essential to note that the toroidal viscosities are given in the following form for both of the Pfirsch-Schlüter and plateau regimes :

$$\begin{bmatrix} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} = \frac{n_a m_a}{\tau_{aa}} c_{ta} \begin{bmatrix} \hat{\mu}_{a1} & \hat{\mu}_{a2} \\ \hat{\mu}_{a2} & \hat{\mu}_{a3} \end{bmatrix} \left( \begin{bmatrix} \langle u_{\parallel a} B \rangle \\ \frac{2}{5p_a} \langle q_{\parallel a} B \rangle \end{bmatrix} - \langle G_{ta} \rangle \frac{c}{e_a} \begin{bmatrix} X_{a1} \\ X_{a2} \end{bmatrix} \right) \quad (99)$$

where  $c_{ta}$  and  $\langle G_{ta} \rangle$  are given in Appendix C and  $\hat{\mu}_{aj}$  ( $j = 1, 2, 3$ ) are the same as in Eq.(78). Here and hereafter, we consider the toroidal viscosities and the nonaxisymmetric fluxes only for the Pfirsch-Schlüter and plateau regimes since the expressions similar to Eq.(99) have not been obtained yet for the banana regime.

Appendix C shows that the ratio between the toroidal and parallel viscosities  $c_{ta}/c_a$  is related to the geometrical factor  $\langle G_a \rangle$  in Eq.(78) by the following equation :

$$1 - \frac{c_{ta}}{c_a} \frac{\langle B^2 \rangle}{B^\zeta \langle B_\zeta \rangle} = \frac{B^\theta \langle G_a \rangle}{\langle B_\zeta \rangle} \quad (100)$$

which is an essential relation for showing the Onsager symmetry of the total neoclassical transport.

Using Eqs.(78) and (99), the toroidal viscosities are written in terms of the parallel viscosities and the thermodynamic forces as

$$\begin{bmatrix} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} = \frac{c_{ta}}{c_a} \begin{bmatrix} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} + \frac{n_a m_a}{\tau_{aa}} c_{ta} (\langle G_a \rangle - \langle G_{ta} \rangle) \frac{c}{e_a} \begin{bmatrix} \hat{\mu}_{a1} \hat{\mu}_{a2} \\ \hat{\mu}_{a2} \hat{\mu}_{a3} \end{bmatrix} \begin{bmatrix} X_{a1} \\ X_{a2} \end{bmatrix}. \quad (101)$$

Then, from Eqs.(43), (84), and (101), we obtain the nonaxisymmetric transport equations relating  $(J_{e1}^{na}, J_{e2}^{na}, J_{i1}^{na}, J_{i2}^{na})$  to  $(X_{e1}, X_{e2}, X_{i1}, X_{i2}, X_E)$  :

$$\begin{bmatrix} J_{e1}^{na} \\ J_{e2}^{na} \\ J_{i1}^{na} \\ J_{i2}^{na} \end{bmatrix} = \begin{bmatrix} (L^{na})_{11}^{ee} & (L^{na})_{12}^{ee} & (L^{na})_{11}^{ei} & (L^{na})_{12}^{ei} & (L^{na})_{1E}^e \\ (L^{na})_{21}^{ee} & (L^{na})_{22}^{ee} & (L^{na})_{21}^{ei} & (L^{na})_{22}^{ei} & (L^{na})_{2E}^e \\ (L^{na})_{11}^{ie} & (L^{na})_{12}^{ie} & (L^{na})_{11}^{ii} & (L^{na})_{12}^{ii} & (L^{na})_{1E}^i \\ (L^{na})_{21}^{ie} & (L^{na})_{22}^{ie} & (L^{na})_{21}^{ii} & (L^{na})_{22}^{ii} & (L^{na})_{2E}^i \end{bmatrix} \begin{bmatrix} X_{e1} \\ X_{e2} \\ X_{i1} \\ X_{i2} \\ X_E \end{bmatrix} \quad (102)$$

where the nonaxisymmetric transport coefficients are given by

$$\begin{aligned} & \begin{bmatrix} (L^{na})_{11}^{ee} & (L^{na})_{12}^{ee} & (L^{na})_{11}^{ei} & (L^{na})_{12}^{ei} & (L^{na})_{1E}^e \\ (L^{na})_{21}^{ee} & (L^{na})_{22}^{ee} & (L^{na})_{21}^{ei} & (L^{na})_{22}^{ei} & (L^{na})_{2E}^e \end{bmatrix} \\ &= -\frac{c_{te}}{c_e} \frac{\langle B^2 \rangle}{B^\zeta \langle B_\zeta \rangle} \begin{bmatrix} (L^{bp})_{11}^{ee} & (L^{bp})_{12}^{ee} & (L^{bp})_{11}^{ei} & (L^{bp})_{12}^{ei} & (L^{bp})_{1E}^e \\ (L^{bp})_{21}^{ee} & (L^{bp})_{22}^{ee} & (L^{bp})_{21}^{ei} & (L^{bp})_{22}^{ei} & (L^{bp})_{2E}^e \end{bmatrix} \\ &+ \frac{n_e m_e c^2}{\tau_{ee} e^2} c_{te} \frac{(\langle G_e \rangle - \langle G_{te} \rangle)}{B^\theta B^\zeta} \begin{bmatrix} \hat{\mu}_{e1} \hat{\mu}_{e2} & 0 & 0 & 0 \\ \hat{\mu}_{e2} \hat{\mu}_{e3} & 0 & 0 & 0 \end{bmatrix} \end{aligned} \quad (103)$$

$$\begin{aligned} & \begin{bmatrix} (L^{na})_{11}^{ie} & (L^{na})_{12}^{ie} & (L^{na})_{11}^{ii} & (L^{na})_{12}^{ii} & (L^{na})_{1E}^i \\ (L^{na})_{21}^{ie} & (L^{na})_{22}^{ie} & (L^{na})_{21}^{ii} & (L^{na})_{22}^{ii} & (L^{na})_{2E}^i \end{bmatrix} \\ &= -\frac{c_{ti}}{c_i} \frac{\langle B^2 \rangle}{B^\zeta \langle B_\zeta \rangle} \begin{bmatrix} (L^{bp})_{11}^{ie} & (L^{bp})_{12}^{ie} & (L^{bp})_{11}^{ii} & (L^{bp})_{12}^{ii} & (L^{bp})_{1E}^i \\ (L^{bp})_{21}^{ie} & (L^{bp})_{22}^{ie} & (L^{bp})_{21}^{ii} & (L^{bp})_{22}^{ii} & (L^{bp})_{2E}^i \end{bmatrix} \\ &+ \frac{n_i m_i c^2}{\tau_{ii} e_i^2} c_{ti} \frac{(\langle G_i \rangle - \langle G_{ti} \rangle)}{B^\theta B^\zeta} \begin{bmatrix} 0 & 0 & \hat{\mu}_{i1} & \hat{\mu}_{i2} & 0 \\ 0 & 0 & \hat{\mu}_{i2} & \hat{\mu}_{i3} & 0 \end{bmatrix}. \end{aligned} \quad (104)$$

It is found that  $(L^{na})_{jk}^{aa} = (L^{na})_{kj}^{aa}$  is always valid although  $(L^{na})_{jk}^{ei} = (L^{na})_{kj}^{ie}$  is satisfied only when  $c_{te}/c_e = c_{ti}/c_i$ . We see from Eq.(100) that the condition  $c_{te}/c_e = c_{ti}/c_i$  is equivalent to  $\langle G_e \rangle = \langle G_i \rangle$ , which holds if the electrons and the ions belong to the same collisionality regime.

Finally, combining the banana-plateau transport equations (84) with the nonaxisymmetric transport equations (102), and using the relation (100), the transport equations for the

total of the banana-plateau and nonaxisymmetric fluxes ( $J_{e1}^{\text{bn}}, J_{e2}^{\text{bn}}, J_{i1}^{\text{bn}}, J_{i2}^{\text{bn}}, J_E^{\text{bp}}$ ) are obtained as

$$\begin{bmatrix} J_{e1}^{\text{bn}} \\ J_{e2}^{\text{bn}} \\ J_{i1}^{\text{bn}} \\ J_{i2}^{\text{bn}} \\ J_E^{\text{bp}} \end{bmatrix} \equiv \begin{bmatrix} J_{e1}^{\text{bp}} + J_{e1}^{\text{na}} \\ J_{e2}^{\text{bp}} + J_{e2}^{\text{na}} \\ J_{i1}^{\text{bp}} + J_{i1}^{\text{na}} \\ J_{i2}^{\text{bp}} + J_{i2}^{\text{na}} \\ J_E^{\text{bp}} \end{bmatrix} = \begin{bmatrix} (L^{\text{bn}})_{11}^{ee} & (L^{\text{bn}})_{12}^{ee} & (L^{\text{bn}})_{11}^{ei} & (L^{\text{bn}})_{12}^{ei} & (L^{\text{bn}})_{1E}^e \\ (L^{\text{bn}})_{21}^{ee} & (L^{\text{bn}})_{22}^{ee} & (L^{\text{bn}})_{21}^{ei} & (L^{\text{bn}})_{22}^{ei} & (L^{\text{bn}})_{2E}^e \\ (L^{\text{bn}})_{11}^{ie} & (L^{\text{bn}})_{12}^{ie} & (L^{\text{bn}})_{11}^{ii} & (L^{\text{bn}})_{12}^{ii} & (L^{\text{bn}})_{1E}^i \\ (L^{\text{bn}})_{21}^{ie} & (L^{\text{bn}})_{22}^{ie} & (L^{\text{bn}})_{21}^{ii} & (L^{\text{bn}})_{22}^{ii} & (L^{\text{bn}})_{2E}^i \\ (L^{\text{bn}})_{E1}^e & (L^{\text{bn}})_{E2}^e & (L^{\text{bn}})_{E1}^i & (L^{\text{bn}})_{E2}^i & (L^{\text{bn}})_{EE} \end{bmatrix} \begin{bmatrix} X_{e1} \\ X_{e2} \\ X_{i1} \\ X_{i2} \\ X_E \end{bmatrix} \quad (105)$$

where the total transport coefficients are given by

$$\begin{bmatrix} (L^{\text{bn}})_{11}^{ee} & (L^{\text{bn}})_{12}^{ee} & (L^{\text{bn}})_{11}^{ei} & (L^{\text{bn}})_{12}^{ei} & (L^{\text{bn}})_{1E}^e \\ (L^{\text{bn}})_{21}^{ee} & (L^{\text{bn}})_{22}^{ee} & (L^{\text{bn}})_{21}^{ei} & (L^{\text{bn}})_{22}^{ei} & (L^{\text{bn}})_{2E}^e \end{bmatrix} = \frac{B^\theta \langle G_e \rangle}{\langle B_\zeta \rangle} \begin{bmatrix} (L^{\text{bp}})_{11}^{ee} & (L^{\text{bp}})_{12}^{ee} & (L^{\text{bp}})_{11}^{ei} & (L^{\text{bp}})_{12}^{ei} & (L^{\text{bp}})_{1E}^e \\ (L^{\text{bp}})_{21}^{ee} & (L^{\text{bp}})_{22}^{ee} & (L^{\text{bp}})_{21}^{ei} & (L^{\text{bp}})_{22}^{ei} & (L^{\text{bp}})_{2E}^e \end{bmatrix} + \frac{n_e m_e c^2}{\tau_{ee} e^2} \frac{(\langle G_e \rangle - \langle G_{te} \rangle)}{B^\theta B^\zeta} \begin{bmatrix} \hat{\mu}_{e1} & \hat{\mu}_{e2} & 0 & 0 & 0 \\ \hat{\mu}_{e2} & \hat{\mu}_{e3} & 0 & 0 & 0 \end{bmatrix} \quad (106)$$

$$\begin{bmatrix} (L^{\text{bn}})_{11}^{ie} & (L^{\text{bn}})_{12}^{ie} & (L^{\text{bn}})_{11}^{ii} & (L^{\text{bn}})_{12}^{ii} & (L^{\text{bn}})_{1E}^i \\ (L^{\text{bn}})_{21}^{ie} & (L^{\text{bn}})_{22}^{ie} & (L^{\text{bn}})_{21}^{ii} & (L^{\text{bn}})_{22}^{ii} & (L^{\text{bn}})_{2E}^i \end{bmatrix} = \frac{B^\theta \langle G_i \rangle}{\langle B_\zeta \rangle} \begin{bmatrix} (L^{\text{bp}})_{11}^{ie} & (L^{\text{bp}})_{12}^{ie} & (L^{\text{bp}})_{11}^{ii} & (L^{\text{bp}})_{12}^{ii} & (L^{\text{bp}})_{1E}^i \\ (L^{\text{bp}})_{21}^{ie} & (L^{\text{bp}})_{22}^{ie} & (L^{\text{bp}})_{21}^{ii} & (L^{\text{bp}})_{22}^{ii} & (L^{\text{bp}})_{2E}^i \end{bmatrix} + \frac{n_i m_i c^2}{\tau_{ii} e_i^2} \frac{(\langle G_i \rangle - \langle G_{ti} \rangle)}{B^\theta B^\zeta} \begin{bmatrix} 0 & 0 & \hat{\mu}_{i1} & \hat{\mu}_{i2} & 0 \\ 0 & 0 & \hat{\mu}_{i2} & \hat{\mu}_{i3} & 0 \end{bmatrix} \quad (107)$$

$$\begin{bmatrix} (L^{\text{bn}})_{E1}^e & (L^{\text{bn}})_{E2}^e & (L^{\text{bn}})_{E1}^i & (L^{\text{bn}})_{E2}^i & (L^{\text{bn}})_{EE} \end{bmatrix} = \begin{bmatrix} (L^{\text{bp}})_{E1}^e & (L^{\text{bp}})_{E2}^e & (L^{\text{bp}})_{E1}^i & (L^{\text{bp}})_{E2}^i & (L^{\text{bp}})_{EE} \end{bmatrix}. \quad (108)$$

Thus, from the above definitions and the symmetry properties given by Eq.(94), we can directly confirm that the total banana-plateau and nonaxisymmetric transport coefficients satisfy the following Onsager symmetry :

$$(L^{\text{bn}})_{jk}^{ab} = (L^{\text{bn}})_{kj}^{ba}, \quad (L^{\text{bn}})_{Ej}^a = -(L^{\text{bn}})_{jE}^a \quad (a, b = e, i; j, k = 1, 2). \quad (109)$$

Here, we should note that these transport coefficients contain terms of different orders in  $(m_e/m_i)^{1/2}$ . As seen from Eqs.(85)–(93) and Eqs.(106)–(108), the ion-ion coefficients  $(L^{\text{bn}})_{jk}^{ii}$  are  $\mathcal{O}((m_i/m_e)^{1/2})$  larger than the other coefficients.

When the ambipolarity condition is imposed in the nonaxisymmetric case, we obtain, in the similar way as in Eqs.(71)–(73), the radial electric field :

$$\begin{aligned} -\Phi' &= \frac{p'_i}{n_i e_i} + \frac{X_{i1}}{e_i} \\ &= \frac{p'_i}{n_e e} - \left( \sum_{a,b=e,i} e_a e_b (L^{\text{bn}})_{11}^{ab} \right)^{-1} \\ &\quad \times \sum_{a=e,i} e_a \left[ (L^{\text{bn}})_{11}^{ae} X_{e1}^* + (L^{\text{bn}})_{12}^{ae} X_{e2} + (L^{\text{bn}})_{12}^{ai} X_{i2} + (L^{\text{bn}})_{1E}^a X_E \right] \end{aligned} \quad (110)$$

and the reduced set of the transport equations :

$$\begin{bmatrix} J_{e1}^{\text{bn}} \\ J_{e2}^{\text{bn}} \\ J_{i2}^{\text{bn}} \\ J_E^{\text{bp}} \end{bmatrix} = \begin{bmatrix} (\mathcal{L}^{\text{bn}})_{11}^{ee} & (\mathcal{L}^{\text{bn}})_{12}^{ee} & (\mathcal{L}^{\text{bn}})_{12}^{ei} & (\mathcal{L}^{\text{bn}})_{1E}^e \\ (\mathcal{L}^{\text{bn}})_{21}^{ee} & (\mathcal{L}^{\text{bn}})_{22}^{ee} & (\mathcal{L}^{\text{bn}})_{22}^{ei} & (\mathcal{L}^{\text{bn}})_{2E}^e \\ (\mathcal{L}^{\text{bn}})_{21}^{ie} & (\mathcal{L}^{\text{bn}})_{22}^{ie} & (\mathcal{L}^{\text{bn}})_{22}^{ii} & (\mathcal{L}^{\text{bn}})_{2E}^i \\ (\mathcal{L}^{\text{bn}})_{E1}^e & (\mathcal{L}^{\text{bn}})_{E2}^e & (\mathcal{L}^{\text{bn}})_{E2}^i & (\mathcal{L}^{\text{bn}})_{EE} \end{bmatrix} \begin{bmatrix} X_{e1}^* \\ X_{e2} \\ X_{i2} \\ X_E \end{bmatrix} \quad (111)$$

where the transport coefficients are defined by

$$\begin{aligned} (\mathcal{L}^{\text{bn}})_{jk}^{ab} &= (L^{\text{bn}})_{jk}^{ab} - \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{j1}^{aA} \right) \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{1k}^{Ab} \right) \left( \sum_{A,B=e,i} e_A e_B (L^{\text{bn}})_{11}^{AB} \right)^{-1} \\ (\mathcal{L}^{\text{bn}})_{jE}^a &= (L^{\text{bn}})_{jE}^a - \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{j1}^{aA} \right) \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{1E}^A \right) \left( \sum_{A,B=e,i} e_A e_B (L^{\text{bn}})_{11}^{AB} \right)^{-1} \\ (\mathcal{L}^{\text{bn}})_{Ej}^a &= (L^{\text{bn}})_{Ej}^a - \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{E1}^A \right) \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{1j}^{Aa} \right) \left( \sum_{A,B=e,i} e_A e_B (L^{\text{bn}})_{11}^{AB} \right)^{-1} \\ (\mathcal{L}^{\text{bn}})_{EE} &= (L^{\text{bn}})_{EE} - \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{E1}^A \right) \left( \sum_{A=e,i} e_A (L^{\text{bn}})_{1E}^A \right) \left( \sum_{A,B=e,i} e_A e_B (L^{\text{bn}})_{11}^{AB} \right)^{-1} \end{aligned} \quad (112)$$

The Onsager symmetry still holds for the above coefficients :

$$(\mathcal{L}^{\text{bn}})_{jk}^{ab} = (\mathcal{L}^{\text{bn}})_{kj}^{ba} \quad (\mathcal{L}^{\text{bn}})_{jE}^a = -(\mathcal{L}^{\text{bn}})_{Ej}^a. \quad (113)$$

As mentioned earlier, terms of different orders in  $(m_e/m_i)^{1/2}$  are included in the transport coefficients  $(L^{\text{bn}})_{jk}^{ab}$ . Therefore, the coefficients  $(\mathcal{L}^{\text{bn}})_{jk}^{ab}$  for the reduced transport equations also contain different order terms. To the lowest order of  $(m_e/m_i)^{1/2}$ , Eq.(110) is approximated by

$$-\Phi' \simeq \frac{p'_i}{n_e e} - \frac{(L^{\text{na}})_{12}^{ii} X_{i2}}{(L^{\text{na}})_{11}^{ii} e_i} \quad (114)$$

which is the same one given in Ref.8 and in Ref.11. In Ref.11, Balescu and Fantechi used this approximate expression for the radial electric field instead of Eq.(110) to derive the reduced set of the transport equations. Then, their resultant transport coefficients are different from ours in Eq.(112) and do not satisfy the Onsager symmetry since part of  $\mathcal{O}((m_e/m_i)^{1/2})$  terms in Eq.(112) are inconsistently neglected.

## V. CONCLUSIONS AND DISCUSSION

In this work, we have investigated the entropy production, the full transport equations and their Onsager symmetry for the neoclassical transport processes in magnetically confined plasmas with general toroidal configurations. It was clearly shown that, for both the



classical and neoclassical transport processes, the kinetic form of the entropy production defined from the linearized collision operator is equivalent to its thermodynamic form written as the inner product of the thermodynamic forces and their conjugate transport fluxes. The entropy production from the gyroangle-dependent distribution function corresponds to the sum of the products of the classical radial particle and heat fluxes and the radial gradient thermodynamic forces, while the magnetic surface average of the entropy production from the gyroangle-averaged distribution function is given by the sum of the products of the thermodynamic forces and their conjugate fluxes which consist of the Pfirsch-Schlüter, banana-plateau, nonaxisymmetric parts of the neoclassical radial fluxes and the parallel current. This equivalence between the kinetic and thermodynamic forms of the entropy production for the full neoclassical transport fluxes were not confirmed by Balescu in Chap.17 of Ref.3. The reason is now discussed.

In deriving the thermodynamic form of the entropy production, we used the linearized drift kinetic equation without employing the Hermitian moment expansion of the distribution function. Balescu expressed the kinetic form of the entropy production as the quadratic form of only the vector Hermitian moment part of the distribution function which corresponds to the  $l = 1$  part of the Legendre polynomial expansion. However, since the neoclassical banana-plateau and nonaxisymmetric fluxes are caused by the parallel and toroidal viscosities, the tensor Hermitian moment part (or the  $l = 2$  part of the Legendre polynomial expansion) needs to be included for deriving the neoclassical thermodynamic form of the entropy production. Furthermore, as shown in Appendix D, we find that it is necessary to include all the tensor moments including higher order parts with  $l = 3, 4, 5, \dots$  in the kinetic form of the entropy production to derive the neoclassical thermodynamic form in the banana and plateau regimes. This is intuitively understandable by considering the resonant particles responsible for the neoclassical fluxes in the plateau regime. The resonant particle distribution is highly anisotropic in velocity space and is approximated by a delta function in pitch angle so that the all Hermitian moments (or all  $l$ th-order Legendre polynomials) are required. As shown in Appendix D, the operator  $v_{\parallel} \mathbf{n} \cdot \nabla$  in the drift kinetic equation (38) introduces the anisotropic distribution in the velocity space and it connects the  $l$ th-order moment with  $(l \pm 1)$ th-order moments in contrast with the linearized collision operator which is isotropic in the velocity space and connects the  $l$ th-order moment with the same  $l$ th-order moment alone. In the Pfirsch-Schlüter regime, the collision operator dominates  $v_{\parallel} \mathbf{n} \cdot \nabla$  and the distribution function has small contributions from higher order moments representing the anisotropy. Then, in the Pfirsch-Schlüter regime, the  $l = 1$  vector moment is enough to express the entropy production as in Ref.3, while the negligibly small viscosity-induced neoclassical fluxes are included in the  $l = 2$  tensor moment part. On the other hand, as the collision frequency decreases, the operator  $v_{\parallel} \mathbf{n} \cdot \nabla$  is comparable to, and then dominates, the collision operator and all  $l$ -order moments are required in the kinetic entropy production functional to obtain its neoclassical thermodynamic form in the banana and plateau regimes.

We also proved from the formal solution of the linearized drift kinetic equation with the self-adjoint linearized collision operator that the Onsager symmetry is robustly valid for the neoclassical transport equations for general toroidal plasmas consisting of electrons and

multi-species ions with arbitrary collision frequencies. Furthermore, we derived in Sec.IV, in the case of a single ion species, the full banana-plateau transport coefficients for all collisionality regimes and the full nonaxisymmetric transport coefficients for the Pfirsch-Schlüter and plateau regimes. The symmetry properties of these transport matrices were separately examined and the Onsager symmetry for their total transport equations was confirmed. We discussed the effects of the ambipolarity condition on the transport equations in detail for both axisymmetric and nonaxisymmetric configurations. The ambipolarity condition reduces by one the number of the conjugate pairs of the transport fluxes and the thermodynamic forces. In the axisymmetric case, the intrinsic ambipolarity holds and the radial electric field does not affect the transport. On the other hand, for broken toroidal symmetry, a radial current is a function of the thermodynamic forces ( $X_{aj}, X_E$ ) in which the radial electric field is included. When the ambipolarity condition is imposed in the nonaxisymmetric case, the radial electric field is given by a linear combination of the other thermodynamic forces. We showed that the Onsager symmetry is satisfied whether the conjugate pairs of the fluxes and forces are reduced by the ambipolarity condition or not.

Balescu and Fantechi derived the full neoclassical transport coefficients for the plateau regime in the nonaxisymmetric configuration and claimed that the Onsager symmetry is slightly broken by the nonaxisymmetry. They showed the transport equations only for the reduced pairs of the fluxes and forces, in which the radial electric field is eliminated by the ambipolar condition. There, terms of  $\mathcal{O}((m_e/m_i)^{1/2})$  were neglected in expressing the radial electric field in terms of the other forces as in Eq.(114). Then, the resultant transport coefficients in Ref.11 deviate from those in Eq.(112) and the Onsager symmetry is broken in them since part of  $\mathcal{O}((m_e/m_i)^{1/2})$  terms necessary for the symmetry are dropped.

The banana-plateau and nonaxisymmetric transport equations obtained here are valid whether electrons and ions belong to the same collisionality regime or not. When both species are in the same collisionality regime, we find that the geometrical factors for electrons and ions coincide with each other  $\langle G_e \rangle = \langle G_i \rangle$  and that, as far as the radial fluxes and the radial forces are concerned, the Onsager symmetry is separately valid for the banana-plateau transport matrix  $(L^{\text{bp}})_{jk}^{ab}$  and for the nonaxisymmetric transport matrix  $(L^{\text{na}})_{jk}^{ab}$ . When electron and ion collisionality regimes are different,  $\langle G_e \rangle \neq \langle G_i \rangle$  and the mixed electron-ion coefficients in each matrix are not symmetric although the total matrix  $L^{\text{bn}}$  are symmetric. In the latter case, the radial electric field appears in the thermodynamic forces for the banana-plateau radial particle and heat fluxes and the bootstrap current in the nonaxisymmetric systems.

Since we proved the robust validity of the Onsager symmetry for the neoclassical transport equations even in the nonaxisymmetric cases, this symmetry property can be utilized for the calculation of the nonaxisymmetric transport coefficients in the banana regime which were not given in Sec.IV. For example, from the banana-plateau transport coefficients  $(L^{\text{bp}})_{jE}^a$  and  $(L^{\text{bp}})_{Ej}^a$  ( $a = e, i$ ;  $j = 1, 2$ ) for the banana regime given in Sec.IV, we can immediately obtain part of the nonaxisymmetric transport coefficients for the banana regime as  $(L^{\text{na}})_{jE}^a = -(L^{\text{bp}})_{jE}^a - (L^{\text{bp}})_{Ej}^a$  ( $a = e, i$ ;  $j = 1, 2$ ).

In the previous work, we investigated the neoclassical and anomalous transport in weakly

turbulent plasma and described the entropy production and the Onsager symmetry in electrostatic turbulence.<sup>16</sup> We are also investigating a unified description of the transport equations, the entropy production, and the Onsager symmetry for neoclassical and turbulent processes with both electrostatic and electromagnetic fluctuations, which we will report on in a future work.

## **ACKNOWLEDGMENTS**

The author (HS) thanks Prof. M. Okamoto for his encouragement of this work. This work is supported in part by the Grant-in-Aid from the Japanese Ministry of Education, Science and Culture, and in part by the U.S. Department of Energy Grant DE-FG05-80ET-53088.

## APPENDIX A: HAMADA COORDINATES AND INCOMPRESSIBLE FLOWS

In general toroidal configurations, the magnetic field is written in the covariant form :

$$\mathbf{B} = B^\theta \partial \mathbf{x} / \partial \theta + B^\zeta \partial \mathbf{x} / \partial \zeta \quad (\text{A1})$$

where  $\theta$  and  $\zeta$  are the poloidal and toroidal angle variables, respectively, and corresponding basis vectors for the covariant representation are given by  $\partial \mathbf{x} / \partial \theta$  and  $\partial \mathbf{x} / \partial \zeta$ . Throughout this paper, we employ the Hamada coordinates  $(V, \theta, \zeta)$ , where the flux label  $V$  represents the volume enclosed by the flux surface, and the periods of the angle variables are normalized as  $\oint d\theta = \oint d\zeta = 1$ . The Jacobian is unity  $\nabla V \cdot \nabla \theta \times \nabla \zeta = 1$  so that the flux surface average is simply written as  $\langle \cdot \rangle = \oint d\theta \oint d\zeta \cdot$ . The poloidal and toroidal fields are given by  $\mathbf{B}_p = B^\theta \partial \mathbf{x} / \partial \theta = B^\theta \nabla \zeta \times \nabla V$  and  $\mathbf{B}_t = B^\zeta \partial \mathbf{x} / \partial \zeta = B^\zeta \nabla \theta \times \nabla V$ . The contravariant poloidal and toroidal components are the surface flux quantities  $B^\theta = \mathbf{B} \cdot \nabla \theta = \chi'(V)$  and  $B^\zeta = \mathbf{B} \cdot \nabla \zeta = \psi'(V)$ , where  $\chi$  and  $\psi$  are the poloidal and toroidal fluxes, respectively, and  $' \equiv \partial / \partial V$ . The vector product of  $\mathbf{B}$  and  $\nabla V$  is given by the linear combination of  $\mathbf{B}$  and  $\partial \mathbf{x} / \partial \zeta$  as

$$\mathbf{B} \times \nabla V = \frac{B_\zeta}{B^\theta} \mathbf{B} - \frac{B^2}{B^\theta} \frac{\partial \mathbf{x}}{\partial \zeta} \quad (\text{A2})$$

where  $B_\zeta = \mathbf{B} \cdot \partial \mathbf{x} / \partial \zeta$ .

When a solenoidal vector field  $\mathbf{U}$  ( $\nabla \cdot \mathbf{U} = 0$ ) is tangential to magnetic surfaces  $\mathbf{U} \cdot \nabla V = 0$  and satisfies  $\mathbf{U} \times \mathbf{B} = \nabla K(V)$  with some flux quantity  $K(V)$ , it is written in the Hamada coordinates as

$$\mathbf{U} = U^\theta \partial \mathbf{x} / \partial \theta + U^\zeta \partial \mathbf{x} / \partial \zeta \quad (\text{A3})$$

where the both contravariant components  $U^\theta = \mathbf{U} \cdot \nabla \theta$  and  $U^\zeta = \mathbf{U} \cdot \nabla \zeta$  are flux surface quantities. Since both of the flow velocity  $\mathbf{u}_a$  and the heat flow  $\mathbf{q}_a$  are incompressible  $\nabla \cdot \mathbf{u}_a = \nabla \cdot \mathbf{q}_a = 0$  (see Eq.(D12)) and satisfies the above conditions (see Eq.(20)) to the lowest order in  $\delta$ , all of their contravariant components  $u_a^\theta = \mathbf{u}_a \cdot \nabla \theta$ ,  $u_a^\zeta = \mathbf{u}_a \cdot \nabla \zeta$ ,  $q_a^\theta = \mathbf{q}_a \cdot \nabla \theta$ ,  $q_a^\zeta = \mathbf{q}_a \cdot \nabla \zeta$  are flux surface quantities. Also,  $\mathbf{u}_a$  and  $\mathbf{q}_a$  are separated into the parallel and perpendicular components as

$$\mathbf{u}_a = u_{\parallel a} \mathbf{n} + \mathbf{u}_{\perp a}, \quad \mathbf{q}_a = q_{\parallel a} \mathbf{n} + \mathbf{q}_{\perp a} \quad (\text{A4})$$

where the perpendicular components  $\mathbf{u}_{\perp a}$  and  $\mathbf{q}_{\perp a}$  are given by the thermodynamic forces  $X_{a1}$  and  $X_{a2}$  as in Eq.(20). Then, the contravariant poloidal and toroidal flow components are given by the linear combinations of the parallel flow components and the thermodynamic forces as

$$u_a^\theta = \frac{B^\theta}{B} u_{\parallel a} - \frac{B_\zeta}{B^2} \frac{cX_{a1}}{e_a}$$

$$\begin{aligned}
u_a^\zeta &= \frac{B^\zeta}{B} u_{\parallel a} + \frac{B_\theta}{B^2} \frac{cX_{a1}}{e_a} \\
\frac{2}{5} \frac{q_a^\theta}{p_a} &= \frac{B^\theta}{B} \frac{2}{5} \frac{q_{\parallel a}}{p_a} - \frac{B_\zeta}{B^2} \frac{cX_{a2}}{e_a} \\
\frac{2}{5} \frac{q_a^\zeta}{p_a} &= \frac{B^\zeta}{B} \frac{2}{5} \frac{q_{\parallel a}}{p_a} + \frac{B_\theta}{B^2} \frac{cX_{a2}}{e_a}
\end{aligned} \tag{A5}$$

where  $B_\theta = \mathbf{B} \cdot \partial \mathbf{x} / \partial \theta$ . We find from Eq.(A5) that

$$\begin{aligned}
\langle B^2 \rangle u_a^\theta &= B^\theta \langle u_{\parallel a} B \rangle - \langle B_\zeta \rangle \frac{cX_{a1}}{e_a} \\
\langle B^2 \rangle u_a^\zeta &= B^\zeta \langle u_{\parallel a} B \rangle + \langle B_\theta \rangle \frac{cX_{a1}}{e_a} \\
\langle B^2 \rangle \frac{2}{5} \frac{q_a^\theta}{p_a} &= B^\theta \frac{2}{5} \frac{\langle q_{\parallel a} B \rangle}{p_a} - \langle B_\zeta \rangle \frac{cX_{a2}}{e_a} \\
\langle B^2 \rangle \frac{2}{5} \frac{q_a^\zeta}{p_a} &= B^\zeta \frac{2}{5} \frac{\langle q_{\parallel a} B \rangle}{p_a} + \langle B_\theta \rangle \frac{cX_{a2}}{e_a}.
\end{aligned} \tag{A6}$$

The following formulae are obtained from Eqs.(A5) and (A6):

$$\begin{aligned}
\frac{u_{\parallel a}}{B} &= \frac{\langle u_{\parallel a} B \rangle}{\langle B^2 \rangle} + \frac{1}{B^\theta} \left( \frac{B_\zeta}{B^2} - \frac{\langle B_\zeta \rangle}{\langle B^2 \rangle} \right) \frac{cX_{a1}}{e_a} \\
&= \frac{\langle u_{\parallel a} B \rangle}{\langle B^2 \rangle} - \frac{1}{B^\zeta} \left( \frac{B_\theta}{B^2} - \frac{\langle B_\theta \rangle}{\langle B^2 \rangle} \right) \frac{cX_{a1}}{e_a} \\
\frac{2}{5} \frac{q_{\parallel a}}{p_a} \frac{1}{B} &= \frac{2}{5} \frac{\langle q_{\parallel a} B \rangle}{p_a \langle B^2 \rangle} + \frac{1}{B^\theta} \left( \frac{B_\zeta}{B^2} - \frac{\langle B_\zeta \rangle}{\langle B^2 \rangle} \right) \frac{cX_{a2}}{e_a} \\
&= \frac{2}{5} \frac{\langle q_{\parallel a} B \rangle}{p_a \langle B^2 \rangle} - \frac{1}{B^\zeta} \left( \frac{B_\theta}{B^2} - \frac{\langle B_\theta \rangle}{\langle B^2 \rangle} \right) \frac{cX_{a2}}{e_a}.
\end{aligned} \tag{A7}$$

## APPENDIX B: PFIRSCH-SCHLÜTER TRANSPORT EQUATIONS

The relations between the parallel friction forces and the parallel flows are given by the 13M approximation in the same form as in Eq.(27), and are written as

$$\begin{bmatrix} F_{\parallel a1} \\ -F_{\parallel a2} \end{bmatrix} = \sum_b \begin{bmatrix} l_{11}^{ab} & l_{12}^{ab} \\ l_{21}^{ab} & l_{22}^{ab} \end{bmatrix} \begin{bmatrix} u_{\parallel b} \\ -\frac{2}{5p_b} q_{\parallel b} \end{bmatrix} \quad (\text{B1})$$

where the coefficients  $l_{jk}^{ab}$  are defined in Eq.(28). Then, the Pfirsch-Schlüter particle and heat fluxes ( $J_{a1}^{\text{PS}}, J_{a2}^{\text{PS}}$ ) defined in Eq.(43) are rewritten as

$$\begin{bmatrix} J_{a1}^{\text{PS}} \\ J_{a2}^{\text{PS}} \end{bmatrix} = -\frac{c}{e_a B^\theta} \sum_b \begin{bmatrix} l_{11}^{ab} & -l_{12}^{ab} \\ -l_{21}^{ab} & l_{22}^{ab} \end{bmatrix} \begin{bmatrix} \left\langle \frac{u_{\parallel a}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \\ \left\langle \frac{2}{5p_a} \frac{q_{\parallel a}}{B} \left( B_\zeta - \langle B_\zeta \rangle \frac{B^2}{\langle B^2 \rangle} \right) \right\rangle \end{bmatrix}. \quad (\text{B2})$$

Substituting Eq.(A7) into Eq.(B2), we obtain the Pfirsch-Schlüter transport equations:

$$\begin{bmatrix} J_{a1}^{\text{PS}} \\ J_{a2}^{\text{PS}} \end{bmatrix} = \sum_b \begin{bmatrix} (L^{\text{PS}})_{11}^{ab} & (L^{\text{PS}})_{12}^{ab} \\ (L^{\text{PS}})_{21}^{ab} & (L^{\text{PS}})_{22}^{ab} \end{bmatrix} \begin{bmatrix} X_{a1} \\ X_{a2} \end{bmatrix} \quad (\text{B3})$$

where the Pfirsch-Schlüter transport coefficients  $(L^{\text{PS}})_{jk}^{ab}$  are given by

$$(L^{\text{PS}})_{jk}^{ab} = \frac{(-1)^{j+k-1}}{e_a e_b} \left( \frac{c}{B^\theta} \right)^2 \left( \left\langle \frac{B_\zeta^2}{B^2} \right\rangle - \frac{\langle B_\zeta \rangle^2}{\langle B^2 \rangle} \right) l_{jk}^{ab}, \quad (j, k = 1, 2). \quad (\text{B4})$$

From Eqs.(29) and (B4), we have the Onsager symmetry for the Pfirsch-Schlüter transport as

$$(L^{\text{PS}})_{jk}^{ab} = (L^{\text{PS}})_{kj}^{ba}, \quad (j, k = 1, 2). \quad (\text{B5})$$

The momentum conservation property described by Eq.(30) reduces to

$$\sum_a e_a (L^{\text{PS}})_{1k}^{ab} = 0, \quad (k = 1, 2). \quad (\text{B6})$$

which implies the intrinsic ambipolarity of the Pfirsch-Schlüter particle fluxes. We easily find from Eqs.(32), (B3), (B5), and (B6) that the Pfirsch-Schlüter transport equations for the pairs  $(J_{a1}^{\text{PS}}, X_{a1}^*)_{a \neq I}$ ,  $(J_{a2}^{\text{PS}}, X_{a2})$  are given by

$$\begin{aligned} J_{a1(a \neq I)}^{\text{PS}} &= \sum_{b \neq I} (L^{\text{PS}})_{11}^{ab} X_{b1}^* + \sum_b (L^{\text{PS}})_{12}^{ab} X_{b2} \\ J_{a2}^{\text{PS}} &= \sum_{b \neq I} (L^{\text{PS}})_{21}^{ab} X_{b1}^* + \sum_b (L^{\text{PS}})_{22}^{ab} X_{b2} \end{aligned} \quad (\text{B7})$$

which shows that the transport coefficients  $(L^{\text{PS}})_{jk}^{ab}$  are the same as in Eq.(B3) except for the limitation  $(a, j), (b, k) \neq (I, 1)$  and that the Onsager symmetry is valid for both of the conjugate pairs.

## APPENDIX C: PARALLEL AND TOROIDAL VISCOSITIES

It is shown from the solution of the linearized drift kinetic equation that the parallel viscosities in general toroidal configurations for all collision frequencies in the Pfirsch-Schlüter, plateau, and banana regimes are given in the following form :

$$\begin{bmatrix} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} = \frac{n_a m_a}{\tau_{aa}} c_a \begin{bmatrix} \hat{\mu}_{a1} & \hat{\mu}_{a2} \\ \hat{\mu}_{a2} & \hat{\mu}_{a3} \end{bmatrix} \left( \begin{bmatrix} \langle u_{\parallel a} B \rangle \\ \frac{2}{5p_a} \langle q_{\parallel a} B \rangle \end{bmatrix} - \langle G_a \rangle \frac{c}{e_a} \begin{bmatrix} X_{a1} \\ X_{a2} \end{bmatrix} \right) \quad (\text{C1})$$

where  $c_a$  and  $\hat{\mu}_{aj}$  ( $j = 1, 2, 3$ ) are the dimensionless parameters for the viscosity coefficients and  $\langle G_a \rangle$  represents the geometrical factor. The collision frequency  $\tau_{aa}$  is defined in Ref.2. The dimensionless viscosity parameters  $\hat{\mu}_{aj}$  ( $j = 1, 2, 3$ ) are written as

$$\hat{\mu}_{aj} = \begin{cases} \{x_a^2(x_a^2 - \frac{5}{2})^{j-1}(\nu_T^a \tau_{aa})^{-1}\} & \text{(for the Pfirsch-Schlüter regime)} \\ \delta_{j1} + \frac{1}{2}\delta_{j2} + \frac{13}{4}\delta_{j3} & \text{(for the plateau regime)} \\ \{(x_a^2 - \frac{5}{2})^{j-1}(\nu_D^a \tau_{aa})\} & \text{(for the banana regime)} \end{cases} \quad (\text{C2})$$

where the frequencies  $\nu_T^a$  and  $\nu_D^a$  are given in Ref.2, and the velocity-space average  $\{\cdot\} \equiv \frac{8}{3\sqrt{\pi}} \int_0^\infty dx_a x_a^4 e^{-x_a^2}$ . is used. The dimensionless coefficient  $c_a$  is given by

$$c_a = \begin{cases} \frac{3}{5}(v_{Ta} \tau_{aa})^2 \langle (\mathbf{B} \cdot \nabla \ln B)^2 \rangle / \langle B^2 \rangle & \text{(for the Pfirsch-Schlüter regime)} \\ \sqrt{\pi} v_{Ta} \tau_{aa} \langle B^2 \rangle^{-1/2} \sum_{m,n} 2\pi |(\ln B)_{mn}|^2 |mB^\theta - nB^\zeta| & \text{(for the plateau regime)} \\ f_t / f_c & \text{(for the banana regime)} \end{cases} \quad (\text{C3})$$

where  $f_c$  and  $f_t \equiv 1 - f_c$  denote the fractions of circulating and trapped particles defined in Ref.2 (or in Ref.7), respectively, and  $(\ln B)_{mn}$  are the coefficients in the Fourier expansion of  $\ln B$  :

$$\ln B = \sum_{m,n} (\ln B)_{mn} \exp[2\pi(m\theta - n\zeta)].$$

The geometrical factor  $\langle G_a \rangle$  is written as

$$\langle G_a \rangle = \begin{cases} \langle (\mathbf{B} \cdot \nabla \ln B)^2 \rangle^{-1} \left[ \langle B_\zeta \rangle \left\langle \frac{\partial \ln B}{\partial \theta} (\mathbf{B} \cdot \nabla \ln B) \right\rangle - \langle B_\theta \rangle \left\langle \frac{\partial \ln B}{\partial \zeta} (\mathbf{B} \cdot \nabla \ln B) \right\rangle \right] & \text{(for the Pfirsch-Schlüter regime)} \\ \left[ \sum_{m,n} |(\ln B)_{mn}|^2 |mB^\theta - nB^\zeta| \right]^{-1} \sum_{m,n} |(\ln B)_{mn}|^2 (m \langle B_\zeta \rangle + n \langle B_\theta \rangle) \frac{mB^\theta - nB^\zeta}{|mB^\theta - nB^\zeta|} & \text{(for the plateau regime).} \end{cases} \quad (\text{C4})$$

The geometrical factor  $\langle G_a \rangle$  for the banana regime is given in Ref.7 (or in Ref.9) as  $\tilde{G}_b$ . When we put  $(\ln B)_{mn} = 0$  for all  $n \neq 0$  in Eqs.(C3) and (C4), the expressions for axisymmetric systems are reproduced. In the axisymmetric case,  $\langle G_a \rangle$  is independent of the collision frequency and is given by  $\langle G_a \rangle = \langle B_\zeta \rangle / B^\theta$  for all particle species.

It is shown that the toroidal viscosities are given in the following form for both of the Pfirsch-Schlüter and plateau regimes :

$$\begin{bmatrix} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} = \frac{n_a m_a}{\tau_{aa}} c_{ta} \begin{bmatrix} \hat{\mu}_{a1} & \hat{\mu}_{a2} \\ \hat{\mu}_{a2} & \hat{\mu}_{a3} \end{bmatrix} \left( \left[ \begin{array}{c} \langle u_{\parallel a} B \rangle \\ \frac{2}{5 p_a} \langle q_{\parallel a} B \rangle \end{array} \right] - \langle G_{ta} \rangle \frac{c}{e_a} \begin{bmatrix} X_{a1} \\ X_{a2} \end{bmatrix} \right) \quad (\text{C5})$$

where  $\hat{\mu}_{aj}$  ( $j = 1, 2, 3$ ) are the same as given by Eq.(C2) and

$$c_{ta} = \begin{cases} \frac{3}{5} (v_{Ta} \tau_{aa})^2 \langle (\mathbf{B}_t \cdot \nabla \ln B)(\mathbf{B} \cdot \nabla \ln B) \rangle / \langle B^2 \rangle & (\text{for the Pfirsch-Schlüter regime}) \\ \sqrt{\pi} v_{Ta} \tau_{aa} \langle B^2 \rangle^{-1/2} \sum_{m,n} 2\pi |(\ln B)_{mn}|^2 (-n B^\zeta) \frac{m B^\theta - n B^\zeta}{|m B^\theta - n B^\zeta|} & (\text{for the plateau regime}) \end{cases} \quad (\text{C6})$$

$$\langle G_{ta} \rangle = \begin{cases} \langle (\mathbf{B}_t \cdot \nabla \ln B)(\mathbf{B} \cdot \nabla \ln B) \rangle^{-1} \left[ \langle B_\zeta \rangle \left\langle \frac{\partial \ln B}{\partial \theta} (\mathbf{B}_t \cdot \nabla \ln B) \right\rangle - \langle B_\theta \rangle \left\langle \frac{\partial \ln B}{\partial \zeta} (\mathbf{B}_t \cdot \nabla \ln B) \right\rangle \right] & (\text{for the Pfirsch-Schlüter regime}) \\ \left[ \sum_{m,n} |(\ln B)_{mn}|^2 \frac{(-n B^\zeta)(m B^\theta - n B^\zeta)}{|m B^\theta - n B^\zeta|} \right]^{-1} \sum_{m,n} |(\ln B)_{mn}|^2 (m \langle B_\zeta \rangle + n \langle B_\theta \rangle) \frac{(-n B^\zeta)}{|m B^\theta - n B^\zeta|} & (\text{for the plateau regime}). \end{cases} \quad (\text{C7})$$

The expressions similar to Eq.(C5) has not been obtained yet for the banana regime. We find from Eqs.(C3), (C4), and (C6) that the ratio between the toroidal and parallel viscosities  $c_{ta}/c_a$  is related to the geometrical factor  $\langle G_a \rangle$  by the following equation :

$$1 - \frac{c_{ta}}{c_a} \frac{\langle B^2 \rangle}{B^\zeta \langle B_\zeta \rangle} = \frac{B^\theta \langle G_a \rangle}{\langle B_\zeta \rangle} \quad (\text{C8})$$

which is an essential relation for showing the Onsager symmetry of the total neoclassical transport. We find from Eqs.(C1) and (C5) that the toroidal viscosities are written in terms of the parallel viscosities and the thermodynamic forces as

$$\begin{bmatrix} \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B}_t \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} = \frac{c_{ta}}{c_a} \begin{bmatrix} \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\pi}_a \rangle \\ \langle \mathbf{B} \cdot \nabla \cdot \boldsymbol{\Theta}_a \rangle \end{bmatrix} + \frac{n_a m_a}{\tau_{aa}} c_{ta} (\langle G_a \rangle - \langle G_{ta} \rangle) \frac{c}{e_a} \begin{bmatrix} \hat{\mu}_{a1} & \hat{\mu}_{a2} \\ \hat{\mu}_{a2} & \hat{\mu}_{a3} \end{bmatrix} \begin{bmatrix} X_{a1} \\ X_{a2} \end{bmatrix}. \quad (\text{C9})$$



## APPENDIX D: DRIFT KINETIC EQUATION AND LEGENDRE POLYNOMIAL EXPANSION

The linearized drift kinetic equation is written as

$$v_{\parallel} \mathbf{n} \cdot \nabla \bar{f}_{a1} + \mathbf{v}_{da} \cdot \nabla f_{a0} - \frac{e_a}{T_a} v_{\parallel} E_{\parallel} f_{a0} = C_a^L(\bar{f}_{a1}) \quad (\text{D1})$$

where  $\bar{f}_{a1}$  and  $f_{a0}$  are regarded as functions of the phase space variables  $(\mathbf{x}, E, \mu)$  ( $E \equiv \frac{1}{2} m_a v^2 + e_a \Phi$ : the particle's energy;  $\mu = m_a v_{\perp}^2 / 2B$ : the magnetic moment). Using  $(\mathbf{x}, v, \xi)$  ( $\xi \equiv v_{\parallel} / v$ ) as the phase space variables instead of  $(\mathbf{x}, E, \mu)$ , we consider the expansion by the Legendre polynomials  $P_l(\xi)$  ( $P_0(\xi) = 1, P_1(\xi) = \xi, P_2(\xi) = \frac{3}{2}\xi^2 - \frac{1}{2}, \dots$ ) for an arbitrary function  $F(\mathbf{x}, v, \xi)$  as

$$F(\mathbf{x}, v, \xi) = \sum_{l=0}^{\infty} F^{(l)}(\mathbf{x}, v, \xi)$$

$$F^{(l)}(\mathbf{x}, v, \xi) = P_l(\xi) \frac{2l+1}{2} \int_{-1}^1 d\eta P_l(\eta) F(\mathbf{x}, v, \eta). \quad (\text{D2})$$

The  $l$ th Legendre component corresponds to the  $l$ -order tensor Hermitian moment part in the Hermitian moment representation employed by Balescu in Ref.3. The first term in the left-hand side of Eq.(D1) is rewritten in the new phase space variables as

$$v \left( \xi \mathbf{n} \cdot \nabla \bar{f}_{a1} - (\mathbf{n} \cdot \nabla \ln B) \frac{1 - \xi^2}{2} \frac{\partial \bar{f}_{a1}}{\partial \xi} \right) \equiv \mathcal{A} \bar{f}_{a1}. \quad (\text{D3})$$

We have the following formulae for the Legendre polynomials :

$$\xi P_l(\xi) = \frac{1}{2l+1} [l P_{l-1}(\xi) + (l+1) P_{l+1}(\xi)]$$

$$(1 - \xi^2) \frac{dP_l(\xi)}{d\xi} = \frac{l(l+1)}{2l+1} [P_{l-1}(\xi) - P_{l+1}(\xi)]. \quad (\text{D4})$$

We find from Eqs.(D3) and (D4) that the operator  $\mathcal{A}$  ( $\equiv v_{\parallel} \mathbf{n} \cdot \nabla$  with  $(\mathbf{x}, E, \mu)$  as the phase space variables) in the drift kinetic equation (D1) transforms the  $l$ th Legendre component to the linear combination of the  $(l \pm 1)$ th components :

$$\mathcal{A} F^{(l)} = (\mathcal{A} F)^{(l-1)} + (\mathcal{A} F)^{(l+1)}. \quad (\text{D5})$$

Contrastively, from the velocity space isotropy of the collision operator described in Eq.(3), the operator  $C_a^L$  transforms the  $l$ th component to only the  $l$ th one. The second and third terms in the right-hand side of Eq.(D1) are rewritten from Eqs.(18) and (20) as

$$\left[ \frac{2}{3} + \frac{1}{3} P_2(\xi) \right] x_a^2 \left[ \nabla \cdot \mathbf{u}_{\perp a} + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5p_a} \nabla \cdot \mathbf{q}_{\perp a} \right] f_{a0} \quad (\text{D6})$$

and

$$-\frac{e_a E_{\parallel}}{T_a} v \xi f_{a0} \quad (\text{D7})$$

respectively. The former is proportional to the divergence of the diamagnetic flows and contains only the 0th and 2nd Legendre polynomial components while the latter is proportional to the parallel electric field and contains only the 1st Legendre component.

Now, let us write the drift kinetic equation (D1) by each Legendre component, separately. The 0th Legendre component (or the scalar moment part) of Eq.(D1) is given by

$$(\mathcal{A} \bar{f}_{a1}^{(l=1)})^{(l=0)} + \frac{2}{3} x_a^2 \left[ \nabla \cdot \mathbf{u}_{\perp a} + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5 p_a} \nabla \cdot \mathbf{q}_{\perp a} \right] f_{a0} = C_a^L(\bar{f}_{a1}^{(l=0)}). \quad (\text{D8})$$

Here, the  $l = 1$  Legendre component  $\bar{f}_{a1}^{(l=1)}$  of the distribution function is expanded by the Laguerre polynomials  $L_j^{(3/2)}(x_a^2)$  ( $L_0^{(3/2)}(x_a^2) = 1, L_1^{(3/2)}(x_a^2) = \frac{5}{2} - x_a^2, \dots$ ) as

$$\bar{f}_{a1}^{(l=1)} = \frac{2}{v T_a} \xi x_a \left[ u_{\parallel a} + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5} \frac{q_{\parallel a}}{p_a} \right] f_{a0} + \bar{f}_{a1}^{(l=1, j \geq 2)} \quad (\text{D9})$$

where  $\bar{f}_{a1}^{(l=1, j \geq 2)}$  denotes the sum of the  $j$ th Laguerre polynomial components with  $j \geq 2$ , which is neglected in the 13M approximation. The first term of Eq.(D8) is rewritten as

$$(\mathcal{A} \bar{f}_{a1}^{(l=1)})^{(l=0)} = \frac{2}{3} x_a^2 \left[ \nabla \cdot (u_{\parallel a} \mathbf{n}) + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5 p_a} \nabla \cdot (q_{\parallel a} \mathbf{n}) \right] f_{a0} + (\mathcal{A} \bar{f}_{a1}^{(l=1, j \geq 2)})^{(l=0)}. \quad (\text{D10})$$

Then, Eq.(D8) reduces to

$$\frac{2}{3} x_a^2 \left[ \nabla \cdot \mathbf{u}_a + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5 p_a} \nabla \cdot \mathbf{q}_a \right] f_{a0} + (\mathcal{A} \bar{f}_{a1}^{(l=1, j \geq 2)})^{(l=0)} = C_a^L(\bar{f}_{a1}^{(l=0)}). \quad (\text{D11})$$

Integrating Eq.(D11) multiplied by 1 and  $x_a^2$  over the velocity space, we obtain the incompressibility of  $\mathbf{u}_a$  and  $\mathbf{q}_a$  :

$$\nabla \cdot \mathbf{u}_a = \nabla \cdot \mathbf{q}_a = 0 \quad (\text{D12})$$

where we used the particle number and energy conservation by the collision operator described in Eq.(2). (Exactly speaking,  $\nabla \cdot \mathbf{q}_a = 0$  is valid to the lowest order of the small mass ratio as in Eq.(9).) Thus, we have

$$(\mathcal{A} \bar{f}_{a1}^{(l=1, j \geq 2)})^{(l=0)} = C_a^L(\bar{f}_{a1}^{(l=0)}). \quad (\text{D13})$$

from which the contribution from the scalar moment part  $\bar{f}_{a1}^{(l=0)}$  of the distribution function to the kinetic form of the entropy production is given by

$$\bar{\sigma}_a^{(l=0)} \equiv - \int d^3 v \frac{\bar{f}_{a1}^{(l=0)}}{f_{a0}} C_a^L(\bar{f}_{a1}^{(l=0)}) = - \int d^3 v \frac{\bar{f}_{a1}^{(l=0)}}{f_{a0}} \mathcal{A} \bar{f}_{a1}^{(l=1, j \geq 2)}. \quad (\text{D14})$$

In the 13M approximation,  $\bar{\sigma}_a^{(l=0)}$  vanishes.

Taking the 1st Legendre component (or the vector moment part) of Eq.(D1), we have

$$[\mathcal{A}(\bar{f}_{a1}^{(l=0)} + \bar{f}_{a1}^{(l=2)})]^{(l=1)} - \frac{e_a E_{\parallel a}}{T_a} v \xi f_{a0} = C_a^L(\bar{f}_{a1}^{(l=1)}) \quad (\text{D15})$$

We obtain the following parallel momentum balance equations from velocity integration of Eq.(D15) multiplied by  $m_a v \xi$  and  $m_a v \xi (x_a^2 - \frac{5}{2})$  :

$$\begin{aligned} \mathbf{n} \cdot (\nabla p_{a1} + \nabla \cdot \boldsymbol{\pi}_a) - n_a e_a E_{\parallel} &= F_{\parallel a1} \\ \mathbf{n} \cdot (\nabla \theta_{a1} + \nabla \cdot \boldsymbol{\Theta}_a) &= F_{\parallel a2}. \end{aligned} \quad (\text{D16})$$

Using Eqs.(D9) and (D15), the contribution from the vector moment part  $\bar{f}_{a1}^{(l=1)}$  to the kinetic form of the entropy production is given by

$$\begin{aligned} \bar{\sigma}_a^{(l=1)} &\equiv - \int d^3 v \frac{\bar{f}_{a1}^{(l=1)}}{f_{a0}} C_a^L(\bar{f}_{a1}^{(l=1)}) \\ &= \frac{1}{T_a} n_a e_a u_{\parallel a} E_{\parallel} - \int d^3 v \frac{\bar{f}_{a1}^{(l=1)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l=0)} + \bar{f}_{a1}^{(l=2)}) \\ &= \frac{1}{T_a} n_a e_a u_{\parallel a} E_{\parallel} - \frac{1}{T_a} \left[ u_{\parallel a} \mathbf{n} \cdot (\nabla p_{a1} + \nabla \cdot \boldsymbol{\pi}_a) + \frac{2}{5} \frac{q_{\parallel a}}{p_a} \mathbf{n} \cdot (\nabla \theta_{a1} + \nabla \cdot \boldsymbol{\Theta}_a) \right] \\ &\quad - \int d^3 v \frac{\bar{f}_{a1}^{(l=1, j \geq 2)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l=0)} + \bar{f}_{a1}^{(l=2)}) \end{aligned} \quad (\text{D17})$$

where the last integral vanishes in the 13M approximation.

The 2nd Legendre component (or the tensor moment part) of Eq.(D1) is given by

$$[\mathcal{A}(\bar{f}_{a1}^{(l=1)} + \bar{f}_{a1}^{(l=3)})]^{(l=2)} + \frac{1}{3} P_2(\xi) x_a^2 \left[ \nabla \cdot \mathbf{u}_{\perp a} + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5 p_a} \nabla \cdot \mathbf{q}_{\perp a} \right] f_{a0} = C_a^L(\bar{f}_{a1}^{(l=2)}) \quad (\text{D18})$$

which is rewritten by Eqs.(D9) and (D12) as

$$2 P_2(\xi) x_a^2 \left[ \mathbf{u}_a + \left( x_a^2 - \frac{5}{2} \right) \frac{2}{5 p_a} \mathbf{q}_a \right] \cdot (\nabla \ln B) f_{a0} + [\mathcal{A}(\bar{f}_{a1}^{(l=1, j \geq 2)} + \bar{f}_{a1}^{(l=3)})]^{(l=2)} = C_a^L(\bar{f}_{a1}^{(l=2)}). \quad (\text{D19})$$

The contribution from the tensor moment part  $\bar{f}_{a1}^{(l=2)}$  to the kinetic form of the entropy production is given by

$$\bar{\sigma}_a^{(l=2)} \equiv - \int d^3 v \frac{\bar{f}_{a1}^{(l=2)}}{f_{a0}} C_a^L(\bar{f}_{a1}^{(l=2)})$$

$$\begin{aligned}
&= -\frac{1}{T_a} \left[ (p_{\parallel a} - p_{\perp a}) \mathbf{u}_a + (\Theta_{\parallel a} - \Theta_{\perp a}) \frac{2}{5p_a} \mathbf{q}_a \right] \cdot \nabla \ln B \\
&\quad - \int d^3v \frac{\bar{f}_{a1}^{(l=2)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l=1, j \geq 2)} + \bar{f}_{a1}^{(l=3)}). \tag{D20}
\end{aligned}$$

Similarly, for higher order tensor moment parts with  $l \geq 3$ , we have

$$[\mathcal{A}(\bar{f}_{a1}^{(l-1)} + \bar{f}_{a1}^{(l+1)})]^{(l)} = C_a^L(\bar{f}_{a1}^{(l)}) \quad (\text{for } l \geq 3) \tag{D21}$$

from which we obtain

$$\bar{\sigma}_a^{(l)} \equiv - \int d^3v \frac{\bar{f}_{a1}^{(l)}}{f_{a0}} C_a^L(\bar{f}_{a1}^{(l)}) = - \int d^3v \frac{\bar{f}_{a1}^{(l)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l-1)} + \bar{f}_{a1}^{(l+1)}) \quad (\text{for } l \geq 3). \tag{D22}$$

Taking the magnetic surface average of each order moment part of the entropy production in Eqs.(D14), (D17), (D20), and (D22) gives

$$\begin{aligned}
\langle \bar{\sigma}_a^{(l=0)} \rangle &= - \left\langle \int d^3v \frac{\bar{f}_{a1}^{(l=0)}}{f_{a0}} \mathcal{A} \bar{f}_{a1}^{(l=1, j \geq 2)} \right\rangle \\
\langle \bar{\sigma}_a^{(l=1)} \rangle &= \frac{1}{T_a} n_a e_a \langle u_{\parallel a} E_{\parallel} \rangle - \frac{1}{T_a} \left\langle u_{\parallel a} \mathbf{n} \cdot (\nabla p_{a1} + \nabla \cdot \boldsymbol{\pi}_a) + \frac{2}{5} \frac{q_{\parallel a}}{p_a} \mathbf{n} \cdot (\nabla \theta_{a1} + \nabla \cdot \boldsymbol{\Theta}_a) \right\rangle \\
&\quad - \left\langle \int d^3v \frac{\bar{f}_{a1}^{(l=1, j \geq 2)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l=0)} + \bar{f}_{a1}^{(l=2)}) \right\rangle \\
\langle \bar{\sigma}_a^{(l=2)} \rangle &= -\frac{1}{T_a} \left\langle \left[ (p_{\parallel a} - p_{\perp a}) \mathbf{u}_a + (\Theta_{\parallel a} - \Theta_{\perp a}) \frac{2}{5p_a} \mathbf{q}_a \right] \cdot \nabla \ln B \right\rangle \\
&\quad - \left\langle \int d^3v \frac{\bar{f}_{a1}^{(l=2)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l=1, j \geq 2)} + \bar{f}_{a1}^{(l=3)}) \right\rangle \\
\langle \bar{\sigma}_a^{(l)} \rangle &= - \left\langle \int d^3v \frac{\bar{f}_{a1}^{(l)}}{f_{a0}} \mathcal{A}(\bar{f}_{a1}^{(l-1)} + \bar{f}_{a1}^{(l+1)}) \right\rangle \quad (\text{for } l \geq 3) \tag{D23}
\end{aligned}$$

which are summed up to

$$\begin{aligned}
\langle \bar{\sigma}_a \rangle &= \sum_{l=0}^{\infty} \langle \bar{\sigma}_a^{(l)} \rangle \\
&= \frac{1}{T_a} n_a e_a \langle u_{\parallel a} E_{\parallel} \rangle - \frac{1}{T_a} \left\langle u_{\parallel a} \mathbf{n} \cdot (\nabla p_{1a} + \nabla \cdot \boldsymbol{\pi}_a) + \frac{2}{5} \frac{q_{\parallel a}}{p_a} \mathbf{n} \cdot (\nabla \theta_{1a} + \nabla \cdot \boldsymbol{\Theta}_a) \right\rangle \\
&\quad - \frac{1}{T_a} \left\langle \left[ (p_{\parallel a} - p_{\perp a}) \mathbf{u}_a + (\Theta_{\parallel a} - \Theta_{\perp a}) \frac{2}{5p_a} \mathbf{q}_a \right] \cdot \nabla \ln B \right\rangle. \tag{D24}
\end{aligned}$$

Here, we have used the following cancellation formula for arbitrary functions  $\chi$  and  $\psi$  of  $(\mathbf{x}, v, \xi)$  :

$$\left\langle \int d^3v \frac{\chi}{f_{a0}} \mathcal{A}\psi \right\rangle + \left\langle \int d^3v \frac{\psi}{f_{a0}} \mathcal{A}\chi \right\rangle = 0 \quad (\text{D25})$$

which is derived from the following properties of  $\mathcal{A}$  :

$$\begin{aligned} \mathcal{A}(\chi\psi) &= \chi\mathcal{A}\psi + \psi\mathcal{A}\chi \\ \left\langle \int d^3v \mathcal{A}F(\mathbf{x}, v, \xi) \right\rangle &= 0 \\ \mathcal{A}G(V, v) &= 0 \end{aligned} \quad (\text{D26})$$

where  $F$  and  $G$  are arbitrary functions of  $(\mathbf{x}, v, \xi)$  and  $(V, v)$ , respectively. After some calculations, Eq. (D24) is rewritten in the same thermodynamic form as in Eq. (46) :

$$T_a \langle \tilde{\sigma}_a \rangle = J_{a1}^{\text{nc1}} X_{a1} + J_{a2}^{\text{nc1}} X_{a2} + J_{a3} X_{a3} \quad (\text{D27})$$

of which the species sum is given by

$$\sum_a T_a \langle \tilde{\sigma}_a \rangle = \sum_a (J_{a1}^{\text{nc1}} X_{a1} + J_{a2}^{\text{nc1}} X_{a2}) + J_E X_E. \quad (\text{D28})$$

It should be noted that the above neoclassical thermodynamic forms of the entropy production are derived from the kinetic form only through the magnetic surface average and sum of all the  $l$ -th tensor moment contributions of the distribution function. In Chap.17 of Ref.3, Balescu used only the vector Hermitian moments of the distribution function to calculate the kinetic form of the entropy production, which is written in our notation as

$$\sum_a T_a \left( \langle \tilde{\sigma}_a \rangle + \langle \tilde{\sigma}_a^{(l=1)} \rangle \right). \quad (\text{D29})$$

Here, as is understandable from the form of  $\tilde{f}_{a1}$  shown in Eq.(17), the contribution of the vector Hermitian moments contains  $\tilde{\sigma}_a$  which is defined by Eq.(16) and written in the classical thermodynamic form of Eq.(21). In the Pfirsch-Schlüter regime, the higher order moments representing higher anisotropy become small so that the  $l = 2$  tensor moments corresponding to the viscosities (and therefore to the banana-plateau and nonaxisymmetric transport fluxes) are smaller than the  $l = 1$  vector moment corresponding to the classical and Pfirsch-Schlüter fluxes by a small factor  $v_{Ta}\tau_{aa}/L$  ( $L$ : the scale length of the magnetic configuration). In this case, the  $l \geq 3$  moments are further smaller and the total entropy production is approximated in the lowest order of  $v_{Ta}\tau_{aa}/L$  with the 13M approximation by Eq.(D29) as

$$\begin{aligned} \sum_a T_a \left( \langle \tilde{\sigma}_a \rangle + \langle \tilde{\sigma}_a \rangle \right) &\simeq \sum_a T_a \left( \langle \tilde{\sigma}_a \rangle + \langle \tilde{\sigma}_a^{(l=1)} \rangle \right) \\ &\simeq \sum_a \left( \left( \langle J_{a1}^{\text{cl}} \rangle + J_{a1}^{\text{PS}} \right) X_{a1} + \left( \langle J_{a2}^{\text{cl}} \rangle + J_{a2}^{\text{nc1}} \right) X_{a2} \right) \\ &\quad + J_E^{\text{cl}} X_E \quad (\text{for the Pfirsch-Schlüter regime}) \end{aligned} \quad (\text{D30})$$

which takes the thermodynamic form consisting of the products of the thermodynamic forces and their conjugate classical and Pfirsch-Schlüter fluxes. (Here, it is noted that the classical fluxes  $J_{a_j}^{\text{cl}}$  ( $j = 1, 2$ ) is defined not by the magnetic surface average but by the local quantity in Eq.(22).) However, for small collision frequencies as in the plateau and banana regimes, contributions from higher order tensor moments with  $l \geq 2$  are comparable to the  $l = 1$  vector moment and are indispensable in order to derive the complete neoclassical thermodynamic form including the banana-plateau and nonaxisymmetric transport fluxes.

## References

- <sup>1</sup>F. L. Hinton and R. D. Hazeltine, *Rev. Mod. Phys.* **42**, 239 (1976).
- <sup>2</sup>S. P. Hirshman and D. J. Sigmar, *Nucl. Fusion* **21**, 1079 (1981).
- <sup>3</sup>R. Balescu, *Transport Processes in Plasmas* (2 vols.) (North-Holland, Amsterdam, 1988).
- <sup>4</sup>P. C. Liewer, *Nucl. Fusion* **25**, 543 (1985).
- <sup>5</sup>S. R. de Groot and P. Mazur, *Nonequilibrium Thermodynamics* (North-Holland, Amsterdam, 1962), Chap.4.
- <sup>6</sup>L. M. Kovrizhnykh, *Nucl. Fusion* **24**, 851 (1984).
- <sup>7</sup>K. C. Shaing and J. D. Callen, *Phys. Fluids* **26**, 3315 (1983).
- <sup>8</sup>K. C. Shaing, S. P. Hirshman, and J. D. Callen, *Phys. Fluids* **29**, 521 (1986).
- <sup>9</sup>K. C. Shaing, S. P. Hirshman, and J. S. Tolliver, *Phys. Fluids* **30**, 2548 (1987).
- <sup>10</sup>M. Coronado and H. Wobig, *Phys. Fluids* **29**, 527 (1986).
- <sup>11</sup>R. Balescu and S. Fantechi, *Phys. Fluids B* **2**, 2091 (1990).
- <sup>12</sup>H. Grad, *Phys. Fluids* **10**, 137 (1967).
- <sup>13</sup>M. N. Rosenbluth, R. D. Hazeltine, and H. L. Hinton, *Phys. Fluids* **15**, 116 (1972).
- <sup>14</sup>R. D. Hazeltine and J. D. Meiss, *Plasma Confinement* (Addison-Wesley, Redwood City, California, 1992), p.113.
- <sup>15</sup>R. D. Hazeltine, *Plasma Phys.* **15**, 77 (1973).
- <sup>16</sup>H. Sugama and W. Horton, to be published in *Phys. Plasmas* **2**, (1995).

## Recent Issues of NIFS Series

- NIFS-314 Shao-ping Zhu, R. Horiuchi, T. Sato and The Complexity Simulation Group,  
*Non-Taylor Magnetohydrodynamic Self-Organization*; Oct. 1994
- NIFS-315 M. Tanaka,  
*Collisionless Magnetic Reconnection Associated with Coalescence of Flux Bundles*; Nov. 1994
- NIFS-316 M. Tanaka,  
*Macro-EM Particle Simulation Method and A Study of Collisionless Magnetic Reconnection*; Nov. 1994
- NIFS-317 A. Fujisawa, H. Iguchi, M. Sasao and Y. Hamada,  
*Second Order Focusing Property of 210° Cylindrical Energy Analyzer*;  
Nov. 1994
- NIFS-318 T. Sato and Complexity Simulation Group,  
*Complexity in Plasma - A Grand View of Self- Organization*; Nov. 1994
- NIFS-319 Y. Todo, T. Sato, K. Watanabe, T.H. Watanabe and R. Horiuchi,  
*MHD-Vlasov Simulation of the Toroidal Alfvén Eigenmode*; Nov. 1994
- NIFS-320 A. Kageyama, T. Sato and The Complexity Simulation Group,  
*Computer Simulation of a Magnetohydrodynamic Dynamo II*: Nov. 1994
- NIFS-321 A. Bhattacharjee, T. Hayashi, C.C.Hegna, N. Nakajima and T. Sato,  
*Theory of Pressure-induced Islands and Self-healing in Three-dimensional Toroidal Magnetohydrodynamic Equilibria*; Nov. 1994
- NIFS-322 A. Iiyoshi, K. Yamazaki and the LHD Group,  
*Recent Studies of the Large Helical Device*; Nov. 1994
- NIFS-323 A. Iiyoshi and K. Yamazaki,  
*The Next Large Helical Devices*; Nov. 1994
- NIFS-324 V.D. Pustovitov  
*Quasisymmetry Equations for Conventional Stellarators*; Nov. 1994
- NIFS-325 A. Taniike, M. Sasao, Y. Hamada, J. Fujita, M. Wada,  
*The Energy Broadening Resulting from Electron Stripping Process of a Low Energy Au<sup>-</sup> Beam*; Dec. 1994
- NIFS-326 I. Viniar and S. Sudo,  
*New Pellet Production and Acceleration Technologies for High Speed Pellet Injection System "HIPEL" in Large Helical Device*; Dec. 1994
- NIFS-327 Y. Hamada, A. Nishizawa, Y. Kawasumi, K. Kawahata, K. Itoh, A. Ejiri, K. Toi, K. Narihara, K. Sato, T. Seki, H. Iguchi, A. Fujisawa, K. Adachi, S. Hidekuma, S. Hirokura, K. Ida, M. Kojima, J. Koong, R. Kumazawa,



- H. Kuramoto, R. Liang, T. Minami, H. Sakakita, M. Sasao, K.N. Sato, T. Tsuzuki, J. Xu, I. Yamada, T. Watari,  
*Fast Potential Change in Sawteeth in JIPP T-IIU Tokamak Plasmas*; Dec. 1994
- NIFS-328 V.D. Pustovitov,  
*Effect of Satellite Helical Harmonics on the Stellarator Configuration*; Dec. 1994
- NIFS-329 K. Itoh, S-I. Itoh and A. Fukuyama,  
*A Model of Sawtooth Based on the Transport Catastrophe*; Dec. 1994
- NIFS-330 K. Nagasaki, A. Ejiri,  
*Launching Conditions for Electron Cyclotron Heating in a Sheared Magnetic Field*; Jan. 1995
- NIFS-331 T.H. Watanabe, Y. Todo, R. Horiuchi, K. Watanabe, T. Sato,  
*An Advanced Electrostatic Particle Simulation Algorithm for Implicit Time Integration*; Jan. 1995
- NIFS-332 N. Bekki and T. Karakisawa,  
*Bifurcations from Periodic Solution in a Simplified Model of Two-dimensional Magnetoconvection*; Jan. 1995
- NIFS-333 K. Itoh, S.-I. Itoh, M. Yagi, A. Fukuyama,  
*Theory of Anomalous Transport in Reverse Field Pinch*; Jan. 1995
- NIFS-334 K. Nagasaki, A. Isayama and A. Ejiri  
*Application of Grating Polarizer to 106.4GHz ECH System on Heliotron-E*; Jan. 1995
- NIFS-335 H. Takamaru, T. Sato, R. Horiuchi, K. Watanabe and Complexity Simulation Group,  
*A Self-Consistent Open Boundary Model for Particle Simulation in Plasmas*; Feb. 1995
- NIFS-336 B.B. Kadomtsev,  
*Quantum Telegraph : is it possible?*; Feb. 1995
- NIFS-337 B.B.Kadomtsev,  
*Ball Lightning as Self-Organization Phenomenon*; Feb. 1995
- NIFS-338 Y. Takeiri, A. Ando, O. Kaneko, Y. Oka, K. Tsumori, R. Akiyama, E. Asano, T. Kawamoto, M. Tanaka and T. Kuroda,  
*High-Energy Acceleration of an Intense Negative Ion Beam*; Feb. 1995
- NIFS-339 K. Toi, T. Morisaki, S. Sakakibara, S. Ohdachi, T.Minami, S. Morita, H. Yamada, K. Tanaka, K. Ida, S. Okamura, A. Ejiri, H. Iguchi, K. Nishimura, K. Matsuoka, A. Ando, J. Xu, I. Yamada, K. Narihara,

- R. Akiyama, H. Idei, S. Kubo, T. Ozaki, C. Takahashi, K. Tsumori,  
*H-Mode Study in CHS*; Feb. 1995
- NIFS-340 T. Okada and H. Tazawa,  
*Filamentation Instability in a Light Ion Beam-plasma System with  
External Magnetic Field*; Feb. 1995
- NIFS-341 T. Watanabe, G. Gnudi,  
*A New Algorithm for Differential-Algebraic Equations Based on HIDM*;  
Feb. 13, 1995
- NIFS-342 Y. Nejoh,  
*New Stationary Solutions of the Nonlinear Drift Wave Equation*;  
Feb. 1995
- NIFS-343 A. Ejiri, S. Sakakibara and K. Kawahata,  
*Signal Based Mixing Analysis for the Magnetohydrodynamic Mode  
Reconstruction from Homodyne Microwave Reflectometry*; Mar.. 1995
- NIFS-344 B.B.Kadomtsev, K. Itoh, S.-I. Itoh  
*Fast Change in Core Transport after L-H Transition*; Mar. 1995
- NIFS-345 W.X. Wang, M. Okamoto, N. Nakajima and S. Murakami,  
*An Accurate Nonlinear Monte Carlo Collision Operator*; Mar. 1995
- NIFS-346 S. Sasaki, S. Takamura, S. Masuzaki, S. Watanabe, T. Kato, K. Kadota,  
*Helium I Line Intensity Ratios in a Plasma for the Diagnostics of Fusion  
Edge Plasmas*; Mar. 1995
- NIFS-347 M. Osakabe,  
*Measurement of Neutron Energy on D-T Fusion Plasma Experiments*;  
Apr. 1995
- NIFS-348 M. Sita Janaki, M.R. Gupta and Brahmananda Dasgupta,  
*Adiabatic Electron Acceleration in a Cnoidal Wave*; Apr. 1995
- NIFS-349 J. Xu, K. Ida and J. Fujita,  
*A Note for Pitch Angle Measurement of Magnetic Field in a Toroidal  
Plasma Using Motional Stark Effect*; Apr. 1995
- NIFS-350 J. Uramoto,  
*Characteristics for Metal Plate Penetration of a Low Energy Negative  
Muonlike or Pionlike Particle Beam*: Apr. 1995
- NIFS-351 J. Uramoto,  
*An Estimation of Life Time for A Low Energy Negative Pionlike Particle  
Beam*: Apr. 1995
- NIFS-352 A. Taniike,

*Energy Loss Mechanism of a Gold Ion Beam on a Tandem Acceleration System: May 1995*

- NIFS-353 A. Nishizawa, Y. Hamada, Y. Kawasumi and H. Iguchi,  
*Increase of Lifetime of Thallium Zeolite Ion Source for Single-Ended Accelerator: May 1995*
- NIFS-354 S. Murakami, N. Nakajima, S. Okamura and M. Okamoto,  
*Orbital Aspects of Reachable  $\beta$  Value in NBI Heated Heliotron/Torsatrons; May 1995*
- NIFS-355 H. Sugama and W. Horton,  
*Neoclassical and Anomalous Transport in Axisymmetric Toroidal Plasmas with Electrostatic Turbulence; May 1995*
- NIFS-356 N. Ohyabu  
*A New Boundary Control Scheme for Simultaneous Achievement of H-mode and Radiative Cooling (SHC Boundary); May 1995*
- NIFS-357 Y. Hamada, K.N. Sato, H. Sakakita, A. Nishizawa, Y. Kawasumi, R. Liang, K. Kawahata, A. Ejiri, K. Toi, K. Narihara, K. Sato, T. Seki, H. Iguchi, A. Fujisawa, K. Adachi, S. Hidekuma, S. Hirokura, K. Ida, M. Kojima, J. Koong, R. Kumazawa, H. Kuramoto, T. Minami, M. Sasao, T. Tsuzuki, J.Xu, I. Yamada, and T. Watari,  
*Large Potential Change Induced by Pellet Injection in JIPP T-IIU Tokamak Plasmas; May 1995*
- NIFS-358 M. Ida and T. Yabe,  
*Implicit CIP (Cubic-Interpolated Propagation) Method in One Dimension; May 1995*
- NIFS-359 A. Kageyama, T. Sato and The Complexity Simulation Group,  
*Computer Has Solved A Historical Puzzle: Generation of Earth's Dipole Field; June 1995*
- NIFS-360 K. Itoh, S.-I. Itoh, M. Yagi and A. Fukuyama,  
*Dynamic Structure in Self-Sustained Turbulence; June 1995*
- NIFS-361 K. Kamada, H. Kinoshita and H. Takahashi,  
*Anomalous Heat Evolution of Deuteron Implanted Al on Electron Bombardment; June 1995*
- NIFS-362 V.D. Pustovitov,  
*Suppression of Pfirsch-schlüter Current by Vertical Magnetic Field in Stellarators; June 1995*
- NIFS-363 A. Ida, H. Sanuki and J. Todoroki  
*An Extended K-dV Equation for Nonlinear Magnetosonic Wave in a Multi-Ion Plasma; June 1995*