



INTERNATIONAL ATOMIC ENERGY AGENCY

17th IAEA Fusion Energy Conference
Yokohama, Japan, 19 - 24 October 1998

IAEA-CN-69/THP1/08

NATIONAL INSTITUTE FOR FUSION SCIENCE

Global Mode Analysis of Ideal MHD Modes in $L = 2$ Heliotron/Torsatron Systems

N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto

(Received - Oct. 1, 1998)

NIFS-566

Oct. 1998

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

RESEARCH REPORT NIFS Series

This report is a preprint prepared for presentation at a scientific meeting. Because of the preliminary nature of its content, substantial changes of substance or detail may have to be made before publication. The publisher makes no claim on the understanding that it will not be cited in the literature of its subject or reproduced in any printed form. The view expressed and the statements made remain the responsibility of the author(s). The views do not necessarily reflect those of the government of the Japanese Federation. The organizations sponsoring this meeting cannot be held responsible for any errors or omissions appearing in this preprint.

NAGOYA, JAPAN



INTERNATIONAL ATOMIC ENERGY AGENCY

17th IAEA Fusion Energy Conference
Yokohama, Japan, 19 - 24 October 1998

IAEA-CN-69/THP1/08

**GLOBAL MODE ANALYSIS OF IDEAL MHD MODES
IN $L = 2$ HELIOTRON/TORSATRON SYSTEMS**

N. NAKAJIMA^{†*}, J. CHEN^{*}, K. ICHIGUCHI^{†*}, M. OKAMOTO^{†*}

[†]National Institute for Fusion Science, Toki 509-5292, Japan

^{*}Graduate University for Advanced Studies, Japan

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The view expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designated Member State(s). In particular, the organizations sponsoring this meeting cannot be held responsible for any material reproduced in this preprint.

GLOBAL MODE ANALYSIS OF IDEAL MHD MODES IN $L = 2$ HELIOTRON/TORSATRON SYSTEMS

N. NAKAJIMA^{†*}, J. CHEN^{*}, K. ICHIGUCHI^{†*}, M. OKAMOTO^{†*}

[†]National Institute for Fusion Science, Toki 509-5292, Japan

^{*}Graduate University for Advanced Studies, Japan

Abstract

By means of a global mode analysis of ideal MHD modes for Mercier-unstable equilibria in a planar axis $L = 2/M = 10$ heliotron/torsatron system with an inherently large Shafranov shift, the conjecture from local mode analysis for Mercier-unstable equilibria has been confirmed and the properties of pressure-driven modes have been clarified. According to the degree of the decrease in the local magnetic shear by the Shafranov shift, the Mercier-unstable equilibria are categorized into toroidicity-dominant (strong reduction) and helicity-dominant (weak reduction) equilibria. In both types of equilibria, interchange modes are destabilized for low toroidal mode numbers $n < M$, where M is the toroidal field period of the equilibria, and both poloidally and toroidally localized ballooning modes purely inherent to three-dimensional systems are destabilized for fairly high toroidal mode numbers $n \gg M$. For moderate toroidal mode numbers $n \sim M$, tokamak-like poloidally localized ballooning modes with a weak toroidal mode coupling are destabilized in toroidicity-dominant equilibria, and in contrast, in the helicity-dominant equilibria, interchange modes are destabilized. The interchange modes are localized on the inner side of the torus, because the Shafranov shift enhances the unfavorable magnetic curvature there rather than on the outer side of the torus. A continuous or quasi-point unstable spectrum is briefly discussed.

1. INTRODUCTION

From the local mode analysis of high-mode-number ballooning modes in an $L = 2/M = 10$ planar axis heliotron/torsatron system with an inherently large Shafranov shift (where L and M are the polarity and toroidal field period of the helical coils, respectively)[1], it was previously conjectured[2] that the global structure of pressure-driven modes for Mercier-unstable equilibria would have the following properties: Tokamak-like poloidally localized ballooning modes or interchange modes appear when their typical toroidal mode numbers are relatively low. As the typical toroidal mode numbers become higher, ballooning modes inherent to three-dimensional systems appear with larger growth rates and localized in both the poloidal and toroidal directions.

The purposes of this work are to prove the above conjecture and to clarify the inherent properties of pressure-driven modes, through a global mode analysis of the ideal MHD modes.

2. CATEGORIZATION OF MERCIER-UNSTABLE EQUILIBRIA

The changes in the local magnetic shear and the normal magnetic curvature by the Shafranov shift are related to toroidicity. The Shafranov shift decreases the local magnetic shear on the outside of the torus, leading to the reduction of the field line bending stabilizing effect on ballooning modes[1]. On the other hand, the Shafranov shift enhances (reduces) the local unfavorable normal magnetic curvature on the inner (outer) side of the torus. According to the degree of the decrease in the local magnetic shear by the Shafranov shift, the Mercier-unstable equilibria can be categorized into two types, namely, toroidicity-dominant and helicity-dominant equilibria. The former (latter) equilibria are characterized by properties that the local magnetic shear is strongly (weakly) reduced by the Shafranov shift, so that ballooning modes are easy (difficult) to destabilize. The former (latter) equilibria are basically created with peaked (broad) pressure profiles under the currentless condition.

3. GLOBAL MODE ANALYSIS IN MERCIER-UNSTABLE EQUILIBRIA

The global mode analysis of ideal MHD modes in Mercier-unstable equilibria is done by using the CAS3D2MN code[3] with the shift-and-invert Lanczos algorithm[4]. The three-dimensional equilibria considered here have toroidal field period M , and this is what mainly determines the toroidal period of the local magnetic curvature due to helicity. Therefore, we investigate the properties of pressure-driven modes for various cases of relative magnitude between the typical toroidal mode number n of the perturbation and the toroidal period M of the local magnetic curvature due to helicity: namely, $n < M$, $n \sim M$, and $n \gg M$.

For low toroidal mode numbers $n < M$, interchange modes occur, which feel the average magnetic curvature, for both types of Mercier-unstable equilibria. One of them is shown in Fig. 1.




Fig. 1 (a) $\vec{\xi} \cdot \nabla \psi$ vs. r_N (normalized radial coordinate) with the dominant toroidal mode number, and the corresponding contours of $\tilde{P} = -\nabla P \cdot \vec{\xi}$ on the vertically (b) and horizontally (c) elongated poloidal cross sections.

The radial distribution of the Fourier components of the normal displacement $\vec{\xi} \cdot \nabla \psi$ is shown in Fig. 1(a) with the origin of the poloidal angle on the inner side of the torus. Three resonant mode structures with $n = 4$ are visible and the amplitudes of modes with different toroidal mode numbers are quite small. The mode structure is similar to that of a ballooning mode except that each Fourier mode has both positive and negative parts, which means that the perturbed pressure $\tilde{P} = -\nabla P \cdot \vec{\xi}$ due to this type of interchange mode has a tendency to extend radially on the inner side of the torus and to change phase in the radial direction on the outer side of the torus through poloidal mode coupling as shown in Figs. 1(b) and (c). In other words, this type of interchange mode is anti-ballooning with respect to the poloidal mode coupling. This is because the normal magnetic curvature is more unfavorable on the inner side of the torus than on the outer side of the torus by the Shafranov shift.

For moderate toroidal mode numbers $n \sim M$, the modes begin to feel the local structure of the magnetic curvature. Since the Shafranov shift strongly reduces the stabilizing effects due to the local magnetic shear on the outer side of the torus in the toroidicity-dominant Mercier-unstable equilibrium, tokamak-like poloidally localized ballooning modes with weak toroidal mode coupling occur. This is shown in Fig. 2 with the origin of the poloidal angle on the outer side of the torus. The typical toroidal mode numbers are still so small that the modes can not feel the local structure of the magnetic curvature effectively, and hence the toroidal mode coupling is weak. Three groups of Fourier modes for the normal displacement $\vec{\xi} \cdot \nabla \psi$ with different toroidal mode numbers are visible, namely, $n = 22$, $n = 32$, and $n = 42$ in Fig. 2(a). Each group, however, consists of many Fourier modes with different poloidal mode numbers caused by the

poloidal mode coupling, so that the structure of each group due to the poloidal mode coupling is quite similar to that of ballooning modes in tokamak plasmas. The corresponding contours of the perturbed pressure $\tilde{P} = -\nabla P \cdot \tilde{\xi}$ are shown in Figs. 2(b) and (c).




Fig. 2 The same quantities as in Fig. 1 for $n \sim M$ in a toroidicity-dominant equilibrium.

In contrast, for the helicity-dominant equilibria with a weak reduction of the stabilizing term of ballooning modes by the Shafranov shift, interchange modes still occur as shown in Fig. 3. The toroidal mode coupling of interchange modes becomes stronger as the toroidal mode number increases, as shown in Fig. 3(a) where the origin of the poloidal angle is on the inner side of the torus. Just as for interchange modes with $n < M$, the interchange modes with $n \sim M$ also have a tendency to be radially extended on the inner side of the torus and to change phase in the radial direction on the outer side of the torus, as shown in Figs. 3(b) and (c).




Fig. 3 The same quantities as in Fig. 1 for $n \sim M$ in a helicity-dominant equilibrium.

For fairly high toroidal mode numbers $n \gg M$, the modes can easily distinguish the local fine structure of the magnetic curvature. This results in the appearance of ballooning modes inherent to three-dimensional systems for both types of Mercier-unstable equilibria. These ballooning modes have such strong poloidal and toroidal mode couplings as to localize in both the poloidal and toroidal directions as shown in Fig. 4. There are eight groups of Fourier modes for $\tilde{\xi} \cdot \nabla \psi$ with different toroidal mode numbers as shown in Fig. 4(a), where the origin of the poloidal angle is on the outer side of the torus. Neighboring groups of Fourier modes have opposite phase to each other, just as in Fig. 2(a). This relative phase difference of the neighboring groups leads to the clear localization of the perturbed pressure in the toroidal direction, as shown in Figs. 4(b) and (c). On the outer side of the torus, the perturbed pressure, which localizes on the horizontally elongated poloidal cross section with the locally unfavorable magnetic curvature at the outside of the torus, almost disappears on the vertically elongated poloidal cross section with the locally favorable magnetic curvature at the outside of the torus. Moreover, the strong

toroidal mode coupling causes a type of poloidal localization that is different from the kind only due to poloidal mode coupling as shown in Fig. 4(d). It shows the corresponding contours of the perturbed pressure on the (θ, ζ) plane with one period in the poloidal direction and one-tenth of a period (one field period) in the toroidal direction at $r_N = 0.734$. In Fig. 4(d), it can be seen that regions where the perturbed pressure has large amplitude (indicated by dark diagonal stripes) alternate with regions of quite small amplitude (denoted by white diagonal stripes). Judging from the value of the rotational transform on this flux surface, namely, $\iota = 0.58$, we conclude that these high-amplitude and low-amplitude stripes are aligned along magnetic field lines, and that the strong toroidal mode coupling in addition to the poloidal mode coupling makes the perturbation be localized on selected flux tubes.




Fig. 4 (a) $\vec{\xi} \cdot \nabla \psi$ vs. r_N with the dominant toroidal mode numbers, and the corresponding contours of $\bar{P} = -\nabla P \cdot \vec{\xi}$ on the vertically (b) and horizontally (c) elongated poloidal cross sections, and (d) on the (θ, ζ) plane at $r_N = 0.734$, where the Fourier mode with $n = 208$ has its maximum amplitude.

4. DISCUSSIONS

Since the interchange modes basically localize along mode rational magnetic field lines driven by the average unfavorable magnetic curvature, the toroidal mode coupling merely influences the localization in the toroidal direction but does not have an essential effect on the magnitude of the eigenvalues. This may lead to the existence of a narrow continuous or quasi-point unstable spectrum. In contrast, the ballooning modes basically localize near the locally unfavorable magnetic curvature, so that toroidal mode coupling significantly influences the eigenvalues. Thus, ballooning modes may be unable to have a continuous or quasi-point unstable spectrum, except perhaps for case with extremely high toroidal mode numbers ($n \rightarrow \infty$).

REFERENCES

- ¹ NAKAJIMA, N., Phys. Plasmas **3** (1996) 4545.
- ² NAKAJIMA, N., Phys. Plasmas **3** (1996) 4556.
- ³ SCHWAB, C., Phys. Fluids B **5** (1993) 3195.
- ⁴ CHEN, J., NAKAJIMA, N., and OKAMOTO, M., Comput. Phys. Commun., *to be published*.

Recent Issues of NIFS Series

- NIFS-532 J. Uramoto,
Pair Creations of Negative and Positive Pionlike (Muonlike) Particle or K Mesonlike (Muonlike) Particle in H₂ or D₂ Gas Discharge in Magnetic Field; Dec 1997
- NIFS-533 S. Kawata, C. Boonmee, T. Teramoto, L. Drska, J. Limpouch, R. Liska, M. Sinor,
Computer-Assisted Particle-in-Cell Code Development; Dec 1997
- NIFS-534 Y. Matsukawa, T. Suda, S. Ohnuki and C. Namba,
Microstructure and Mechanical Property of Neutron Irradiated TiNi Shape Memory Alloy; Jan 1998
- NIFS-535 A. Fujisawa, H. Iguchi, H. Idei, S. Kubo, K. Matsuoka, S. Okamura, K. Tanaka, T. Minami, S. Ohdachi, S. Morita, H. Zushi, S. Lee, M. Osakabe, R. Akiyama, Y. Yoshimura, K. Toi, H. Sanuki, K. Itoh, A. Shimizu, S. Takagi, A. Ejiri, C. Takahashi, M. Kojima, S. Hidekuma, K. Ida, S. Nishimura, N. Inoue, R. Sakamoto, S.-I. Itoh, Y. Hamada, M. Fujiwara,
Discovery of Electric Pulsation in a Toroidal Helical Plasma, Jan 1998
- NIFS-536 Lj R. Hadzievski, M.M. Skonc, M. Kono and T. Sato,
Simulation of Weak and Strong Langmuir Collapse Regimes, Jan. 1998
- NIFS-537 H. Sugama, W. Horton,
Nonlinear Electromagnetic Gyrokinetic Equation for Plasmas with Large Mean Flows; Feb. 1998
- NIFS-538 H. Iguchi, T.P. Crowley, A. Fujisawa, S. Lee, K. Tanaka, T. Minami, S. Nishimura, K. Ida, R. Akiyama, Y. Hamada, H., Idei, M. Isobe, M. Kojima, S. Kubo, S. Morita, S. Ohdachi, S. Okamura, M. Osakabe, K. Matsuoka, C. Takahashi and K. Toi,
Space Potential Fluctuations during MHD Activities in the Compact Helical System (CHS); Feb 1998
- NIFS-539 Takashi Yabe and Yan Zhang,
Effect of Ambient Gas on Three-Dimensional Breakup in Coronet Formation Process; Feb. 1998
- NIFS-540 H. Nakamura, K. Ikeda and S. Yamaguchi,
Transport Coefficients of InSb in a Strong Magnetic Field, Feb 1998
- NIFS-541 J. Uramoto,
Development of v_{μ} Beam Detector and Large Area v_{μ} Beam Source by H₂ Gas Discharge (I); Mar 1998
- NIFS-542 J. Uramoto,
Development of \bar{v}_{μ} Beam Detector and Large Area \bar{v}_{μ} Beam Source by H₂ Gas Discharge (II); Mar. 1998
- NIFS-543 J. Uramoto,
Some Problems inside a Mass Analyzer for Pions Extracted from a H₂ Gas Discharge; Mar 1998
- NIFS-544 J. Uramoto,
Simplified v_{μ} \bar{v}_{μ} Beam Detector and v_{μ} \bar{v}_{μ} Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch; Mar 1998
- NIFS-545 J. Uramoto,
Various Neutrino Beams Generated by D₂ Gas Discharge; Mar 1998
- NIFS-546 R. Kanno, N. Nakajima, T. Hayashi and M. Okamoto,
Computational Study of Three Dimensional Equilibria with the Bootstrap Current; Mar. 1998
- NIFS-547 R. Kanno, N. Nakajima and M. Okamoto,
Electron Heat Transport in a Self-Similar Structure of Magnetic Islands; Apr 1998
- NIFS-548 J.E. Rice,
Simulated Impurity Transport in LHD from MIST; May 1998
- NIFS-549 M.M. Skonc, T. Sato, A.M. Maluckov and M.S. Jovanovic,
On Kinetic Complexity in a Three-Wave Interaction; June 1998

- NIFS-550 S. Goto and S. Kida,
Passive Saclar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation; June 1998
- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,
Initial Value Problem of the Toroidal Ion Temperature Gradient Mode ; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watari,
Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device ; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,
Oronization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-: July 1998
- NIFS-554 S. Murakami and H. Saleem,
Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations; July 1998
- NIFS-555 H. Nakamura, K. Ikeda and S. Yamaguchi,
Physical Model of Nernst Element; Aug. 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,
Numerical Computation of Thermoelectric and Thermomagnetic Effects; Aug. 1998
- NIFS-557 T. Yasuhiko, O. Masaki, T. Katsuyoshi, O. Yoshihide, K. Osamu, A. Eiji, K. Toshikazu, A. Ryuichi and T. Masanobu,
Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System; Aug 1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Structure Organization of Polyampholytes; Sep. 1998
- NIFS-559 R. Horiuchi, K. Nishimura and T. Watanabe,
Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations; Sep. 1998
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Kholopenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,
Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection; Oct. 1998
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,
Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron; Oct. 1998
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,
Next Generation Toroidal Devices; Oct 1998
- NIFS-563 Y. Todo and T. Sato,
Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes; Oct. 1998
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watari, T. Shimozuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Idei, Y. Oka, M. Osakabe, T. Seki, K. Tsumon, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Shrimpo, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Morimoto, T. Bigelow and Y.P. Zhao,
Steady State Heating Technology Development for LHD; Oct. 1998
(IAEA-CN-69/FTP/21)
- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyabu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda,
LHD-Type Compact Helical Reactors; Oct. 1998
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems; Oct. 1998
(IAEA-CN-69/THP1/08)