

§4. Simulations of Large-Amplitude Plasma Waves and Particle Acceleration by Use of SNET

Ohsawa, Y., Toida, M., Sato, M., Takahashi, S. (Nagoya Univ.),
Ishiguro, S.

By using SNET and super computers at NIFS, we have studied the behavior of large-amplitude plasma waves and strong particle acceleration processes caused by these waves. The subjects we have studied are 1) ultrarelativistic electron acceleration in large-amplitude Alfvén wave packets [1], 2) parallel electric field in nonlinear magnetosonic waves in a three-component plasma consisting of electrons, positrons, and ions [2], 3) shock formation processes in a collision of two plasmas in an external magnetic field [3], 4) evolution of cylindrical shock wave and the trapping and acceleration of electrons in it [4], and 5) effect of ion composition on the ion acceleration in a shock wave in a two-ion-species plasma [5]. For the investigations of No. 1 – No. 3, we have used a one-dimensional, fully kinetic, relativistic, electromagnetic particle simulation code. For No. 4 and No. 5, respectively, we have used two-dimensional and three-dimensional codes.

Figure 1 is a result of the study No. 2; parallel electric field in nonlinear magnetosonic waves in a three component plasma. This figure shows the integral of the parallel electric field along the magnetic field, F , in a magnetosonic shock wave as a function of the positron-to-electron density ratio n_{p0}/n_{e0} . As n_{p0}/n_{e0} decreases, the magnitude of F decreases, and F becomes zero in a pure electron-positron plasma ($n_{p0}/n_{e0}=1$). This explains the simulation result [6] that the positron acceleration is stronger in a plasma with a lower positron density.




Fig. 1. Magnitude of F as a function of the positron-to-electron density ratio n_{p0}/n_{e0} . The dots and solid line represent the simulation result and theoretical prediction, respectively, and ϵ is the wave amplitude.




Fig. 2. Ion phase spaces (x, v_x) and (x, v_y) for a shock wave in a D-T plasma. Field profiles are also shown.

Figure 2 is a result of three-dimensional simulations with the number of simulation particles $N_i=N_e=10^9$ for the study No. 5; effect of ion composition on the ion acceleration in a shock wave in a two-ion-species plasma. This figure shows ion phase spaces for a shock wave in a deuterium-tritium (D-T) plasma. From the simulations for H-T plasmas and for D-T plasmas, we have found that the frequency difference $\Delta\omega$ between the cutoff frequency of the high-frequency mode of the magnetosonic wave and the resonance frequency of the low-frequency mode plays an important role in the wave evolution and thus in the acceleration of particles. For a D-T plasma with $n_D=n_T$, in which $\Delta\omega$ is smaller than in H-T plasmas, we have a high-frequency-mode shock wave. All the T ions are accelerated by the transverse electric field, and a great number of D ions gain energy from the longitudinal electric field.

- [1] M. Sato and Y. Ohsawa: J. Phys. Soc. Jpn. **77**, 084502 (2008).
- [2] S. Takahashi, M. Sato, and Y. Ohsawa: Phys. Plasmas **15**, 082309 (2008).
- [3] K. Yamauchi and Y. Ohsawa: J. Phys. Soc. Jpn. **77**, 054501 (2008).
- [4] M. Toida, M. Ueno, and Y. Ohsawa: J. Phys. Soc. Jpn. **77**, 084501 (2008).
- [5] M. Toida, H. Higashino, and Y. Ohsawa: Phys. Plasmas **15**, 092305 (2008).
- [6] H. Hasegawa, S. Usami, and Y. Ohsawa: Phys. Plasmas **10**, 3455 (2003); H. Hasegawa and Y. Ohsawa, *ibid.* **12**, 012312 (2005).