

Nonlinear competition between zonal flows and geodesic acoustic modes

K. Itoh¹, S.-I. Itoh², M. Yagi², S. Toda¹, A. Fujisawa¹, N. Kasuya¹, A. Fukuyama³

¹National Institute for Fusion Science, Toki 509-5292, Japan

²Research Institute for Applied Mechanics, Kyushu University, Kasuga 816-8580, Japan

³Department of Nuclear Engineering, Kyoto University, Kyoto, 606-8501, Japan

It has been known that the zonal flows (ZFs) and geodesic acoustic modes (GAMs) are destabilized by microscopic fluctuations in the range of drift wave frequencies [1]. The rates, at which ZFs and GAMs are driven by background microscopic turbulence, have been derived [1,2]. The nonlinear saturation of the Reynolds stress, when the ZFs have finite amplitude, was also derived [3]. That is, when the induced flow has finite amplitude, the drift wave spectrum is deformed so that the energy input from turbulence to the flow is suppressed. This result was derived by renormalizing the higher order kinetic of quasiparticle response. Under the presence of ZFs and GAMs, the deformation of drift wave spectrum causes the modification of the driving rates of ZFs and GAMs. Thus, the nonlinear competition between ZFs and GAMs occurs. The turbulent transport is strongly affected by the inclusion of zonal flows. Thus, the competition between ZFs and GAMs must be taken into account, in order to analyze the turbulent transport in toroidal plasmas.

In order to study the competition between the ZFs and GAMs, the effect of the return flow along the magnetic field line must be taken into account. The coupling of the poloidal flow to the parallel flow enhances the effective inertia of plasma [4]. For ZFs, one has an enhancement factor for the effective inertia $1 + 2q^2$ (in the plateau regime). The back interaction of the ZFs and GAMs on drift wave turbulence closes the set of coupled equations, which determines the level of turbulence and turbulent transport. The self-consistent solution of the microscopic fluctuations, ZFs and GAMs is obtained. The competition between ZFs and GAMs is summarized in the parameter space of the damping rates.

[1] P H Diamond, S.-I. Itoh, K. Itoh, T. S. Hahm: Plasma Phys. Control. Fusion **47** (2005) R35

[2] K. Itoh, K. Hallatschek, S.-I. Itoh: Plasma Phys. Control. Fusion **47** (2005) 451

[3] K. Itoh, et al.: Phys. Plasmas **12** (2005) 062303

[4] K. Itoh and S.-I. Itoh: Plasma Phys. Control. Fusion **38** (1996) 1

This work was partly supported by Grants-in-Aid for Specially-Promoted Research (16002005) and for Scientific Research (19360418).