

Transport analysis of the LHD plasma using the integrated code: TASK3D

A. Wakasa, A. Fukuyama, S. Murakami, C.D. Beidler^{a)}, H. Maassberg^{a)},
M. Yokoyama^{b)}, and M. Sato^{b)}

Department of Nuclear Engineering, Kyoto University, Kyoto 606-8501, Japan
^{a)} *Max-Planck-Institute für Plasmaphysik, EURATOM Ass., Greifswald, Germany*
^{b)} *National Institute for Fusion Science, 322-6 Oroshi-cho, Toki 509-5292, Japan*

wakasa@p-grp.nucleng.kyoto-u.ac.jp

The integrated simulation code in helical plasmas, TASK3D[1], is being developed on the basis of an integrated modeling code for tokamak plasma, TASK[2]. In helical systems, the neoclassical transport is one of the important issues in addition to the anomalous transport, because of strong temperature dependence of heat conductivity and an important role in determining the radial electric field. We have already constructed the neoclassical transport database in LHD, DGN/LHD[3]. The mono-energetic diffusion coefficients are evaluated based on the Monte Carlo method by DCOM code and the mono-energetic diffusion coefficients database is constructed using a neural network technique. Also we apply GSRAKE[4] code, which solves the ripple-averaged drift kinetic equation, to obtain transport coefficients in highly collisionless regime. We have newly incorporated the DGN/LHD module into TASK3D. We will present several results of transport simulation in typical LHD plasmas.

[1] M. Sato, *et al.*, Plasma and Fusion Res. Vol. 3 (2008) S1063.

[2] A. Fukuyama, *et al.*, Proc. of 20th IAEA Fusion Energy Conf. (Villamoura, Portugal, 2004) IAEA-CSP-25/CD/TH/P2-3.

[3] A. Wakasa, *et al.*, Plasma and Fusion Res. Vol. 3 (2008) S1030.

[4] C. D. Beidler, *et al.*, Plasma Phy. Control. Fusion **37** (1995) 463