



INTERNATIONAL ATOMIC ENERGY AGENCY

**FIFTEENTH INTERNATIONAL CONFERENCE ON PLASMA PHYSICS  
AND CONTROLLED NUCLEAR FUSION RESEARCH**

Seville, Spain, 26 September – 1 October 1994

IAEA-CN-60/D-2-II-4

**NATIONAL INSTITUTE FOR FUSION SCIENCE**
**Behavior of Magnetic Islands in 3D MHD  
Equilibria of Helical Devices**

T. Hayashi, T. Sato, N. Nakajima, K. Ichiguchi, P. Merkel,  
J. Nührenberg, U. Schwenn, H. Gardner,  
A. Bhattacharjee and C.C.Hegna

(Received - Aug. 26, 1994 )

NIFS-305

Sep. 1994

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

**RESEARCH REPORT  
NIFS Series**

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of content or of text may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designating Member State(s) or of the designating organization(s). In particular, neither the IAEA nor any other organization or body sponsoring this meeting can be held responsible for any material reproduced in this preprint.

NAGOYA, JAPAN



INTERNATIONAL ATOMIC ENERGY AGENCY

**FIFTEENTH INTERNATIONAL CONFERENCE ON PLASMA PHYSICS  
AND CONTROLLED NUCLEAR FUSION RESEARCH**

Seville, Spain, 26 September – 1 October 1994

---

**IAEA-CN-60/  
D-2-II-4**

**Behavior of Magnetic Islands  
in 3D MHD Equilibria of Helical Devices**

**T.Hayashi, T.Sato, N.Nakajima, K.Ichiguchi**

*National Institute for Fusion Science, Nagoya 464-01, Japan*

**P.Merkel, J.Nührenberg, U.Schwenn**

*Max-Planck-Institut für Plasmaphysik, Garching bei München, Germany*

**H.Gardner**

*Australian National University, Canberra, ACT 0200, Australia*

**A.Bhattacharjee**

*University of Iowa, Iowa City, IA 52242, USA*

**C.C.Hegna**

*University of Wisconsin-Madison, Wisconsin 53706-1687, USA*

**key words:** helical system, heliac, heliotron, torsatron, stellarator, three dimensional equilibrium, beta limit, magnetic island, self-healing of magnetic island, finite pressure effect

## Behavior of Magnetic Islands in 3D MHD Equilibria of Helical Devices

### Abstract

Magnetic island formation in three-dimensional finite- $\beta$  equilibria in the H-1 Helic is studied by using the HINT code. It is found that the size of a dangerous island should increase with  $\beta$  but that a destruction of the equilibrium at low  $\beta$  is avoided because the rotational transform evolves to exclude the rational surface concerned. At higher  $\beta$  there is evidence of near-resonant flux surface deformations which may lead to an equilibrium limit. A reconnected equilibrium at still higher  $\beta$  exhibits a double island structure which is similar to homoclinic phase portraits which have been observed after separatrix reconnection in Hamiltonian systems. Physical mechanism of the island formation in finite- $\beta$  helical equilibria is investigated to confirm there are cases where the global effect of the Pfirsch-Schlüter currents is important. The earlier theory is extended to elucidate the occurrence of the complete self-healing of island when the resistive interchange criterion satisfied.

### 1. Evolution of magnetic islands in a Helic

One of the critical issues in obtaining a high- $\beta$  plasma for helical devices is the formation of magnetic islands induced in nonaxisymmetric finite- $\beta$  equilibria. This issue has been studied by the 3D equilibrium code HINT, which does not a-priori demand the existence of regular nested magnetic surfaces. It has been discovered numerically that breaking of magnetic surfaces due to the finite- $\beta$  effect occurs in a practical configuration of a toroidal helical system, such as heliotron/torsatron [1] and Helic [2], and often imposes severer limitation on the equilibrium beta than the Shafranov shift. In addition, unexpectedly good aspect, which can be called “self-healing”, has also been discovered.

The island issue is especially the case for the Helic configuration, which has a small shear and contains a large number of Fourier harmonics in the magnetic field structure. In this paper, properties of magnetic islands induced due to finite

pressure effects in a H-1 Helic configuration [3] are studied using HINT. H-1 fields have a large degree of flexibility due to an  $l = 1$  helical trim coil, however, this flexibility does not seem to allow varying the phase of the major, stellarator-symmetric island chains. In this study we examine a configuration which includes the island structure corresponding to the  $\iota = 6/5$  resonance ( $\iota = 2/5$  per field period) in the vacuum field and assume the existence of stellarator symmetry. A sequence of results from these simulations are shown in Fig. 1 where one cross section of the H-1 flux surfaces is shown at various plasma  $\beta$ 's starting with Fig. 1(a) for the vacuum field. The  $\iota$  profile for the vacuum field is monotonous and the shear is quite small (Fig. 1(b)).

As the increase of  $\beta$ , minimum of  $\iota$  (vanishing shear) appears at around half the minor radius. As is shown in the following, depending on the value of the minimum  $\iota$  with respect to  $6/5$ , the most dangerous (low order) rational surface relevant, a variety of topology appears for magnetic islands induced in finite  $\beta$  equilibria. Figure 1(c), at  $\beta_0 = 1.5\%$  ( $\langle\beta\rangle \approx 0.5\%$ ), shows that the size of the  $\iota = 6/5$  island should increase with plasma pressure to about  $1/5$  of the plasma radius. If the pressure is increased to  $\beta_0 = 2.0\%$ , in Fig. 1(d), the plasma appears to experience a form of self-healing, however the corresponding  $\iota$  profile shows that this is because the  $6/5$  surface has been excluded from the configuration. It is possible that this exclusion might be due to some self-consistent plasma dynamics and, as such, could be considered as a (new) form of self-healing. In any case, it is a favorable property of the H-1 configuration that reasonably smooth flux surfaces can be obtained at these plasma  $\beta$ 's.

As the pressure is increased further the HINT results show that the minimum value of  $\iota$  remains just above  $6/5$ , however the flux surfaces show marked signs of *near-resonant* deformation (Fig. 1(e)). Figure 1(f) shows the flux surfaces when the equilibrium pressure has increased to  $\beta_0 = 4.4\%$ . The flux surfaces have reconnected to form a double island structure, indicating that the minimum of  $\iota$  has decreased down to  $6/5$ . The topology of the island is analogous to the homoclinic phase portraits observed for Hamiltonian dynamical systems after separatrix reconnection [4], but appears to be new in the context of stellarator magnetic fields. The possibility of separatrix reconnection occurring is due to the corresponding Hamiltonian being degenerate, which, in our case, corresponds to  $\iota' = 0$  across a finite fraction of the plasma radius.

The behavior of islands shown in Fig. 1(f) suggests an existence of a new type of self-organizing process, which is driven by a pressure-driven instability. Based on an equilibrium obtained by HINT, nonlinear simulations are also studied, where a compressible full MHD model is assumed in a 3D geometry. Excitation of strong pressure driven instability with global structure is observed as an increase in  $\beta$ . Interestingly, a tendency is found that the pressure profile spontaneously relaxes to a simple confined structure after a disordered state driven by the instability.

## 2. Physical mechanisms of formation and ‘self-healing’ of magnetic islands

A comparative study is given of some of the computational and analytical results presently available for calculating pressure-induced magnetic islands in 3D equilibria of currentless stellarators. The main goal of this study is to clarify the dominant physical mechanism of island formation, either the ‘local’ (current sheets) or the ‘global’ (Pfirsch-Schlüter currents) effects of the plasma current. For this purpose, we examine the behavior of  $D_R$  (the criteria for the local resistive interchange mode) and the Jacobian  $\sqrt{g_{mn}}$  for results obtained by HINT for the self-healing case of Helias [2].

We find the process can be consistently explained by the ‘global’ effect by supposing that the self-healing process originates from the property that the pressure-induced islands always have the same phase independent of  $\beta$  (‘fixed-phase’ property). This property has been discovered numerically; the phase of the pressure-induced islands is determined independent of the phase of the vacuum islands. The field generated by global effects of the Pfirsch-Schlüter current profile integrated over the whole plasma volume appears to have a specific simple global spatial structure. A coupling of this global-structure field with the non-axisymmetric configuration results in the robust ‘fixed-phase’ property of the pressure-induced islands. Moreover, we analyze the consistency of the equilibrium on the rational magnetic surface  $5/6$  which exists in the Helias configuration for vanishing island thickness. We find that the plasma deforms itself so that the so-called Hamada condition, that  $\int dl/B$  be constant on a rational surface, is satisfied with very good numerical accuracy when the island vanishes completely due to self-healing. The behavior described above for a Helias configuration occurs with the resistive interchange criterion being satisfied. The importance of the global effect is further confirmed by a computation for resistive-interchange

unstable equilibria; we observe the similar process of the self-healing.

On the other hand, in order to elucidate the self-healing process, the earlier theory [5] is extended to include a small vacuum island which may, in general, have different phases than pressure-induced islands. For a negative (stable)  $D_R$  case, the extended theory predicts complete self-healing at a critical value of beta, when the vacuum perturbation and the perturbation caused by the Pfirsch-Schlüter currents are exactly out of phase. Beyond this critical value, the islands reappear but with a flipped phase. This behavior is consistent with the computational results.

An apparent discrepancy remains between computations and analysis for a positive  $D_R$  case. It brings forward following problems to be clarified in the future. It certainly has to be appreciated that for a 3D computation like HINT it is very difficult to achieve the needed accuracy to resolve current sheet investigated by the earlier theories. On the other hand, consistent quantitative estimation of the amplitude as well as the phase of the global effect without assumptions for simplification is possible only by a 3D computation.

## Acknowledgements

The authors wish to thank J.D. Hanson for bringing Ref. [4] to their notice.

## References

- [1 ] HAYASHI, T., TAKEI, A., SATO, T., Phys. Fluids B **4** (1992) 1539.
- [2 ] HAYASHI, T., SATO, T., LOTZ, W., MERKEL, P., NÜHRENBERG, J., SCHWENN, U., STRUMBERGER, E., in Plasma Physics and Controlled Nuclear Fusion Research Vol.2 (IAEA, Vienna, 1993) p. 29.
- [3 ] HAMBERGER, S.M., BLACKWELL, B.D., SHARP, L.E., SHENTON, D.B., Fusion. Technol. **17** (1990) 123.
- [4 ] DEL-CASTILLO-NEGRETE, D., MORRISON, P.J., Phys. Fluids A **5** (1993) 948.
- [5 ] HEGNA, C.C., BHATTACHARJEE, A., Phys. Fluids B **1** (1989) 392.

## FIGURE CAPTION

**Fig. 1** Evolution of magnetic islands and deformation of magnetic surfaces in a Heliac (a) vacuum field, (b) rotational transform profile for vacuum field, (c)  $\beta_0 = 1.5\%$  , (d)  $\beta_0 = 2.0\%$ , (e)  $\beta_0 = 3.8\%$ , (f)  $\beta_0 = 4.4\%$ .




Fig.1



## Recent Issues of NIFS Series

- NIFS-267 H. Iguchi, K. Ida, H. Yamada, K. Itoh, S.-I. Itoh, K. Matsuoka, S. Okamura, H. Sanuki, I. Yamada, H. Takenaga, K. Uchino, K. Muraoka, *The Effect of Magnetic Field Configuration on Particle Pinch Velocity in Compact Helical System (CHS)*; Jan. 1994
- NIFS-268 T. Shikama, C. Namba, M. Kosuda, Y. Maeda, *Development of High Time-Resolution Laser Flash Equipment for Thermal Diffusivity Measurements Using Miniature-Size Specimens*; Jan. 1994
- NIFS-269 T. Hayashi, T. Sato, P. Merkel, J. Nührenberg, U. Schwenn, *Formation and 'Self-Healing' of Magnetic Islands in Finite- $\beta$  Helias Equilibria*; Jan. 1994
- NIFS-270 S. Murakami, M. Okamoto, N. Nakajima, T. Mutoh, *Efficiencies of the ICRF Minority Heating in the CHS and LHD Plasmas*; Jan. 1994
- NIFS-271 Y. Nejoh, H. Sanuki, *Large Amplitude Langmuir and Ion-Acoustic Waves in a Relativistic Two-Fluid Plasma*; Feb. 1994
- NIFS-272 A. Fujisawa, H. Iguchi, A. Taniike, M. Sasao, Y. Hamada, *A 6MeV Heavy Ion Beam Probe for the Large Helical Device*; Feb. 1994
- NIFS-273 Y. Hamada, A. Nishizawa, Y. Kawasumi, K. Narihara, K. Sato, T. Seki, K. Toi, H. Iguchi, A. Fujisawa, K. Adachi, A. Ejiri, S. Hidekuma, S. Hirokura, K. Ida, J. Koong, K. Kawahata, M. Kojima, R. Kumazawa, H. Kuramoto, R. Liang, H. Sakakita, M. Sasao, K. N. Sato, T. Tsuzuki, J. Xu, I. Yamada, T. Watari, I. Negi, *Measurement of Profiles of the Space Potential in JIPP T-IIU Tokamak Plasmas by Slow Poloidal and Fast Toroidal Sweeps of a Heavy Ion Beam*; Feb. 1994
- NIFS-274 M. Tanaka, *A Mechanism of Collisionless Magnetic Reconnection*; Mar. 1994
- NIFS-275 A. Fukuyama, K. Itoh, S.-I. Itoh, M. Yagi and M. Azumi, *Isotope Effect on Confinement in DT Plasmas*; Mar. 1994
- NIFS-276 R.V. Reddy, K. Watanabe, T. Sato and T.H. Watanabe, *Impulsive Alfvén Coupling between the Magnetosphere and Ionosphere*; Apr. 1994
- NIFS-277 J. Uramoto,

*A Possibility of  $\pi^-$  Meson Production by a Low Energy Electron Bunch and Positive Ion Bunch; Apr. 1994*

NIFS-278 K. Itoh, S.-I. Itoh, A. Fukuyama, M. Yagi and M. Azumi,  
*Self-sustained Turbulence and L-mode Confinement in Toroidal Plasmas II; Apr. 1994*

NIFS-279 K. Yamazaki and K.Y.Watanabe,  
*New Modular Heliotron System Compatible with Closed Helical Divertor and Good Plasma Confinement; Apr. 1994*

NIFS-280 S. Okamura, K. Matsuoka, K. Nishimura, K. Tsumori, R. Akiyama, S. Sakakibara, H. Yamada, S. Morita, T. Morisaki, N. Nakajima, K. Tanaka, J. Xu, K. Ida, H. Iguchi, A. Lazaros, T. Ozaki, H. Arimoto, A. Ejiri, M. Fujiwara, H. Idei, O. Kaneko, K. Kawahata, T. Kawamoto, A. Komori, S. Kubo, O. Motojima, V.D. Pustovitov, C. Takahashi, K. Toi and I. Yamada,  
*High-Beta Discharges with Neutral Beam Injection in CHS; Apr. 1994*

NIFS-281 K. Kamada, H. Kinoshita and H. Takahashi,  
*Anomalous Heat Evolution of Deuteron Implanted Al on Electron Bombardment ; May 1994*

NIFS-282 H. Takamaru, T. Sato, K. Watanabe and R. Horiuchi,  
*Super Ion Acoustic Double Layer; May 1994*

NIFS-283 O.Mitarai and S. Sudo  
*Ignition Characteristics in D-T Helical Reactors; June 1994*

NIFS-284 R. Horiuchi and T. Sato,  
*Particle Simulation Study of Driven Magnetic Reconnection in a Collisionless Plasma; June 1994*

NIFS-285 K.Y. Watanabe, N. Nakajima, M. Okamoto, K. Yamazaki, Y. Nakamura, M. Wakatani,  
*Effect of Collisionality and Radial Electric Field on Bootstrap Current in LHD (Large Helical Device); June 1994*

NIFS-286 H. Sanuki, K. Itoh, J. Todoroki, K. Ida, H. Idei, H. Iguchi and H. Yamada,  
*Theoretical and Experimental Studies on Electric Field and Confinement in Helical Systems; June 1994*

NIFS-287 K. Itoh and S.-I. Itoh,  
*Influence of the Wall Material on the H-mode Performance; June 1994*

NIFS-288 K. Itoh, A. Fukuyama, S.-I. Itoh, M. Yagi and M. Azumi  
*Self-Sustained Magnetic Braiding in Toroidal Plasmas: July 1994*

- NIFS-289 Y. Nejoh,  
*Relativistic Effects on Large Amplitude Nonlinear Langmuir Waves in a Two-Fluid Plasma*; July 1994
- NIFS-290 N. Ohyabu, A. Komori, K. Akaishi, N. Inoue, Y. Kubota, A.I. Livshit, N. Noda, A. Sagara, H. Suzuki, T. Watanabe, O. Motojima, M. Fujiwara, A. Iiyoshi,  
*Innovative Divertor Concepts for LHD*; July 1994
- NIFS-291 H. Idei, K. Ida, H. Sanuki, S. Kubo, H. Yamada, H. Iguchi, S. Morita, S. Okamura, R. Akiyama, H. Arimoto, K. Matsuoka, K. Nishimura, K. Ohkubo, C. Takahashi, Y. Takita, K. Toi, K. Tsumori and I. Yamada,  
*Formation of Positive Radial Electric Field by Electron Cyclotron Heating in Compact Helical System*; July 1994
- NIFS-292 N. Noda, A. Sagara, H. Yamada, Y. Kubota, N. Inoue, K. Akaishi, O. Motojima, K. Iwamoto, M. Hashiba, I. Fujita, T. Hino, T. Yamashina, K. Okazaki, J. Rice, M. Yamage, H. Toyoda and H. Sugai,  
*Boronization Study for Application to Large Helical Device*; July 1994
- NIFS-293 Y. Ueda, T. Tanabe, V. Philipps, L. Könen, A. Pospieszczyk, U. Samm, B. Schweer, B. Unterberg, M. Wada, N. Hawkes and N. Noda,  
*Effects of Impurities Released from High Z Test Limiter on Plasma Performance in TEXTOR*; July. 1994
- NIFS-294 K. Akaishi, Y. Kubota, K. Ezaki and O. Motojima,  
*Experimental Study on Scaling Law of Outgassing Rate with A Pumping Parameter*, Aug. 1994
- NIFS-295 S. Bazdenkov, T. Sato, R. Horiuchi, K. Watanabe  
*Magnetic Mirror Effect as a Trigger of Collisionless Magnetic Reconnection*, Aug. 1994
- NIFS-296 K. Itoh, M. Yagi, S.-I. Itoh, A. Fukuyama, H. Sanuki, M. Azumi  
*Anomalous Transport Theory for Toroidal Helical Plasmas*, Aug. 1994 (IAEA-CN-60/D-III-3)
- NIFS-297 J. Yamamoto, O. Motojima, T. Mito, K. Takahata, N. Yanagi, S. Yamada, H. Chikaraishi, S. Imagawa, A. Iwamoto, H. Kaneko, A. Nishimura, S. Satoh, T. Satow, H. Tamura, S. Yamaguchi, K. Yamazaki, M. Fujiwara, A. Iiyoshi and LHD group,  
*New Evaluation Method of Superconductor Characteristics for Realizing the Large Helical Device*; Aug. 1994 (IAEA-CN-60/F-P-3)
- NIFS-298 A. Komori, N. Ohyabu, T. Watanabe, H. Suzuki, A. Sagara, N. Noda, K. Akaishi, N. Inoue, Y. Kubota, O. Motojima, M. Fujiwara and A. Iiyoshi,  
*Local Island Divertor Concept for LHD*; Aug. 1994 (IAEA-CN-60/F-P-4)
- NIFS-299 K. Toi, T. Morisaki, S. Sakakibara, A. Ejiri, H. Yamada, S. Morita,

K. Tanaka, N. Nakajima, S. Okamura, H. Iguchi, K. Ida, K. Tsumori, S. Ohdachi, K. Nishimura, K. Matsuoka, J. Xu, I. Yamada, T. Minami, K. Narihara, R. Akiyama, A. Ando, H. Arimoto, A. Fujisawa, M. Fujiwara, H. Idei, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, T. Ozaki, A. Sagara, C. Takahashi, Y. Takita and T. Watari  
*Impact of Rotational-Transform Profile Control on Plasma Confinement and Stability in CHS*; Aug. 1994 (IAEA-CN-60/A6/C-P-3)

NIFS-300 H. Sugama and W. Horton,  
*Dynamical Model of Pressure-Gradient-Driven Turbulence and Shear Flow Generation in L-H Transition*; Aug. 1994 (IAEA/CN-60/D-P-I-11)

NIFS-301 Y. Hamada, A. Nishizawa, Y. Kawasumi, K.N. Sato, H. Sakakita, R. Liang, K. Kawahata, A. Ejiri, K. Narihara, K. Sato, T. Seki, K. Toi, K. Itoh, H. Iguchi, A. Fujisawa, K. Adachi, S. Hidekuma, S. Hirokura, K. Ida, M. Kojima, J. Koog, R. Kumazawa, H. Kuramoto, T. Minami, I. Negi, S. Ohdachi, M. Sasao, T. Tsuzuki, J. Xu, I. Yamada, T. Watari,  
*Study of Turbulence and Plasma Potential in JIPP T-IIU Tokamak*; Aug. 1994 (IAEA/CN-60/A-2-III-5)

NIFS-302 K. Nishimura, R. Kumazawa, T. Mutoh, T. Watari, T. Seki, A. Ando, S. Masuda, F. Shinpo, S. Murakami, S. Okamura, H. Yamada, K. Matsuoka, S. Morita, T. Ozaki, K. Ida, H. Iguchi, I. Yamada, A. Ejiri, H. Idei, S. Muto, K. Tanaka, J. Xu, R. Akiyama, H. Arimoto, M. Isobe, M. Iwase, O. Kaneko, S. Kubo, T. Kawamoto, A. Lazaros, T. Morisaki, S. Sakakibara, Y. Takita, C. Takahashi and K. Tsumori,  
*ICRF Heating in CHS*; Sep. 1994 (IAEA-CN-60/A-6-I-4)

NIFS-303 S. Okamura, K. Matsuoka, K. Nishimura, K. Tsumori, R. Akiyama, S. Sakakibara, H. Yamada, S. Morita, T. Morisaki, N. Nakajima, K. Tanaka, J. Xu, K. Ida, H. Iguchi, A. Lazaros, T. Ozaki, H. Arimoto, A. Ejiri, M. Fujiwara, H. Idei, A. Iiyoshi, O. Kaneko, K. Kawahata, T. Kawamoto, S. Kubo, T. Kuroda, O. Motojima, V.D. Pustovitov, A. Sagara, C. Takahashi, K. Toi and I. Yamada,  
*High Beta Experiments in CHS*; Sep. 1994 (IAEA-CN-60/A-2-IV-3)

NIFS-304 K. Ida, H. Idei, H. Sanuki, K. Itoh, J. Xu, S. Hidekuma, K. Kondo, A. Sahara, H. Zushi, S.-I. Itoh, A. Fukuyama, K. Adati, R. Akiyama, S. Bessho, A. Ejiri, A. Fujisawa, M. Fujiwara, Y. Hamada, S. Hirokura, H. Iguchi, O. Kaneko, K. Kawahata, Y. Kawasumi, M. Kojima, S. Kubo, H. Kuramoto, A. Lazaros, R. Liang, K. Matsuoka, T. Minami, T. Mizuuchi, T. Morisaki, S. Morita, K. Nagasaki, K. Narihara, K. Nishimura, A. Nishizawa, T. Obiki, H. Okada, S. Okamura, T. Ozaki, S. Sakakibara, H. Sakakita, A. Sagara, F. Sano, M. Sasao, K. Sato, K.N. Sato, T. Saeki, S. Sudo, C. Takahashi, K. Takaka, K. Tsumori, H. Yamada, I. Yamada, Y. Takita, T. Tuzuki, K. Toi and T. Watari,  
*Control of Radial Electric Field in Torus Plasma*; Sep. 1994 (IAEA-CN-60/A-2-IV-2)