

NATIONAL INSTITUTE FOR FUSION SCIENCE

H-Mode Study in CHS

K. Toi, T. Morisaki, S. Sakakibara, S. Ohdachi, T. Minami,
S. Morita, H. Yamada, K. Tanaka, K. Ida, S. Okamura,
A. Ejiri, H. Iguchi, K. Nishimura, K. Matsuoka, A. Ando,
J. Xu, I. Yamada, K. Narihara, R. Akiyama, H. Idei,
S. Kubo, T. Ozaki, C. Takahashi, K. Tsumori

(Received - Jan. 25, 1995)

NIFS-339

Feb.1995

RESEARCH REPORT NIFS Series

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

Presented at Sixth International Toki Conference
on Plasma Physics and Controlled Nuclear Fusion,
Nov.29-Dec.2, 1994, Toki-city, Japan
(to be submitted in Fusion Technology)

H-MODE STUDY IN CHS

K. Toi, T. Morisaki, S. Sakakibara, S. Ohdachi, T. Minami, S. Morita, H. Yamada,
K. Tanaka, K. Ida, S. Okamura, A. Ejiri, H. Iguchi, K. Nishimura, K. Matsuoka,
A. Ando, J. Xu, I. Yamada, K. Narihara, R. Akiyama, H. Idei,
S. Kubo, T. Ozaki, C. Takahashi, K. Tsumori
National Institute for Fusion Science, Nagoya 464-01, Japan

ABSTRACT

In CHS rapid H-mode transition is observed in NBI heated deuterium and hydrogen plasmas without obvious isotope effect, when a net plasma current is ramped up to increase the external rotational transform. The H-mode of CHS has many similarities with those in tokamaks. Recent measurement with fast response Langmuir probes has revealed that the rapid change in floating potential occurs at the transition, but the change follows the formation of edge transport barrier. The presence of $\nu/2\pi = 1$ surface near the edge and sawtooth crash triggered by internal modes may play an important role for determining the H-mode transition in CHS.

Keywords: L-H Transition, Edge Transport Barrier, Heliotron/Torsatron,
Collapse of Magnetic Island, Radial Electric Field

I. INTRODUCTION

Recent confinement study in toroidal magnetic systems such as tokamak or helical system (stellarator or heliotron/torsatron) is addressed to enhanced confinement regimes, because they have a potentiality for minimizing a fusion-reactor size. Discovery of the H-mode in ASDEX accelerated the confinement study in tokamaks[1]. At present, radial electric field shear E_r' or $E \times B$ sheared flow near the edge is thought to be a key factor for the L-H transition and confinement improvement[2]. However, a cause and effect relationship between $E \times B$ sheared flow and the transition has not been established yet. In helical plasmas several experiments for controlling E_r or E_r' externally were carried out in order to improve the plasma confinement, but no H-mode transition was observed so far. A different approach controlling the rotational transform profile was started since 1991 in the CHS heliotron/torsatron, and the H-mode was immediately achieved[3]. The spontaneous H-mode was also observed in W7-AS stellarator[4]. This paper describes the characteristics of the H-mode achieved in CHS along this approach[3,5,6].

II. EXPERIMENTAL SET-UP AND RESULTS

In CHS the H-mode study is carried out on deuterium and hydrogen plasmas heated with co-injected neutral beams (NBI), at the toroidal magnetic field $B_t=1.2$ T and 1.4 T. A net plasma current is induced up to 50 kA mostly by a toroidal inductive voltage. Line averaged electron density just before the transition is adjusted from 1.5 to 3×10^{13} cm^{-3} , and increases up to 3 to 5×10^{13} cm^{-3} in the H-phase. In this experimental series, the external rotational transform is about 0.25 at the magnetic axis and 0.9 at the last closed flux surface(LCFS).

Figure 1 shows a typical H-mode discharge obtained in NBI heated deuterium plasma. In this discharge the plasma current is ramped up to ~ 35 kA in 100 ms by a loop voltage of about 2 V, to increase the external rotational transform. $H\alpha/D\alpha$ emission is depressed within 0.2 ms, which is much shorter than the global energy confinement time(~ 2 -3 ms). The line averaged electron density (n_e) continuously rises during the H-phase till the gas puff is turned off. At the transition edge electron density measured with thermal Li beam probing(LIBP) just inside LCFS is suddenly raised (in 50 to 100 μs) and that just outside LCFS is suddenly reduced, as

shown in Fig.1(b). This clearly indicates a rapid formation of edge transport barrier.

Figure 2 shows time evolution of electron temperature and density profiles obtained with YAG Thomson scattering. At ~ 5 ms after the transition electron temperature near the edge remains unchanged or slightly increases compared with that in L-phase, while electron density is considerably increased there. In the deep H-phase (~ 25 ms after the transition), the electron temperature exhibits a pedestal near the edge and electron density profile evolves to the hollow one. The ion temperature profile also becomes broad. Poloidal rotation of C^{+6} ion is increased in the electron diamagnetic drift direction [6]. The radial electric field is estimated to be ~ 100 V/cm near the edge.

At the L-H transition electron density fluctuations measured with LIBP are clearly suppressed at very near and just outside LCFS. Magnetic fluctuations with various low mode numbers are excited throughout the H-mode discharges. About 10-20 ms before the transition $m/n=1/1$ mode is excited presumably due to appearance of the $1/2\pi=1$ rational surface just inside LCFS. The $m/n=1/1$ mode is thought to be the resistive interchange mode. As shown in Fig.3 this mode amplitude is clearly reduced at the transition, although the $1/2\pi=1$ surface still resides just inside LCFS. Increase in electron temperature at $1/2\pi=1$ rational surface through the increase of $1(a)/2\pi$ in time and a heat pulse due to internal modes may play an important role on suppression of the $m/n=1/1$ mode.

Time evolution of ion saturation current and floating potential across the L-H transition has been measured with Langmuir probes having good time ($2 \mu\text{s}$ sampling) and spatial resolutions (~ 1 mm). Floating potential just outside LCFS, which gradually evolves from positive to negative for 10-15 ms before the transition, is suddenly ($\sim 100-200 \mu\text{s}$) raised at the transition (Fig.4(a)). On the other hand the potential just inside LCFS is suddenly decreased, as shown in Fig.4(b). It is thought that a strong negative E_r is generated at the transition near the edge and presumably E_r' is also enhanced. Note that the change in $H\alpha/D\alpha$ emission is usually slow and often small, because of an integral effect along the line of sight. Most significant point is that in these shots the potential change follows the rapid reduction of ion saturation current which indicates the formation of edge transport barrier. Namely, the strong E_r or E_r' is developed as a result of the L-H transition, not a cause. Above mentioned temporal evolution of floating potential across the transition may be interpreted with the reduction of electron loss enhanced during the L-phase instead of enhanced ion loss. One of possible interpretations of

these observations at the transition is collapse of magnetic islands or healing of braised magnetic surfaces near the edge.

There is an obvious threshold for the plasma current depending on Bt , that is, $I_p \sim 30$ kA at $Bt=1.2$ T and ~ 38 kA at $Bt=1.4$ T. The threshold current also depends on the external $\iota(a)/2\pi$. These results suggest the transition closely correlates with the presence of $\iota/2\pi=1$ surface just inside LCFS. It should be noted that a sawtooth crash initiated by internal MHD mode also plays an important role on the transition. The threshold heating power is within twice of the ITER scaling law[7]. So far we have not recognized obvious isotope effect on the transition character.

III. SUMMARY

The H-mode is achieved in CHS, exhibiting rapid formation of edge transport barrier. Although improvement of global energy confinement is not significant so far, i.e., ~ 20 %, the H-mode in higher beta plasma may provide a route for further improvement of plasma confinement in the heliotron/torsatron through self-stabilization effect. Fast-response Langmuir-probe with good spatial resolution has revealed the generation of enhanced edge radial electric field and E_r' as a result of the formation of edge transport barrier. The cause of generation of enhanced edge E_r or E_r' is under investigation. Collapse of magnetic islands or healing braised magnetic surfaces near the edge may have close connection with the H-mode transition in CHS, through a preferential improvement of edge electron confinement. We are also studying how thus enhanced E_r' contributes to boost a frail transition to more robust one.

ACKNOWLEDGMENTS

Authors thank all members of Torus Experiment Group for many useful discussions and supports. One of them(K.T.) would like to acknowledge Drs. N. Nakajima and H. Sugama, and Prof. M. Wakatani for many stimulating discussions. They also thank Prof. M. Fujiwara and Director General A. Iiyoshi for continuous encouragement.

REFERENCES

1. F. Wagner et al., Phys. Rev. Lett. 49, 1408(1982).
2. K.H. Burrell, Plasma Phys. Control. Fusion, 36, A291 (1994).
3. K. Toi et al., in Plasma Phys. Control. Nucl. Fusion Res.(Proc. 14th IAEA conf., Würzburg, 1992) Vol.2, IAEA, Vienna(1993) 461.
4. Erckmann V et al 1993a Plasma Physics and Controlled Nuclear Fusion Research (Proc. 14th Int. Conf., Würzburg, 1992) vol 2 (Vienna: IAEA)p469.
5. K. Toi et al., Plasma Phys. Control. Fusion, 36, A117 (1994).
6. K. Toi et al., to be published in Plasma Phys. Control. Nucl. Fusion Res.(Proc. 15th IAEA conf., Seville, 1994) paper NO. IAEA-CN-60/A6/C-P-3.
7. H-mode database working group, in Plasma Phys. Control. Nucl. Fusion Res.(Proc. 14th IAEA conf, Würzburg, 1992) Vol.3, IAEA, Vienna (1993) 251.

Figure Captions

- Fig.1(a) Time evolution of $H\alpha/D\alpha$ emission, line averaged electron density at the central and edge chords ($\langle r \rangle / \langle a \rangle \sim 0$ and 0.77), and plasma current in a typical deuterium H-mode discharge, where $B_t = 1.2$ T. Two solid vertical lines denote the first and second L-H transition, and dotted ones the H-L transition.
- (b) Time behaviors of LIBP signals which is roughly proportional to local electron density near the edge, where Li8 is the signal from just outside LCFS and Li6 just inside LCFS.
- Fig.2 Time evolution of electron temperature and density profile measured with YAG Thomson scattering in the cross-section horizontally elongated, where solid circles, open circles and diamonds denote the profiles at $t = 80$ ms, 100 ms and 120 ms respectively. The L-H transition occurs at $t \sim 95$ ms.
- Fig.3 Time evolution of magnetic fluctuation amplitude in the range of 5 to 11 kHz. The fluctuations are dominated by $m/n = 1/1$ mode activity. The transition is seen from LIBP signal from just outside LCFS.
- Fig.4(a) Time evolution of ion saturation current (I_{is}) and floating potential (ϕ_{fl}) measured just outside LCFS, together with $H\alpha/D\alpha$ emission, where these data are averaged over $20 \mu s$ and the L-H transition occurs at 97.05 ms.
- (b) Time evolution of I_{is} and ϕ_{fl} just inside LCFS, when the L-H transition occurs at 97.38 ms.




Fig.1 K. Toi et al.




Fig.2 K. Toi et al.




Fig.3 K. Toi et al.




Fig.4 K. Toi et al.

Recent Issues of NIFS Series

- NIFS-297 J. Yamamoto, O. Motojima, T. Mito, K. Takahata, N. Yanagi, S. Yamada, H. Chikaraishi, S. Imagawa, A. Iwamoto, H. Kaneko, A. Nishimura, S. Satoh, T. Satow, H. Tamura, S. Yamaguchi, K. Yamazaki, M. Fujiwara, A. Iiyoshi and LHD group,
New Evaluation Method of Superconductor Characteristics for Realizing the Large Helical Device; Aug. 1994 (IAEA-CN-60/F-P-3)
- NIFS-298 A. Komori, N. Ohyabu, T. Watanabe, H. Suzuki, A. Sagara, N. Noda, K. Akaishi, N. Inoue, Y. Kubota, O. Motojima, M. Fujiwara and A. Iiyoshi,
Local Island Divertor Concept for LHD; Aug. 1994 (IAEA-CN-60/F-P-4)
- NIFS-299 K. Toi, T. Morisaki, S. Sakakibara, A. Ejiri, H. Yamada, S. Morita, K. Tanaka, N. Nakajima, S. Okamura, H. Iguchi, K. Ida, K. Tsumori, S. Ohdachi, K. Nishimura, K. Matsuoka, J. Xu, I. Yamada, T. Minami, K. Narihara, R. Akiyama, A. Ando, H. Arimoto, A. Fujisawa, M. Fujiwara, H. Idei, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, T. Ozaki, A. Sagara, C. Takahashi, Y. Takita and T. Watari,
Impact of Rotational-Transform Profile Control on Plasma Confinement and Stability in CHS; Aug. 1994 (IAEA-CN-60/A6/C-P-3)
- NIFS-300 H. Sugama and W. Horton,
Dynamical Model of Pressure-Gradient-Driven Turbulence and Shear Flow Generation in L-H Transition; Aug. 1994 (IAEA/CN-60/D-P-I-11)
- NIFS-301 Y. Hamada, A. Nishizawa, Y. Kawasumi, K.N. Sato, H. Sakakita, R. Liang, K. Kawahata, A. Ejiri, K. Narihara, K. Sato, T. Seki, K. Toi, K. Itoh, H. Iguchi, A. Fujisawa, K. Adachi, S. Hidekuma, S. Hirokura, K. Ida, M. Kojima, J. Koog, R. Kumazawa, H. Kuramoto, T. Minami, I. Negi, S. Ohdachi, M. Sasao, T. Tsuzuki, J. Xu, I. Yamada, T. Watari,
Study of Turbulence and Plasma Potential in JIPP T-IIU Tokamak; Aug. 1994 (IAEA/CN-60/A-2-III-5)
- NIFS-302 K. Nishimura, R. Kumazawa, T. Mutoh, T. Watari, T. Seki, A. Ando, S. Masuda, F. Shinpo, S. Murakami, S. Okamura, H. Yamada, K. Matsuoka, S. Morita, T. Ozaki, K. Ida, H. Iguchi, I. Yamada, A. Ejiri, H. Idei, S. Muto, K. Tanaka, J. Xu, R. Akiyama, H. Arimoto, M. Isobe, M. Iwase, O. Kaneko, S. Kubo, T. Kawamoto, A. Lazaros, T. Morisaki, S. Sakakibara, Y. Takita, C. Takahashi and K. Tsumori,
ICRF Heating in CHS; Sep. 1994 (IAEA-CN-60/A-6-I-4)
- NIFS-303 S. Okamura, K. Matsuoka, K. Nishimura, K. Tsumori, R. Akiyama, S. Sakakibara, H. Yamada, S. Morita, T. Morisaki, N. Nakajima, K. Tanaka, J. Xu, K. Ida, H. Iguchi, A. Lazaros, T. Ozaki, H. Arimoto, A. Ejiri, M. Fujiwara, H. Idei, A. Iiyoshi, O. Kaneko, K. Kawahata, T. Kawamoto, S. Kubo, T. Kuroda, O. Motojima, V.D. Pustovitov, A. Sagara, C. Takahashi, K. Toi and I. Yamada,

High Beta Experiments in CHS; Sep. 1994 (IAEA-CN-60/A-2-IV-3)

- NIFS-304 K. Ida, H. Idei, H. Sanuki, K. Itoh, J. Xu, S. Hidekuma, K. Kondo, A. Sahara, H. Zushi, S.-I. Itoh, A. Fukuyama, K. Adati, R. Akiyama, S. Bessho, A. Ejiri, A. Fujisawa, M. Fujiwara, Y. Hamada, S. Hirokura, H. Iguchi, O. Kaneko, K. Kawahata, Y. Kawasumi, M. Kojima, S. Kubo, H. Kuramoto, A. Lazaros, R. Liang, K. Matsuoka, T. Minami, T. Mizuuchi, T. Morisaki, S. Morita, K. Nagasaki, K. Narihara, K. Nishimura, A. Nishizawa, T. Obiki, H. Okada, S. Okamura, T. Ozaki, S. Sakakibara, H. Sakakita, A. Sagara, F. Sano, M. Sasao, K. Sato, K.N. Sato, T. Saeki, S. Sudo, C. Takahashi, K. Tanaka, K. Tsumori, H. Yamada, I. Yamada, Y. Takita, T. Tuzuki, K. Toi and T. Watari, *Control of Radial Electric Field in Torus Plasma; Sep. 1994 (IAEA-CN-60/A-2-IV-2)*
- NIFS-305 T. Hayashi, T. Sato, N. Nakajima, K. Ichiguchi, P. Merkel, J. Nührenberg, U. Schwenn, H. Gardner, A. Bhattacharjee and C.C.Hegna, *Behavior of Magnetic Islands in 3D MHD Equilibria of Helical Devices; Sep. 1994 (IAEA-CN-60/D-2-II-4)*
- NIFS-306 S. Murakami, M. Okamoto, N. Nakajima, K.Y. Watanabe, T. Watari, T. Mutoh, R. Kumazawa and T. Seki, *Monte Carlo Simulation for ICRF Heating in Heliotron/Torsatrons; Sep. 1994 (IAEA-CN-60/D-P-I-14)*
- NIFS-307 Y. Takeiri, A. Ando, O. Kaneko, Y. Oka, K. Tsumori, R. Akiyama, E. Asano, T. Kawamoto, T. Kuroda, M. Tanaka and H. Kawakami, *Development of an Intense Negative Hydrogen Ion Source with a Wide-Range of External Magnetic Filter Field; Sep. 1994*
- NIFS-308 T. Hayashi, T. Sato, H.J. Gardner and J.D. Meiss, *Evolution of Magnetic Islands in a Heliac; Sep. 1994*
- NIFS-309 H. Amo, T. Sato and A. Kageyama, *Intermittent Energy Bursts and Recurrent Topological Change of a Twisting Magnetic Flux Tube; Sep.1994*
- NIFS-310 T. Yamagishi and H. Sanuki, *Effect of Anomalous Plasma Transport on Radial Electric Field in Torsatron/Heliotron; Sep. 1994*
- NIFS-311 K. Watanabe, T. Sato and Y. Nakayama, *Current-profile Flattening and Hot Core Shift due to the Nonlinear Development of Resistive Kink Mode; Oct. 1994*
- NIFS-312 M. Salimullah, B. Dasgupta, K. Watanabe and T. Sato, *Modification and Damping of Alfvén Waves in a Magnetized Dusty Plasma; Oct. 1994*

- NIFS-313 K. Ida, Y. Miura, S.-I. Itoh, J.V. Hofmann, A. Fukuyama, S. Hidekuma, H. Sanuki, H. Idei, H. Yamada, H. Iguchi, K. Itoh,
Physical Mechanism Determining the Radial Electric Field and its Radial Structure in a Toroidal Plasma; Oct. 1994
- NIFS-314 Shao-ping Zhu, R. Horiuchi, T. Sato and The Complexity Simulation Group,
Non-Taylor Magnetohydrodynamic Self-Organization; Oct. 1994
- NIFS-315 M. Tanaka,
Collisionless Magnetic Reconnection Associated with Coalescence of Flux Bundles; Nov. 1994
- NIFS-316 M. Tanaka,
Macro-EM Particle Simulation Method and A Study of Collisionless Magnetic Reconnection; Nov. 1994
- NIFS-317 A. Fujisawa, H. Iguchi, M. Sasao and Y. Hamada,
Second Order Focusing Property of 210° Cylindrical Energy Analyzer; Nov. 1994
- NIFS-318 T. Sato and Complexity Simulation Group,
Complexity in Plasma - A Grand View of Self-Organization; Nov. 1994
- NIFS-319 Y. Todo, T. Sato, K. Watanabe, T.H. Watanabe and R. Horiuchi,
MHD-Vlasov Simulation of the Toroidal Alfvén Eigenmode; Nov. 1994
- NIFS-320 A. Kageyama, T. Sato and The Complexity Simulation Group,
Computer Simulation of a Magnetohydrodynamic Dynamo II; Nov. 1994
- NIFS-321 A. Bhattacharjee, T. Hayashi, C.C.Hegna, N. Nakajima and T. Sato,
Theory of Pressure-induced Islands and Self-healing in Three-dimensional Toroidal Magnetohydrodynamic Equilibria; Nov. 1994
- NIFS-322 A. Iiyoshi, K. Yamazaki and the LHD Group,
Recent Studies of the Large Helical Device; Nov. 1994
- NIFS-323 A. Iiyoshi and K. Yamazaki,
The Next Large Helical Devices; Nov. 1994
- NIFS-324 V.D. Pustovitov
Quasisymmetry Equations for Conventional Stellarators; Nov. 1994
- NIFS-325 A. Taniike, M. Sasao, Y. Hamada, J. Fujita, M. Wada,
The Energy Broadening Resulting from Electron Stripping Process of a Low Energy Au⁻ Beam; Dec. 1994
- NIFS-326 I. Viniar and S. Sudo,
New Pellet Production and Acceleration Technologies for High Speed Pellet Injection System "HIPEL" in Large Helical Device; Dec. 1994

- NIFS-327 Y. Hamada, A. Nishizawa, Y. Kawasumi, K. Kawahata, K. Itoh, A. Ejiri, K. Toi, K. Narihara, K. Sato, T. Seki, H. Iguchi, A. Fujisawa, K. Adachi, S. Hidekuma, S. Hirokura, K. Ida, M. Kojima, J. Koong, R. Kumazawa, H. Kuramoto, R. Liang, T. Minami, H. Sakakita, M. Sasao, K.N. Sato, T. Tsuzuki, J. Xu, I. Yamada, T. Watari,
Fast Potential Change in Sawteeth in JIPP T-IIU Tokamak Plasmas; Dec. 1994
- NIFS-328 V.D. Pustovitov,
Effect of Satellite Helical Harmonics on the Stellarator Configuration; Dec. 1994
- NIFS-329 K. Itoh, S-I. Itoh and A. Fukuyama,
A Model of Sawtooth Based on the Transport Catastrophe; Dec. 1994
- NIFS-330 K. Nagasaki, A. Ejiri,
Launching Conditions for Electron Cyclotron Heating in a Sheared Magnetic Field; Jan. 1995
- NIFS-331 T.H. Watanabe, Y. Todo, R. Horiuchi, K. Watanabe, T. Sato,
An Advanced Electrostatic Particle Simulation Algorithm for Implicit Time Integration; Jan. 1995
- NIFS-332 N. Bekki and T. Karakisawa,
Bifurcations from Periodic Solution in a Simplified Model of Two-dimensional Magnetoconvection; Jan. 1995
- NIFS-333 K. Itoh, S.-I. Itoh, M. Yagi, A. Fukuyama,
Theory of Anomalous Transport in Reverse Field Pinch; Jan. 1995
- NIFS-334 K. Nagasaki, A. Isayama and A. Ejiri
Application of Grating Polarizer to 106.4GHz ECH System on Heliotron-E; Jan. 1995
- NIFS-335 H. Takamaru, T. Sato, R. Horiuchi, K. Watanabe and Complexity Simulation Group,
A Self-Consistent Open Boundary Model for Particle Simulation in Plasmas; Feb. 1995
- NIFS-336 B.B. Kadomtsev,
Quantum Telegraph : is it possible?; Feb. 1995
- NIFS-337 B.B.Kadomtsev,
Ball Lightning as Self-Organization Phenomenon; Feb. 1995
- NIFS-338 Y. Takeiri, A. Ando, O. Kaneko, Y. Oka, K. Tsumori, R. Akiyama, E. Asano, T. Kawamoto, M. Tanaka and T. Kuroda,
High-Energy Acceleration of an Intense Negative Ion Beam; Feb. 1995