



INTERNATIONAL ATOMIC ENERGY AGENCY

17th IAEA Fusion Energy Conference
Yokohama, Japan, 19 - 24 October 1998

IAEA-CN-69/ THP2/22

NATIONAL INSTITUTE FOR FUSION SCIENCE

Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes

Y. Todo and T. Sato

(Received - Sep. 25, 1998)

NIFS-563

Oct. 1998

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

RESEARCH REPORT NIFS Series

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the professional nature of its content, and for changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the author(s). The views do not necessarily reflect those of the government of the designated Member country. In particular, the organizations sponsoring this meeting cannot be held responsible for any inaccuracies appearing in this preprint.

NAGOYA, JAPAN

Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes

Y. TODO, T. SATO

Theory and Computer Simulation Center,
National Institute for Fusion Science,
322-6 Oroshi-cho, Toki-shi 509-5292,
Japan

Abstract

Particle-magnetohydrodynamic and Fokker-Planck-magnetohydrodynamic simulations of fast ions and toroidicity-induced Alfvén eigenmodes (TAE modes) have been carried out. Alpha particle losses induced by TAE mode are investigated with particle-magnetohydrodynamic simulations. Trapped particles near the passing-trapped boundary in the phase space are also lost appreciably in addition to the counter-passing particles. In Fokker-Planck-magnetohydrodynamic simulation source and slowing-down of fast ions are considered. A coherent pulsating behavior of multiple TAE modes, which occurs in neutral beam injection experiments, is observed when the slowing-down time is much longer than the damping time of the TAE modes and the fast-ion pressure is sufficiently high. For a slowing-down time comparable to the damping time, the TAE modes reach steady saturation levels.

1. INTRODUCTION

The toroidicity-induced Alfvén eigenmode (TAE mode) is a shear-Alfvén eigenmode in toroidal plasmas [1]. TAE modes have been observed in many experiments such as the neutral beam injection (NBI) experiments [2, 3], the ion-cyclotron-range-of-frequency (ICRF) heating experiments [4, 5], and D-T fusion experiments at TFTR [6]. Nonlinear evolution of TAE modes is an important issue for fusion reactors, since TAE modes of sufficiently large amplitude can induce fast-ion losses [3, 7]. The first purpose of the present paper is to investigate the TAE-induced fast ion losses with particle-magnetohydrodynamic (MHD) simulation [8]. The second purpose is to examine the pulsating behavior of multiple TAE modes in NBI experiments, which is completely different from the steady behavior of ICRF-driven and alpha-driven TAE modes. It is investigated with Fokker-Planck-MHD simulation [9].

2. PARTICLE-MHD SIMULATION

In the particle-MHD simulation [8], the particle simulation method is used for the fast ions. In the model employed here, plasma is divided into two parts, the background plasma and fast ions. The background plasma is described by the magnetohydrodynamic (MHD) equations and the electromagnetic field is given by the MHD description. This approximation is reasonable under the condition that the fast ion density is much less than the background plasma density.

The initial condition considered here is $B=5\text{T}$, $I_p=2\text{MA}$, $R_0=2.7\text{m}$, $a=0.9\text{m}$, $n=10^{20}\text{m}^{-3}$, and $\langle\beta\rangle=0.88\%$. Fast ion distribution is a slowing-down distribution with the maximum energy of 3.5 MeV, which is isotropic in the velocity space. Nonlinear simulations for $n=2$ TAE modes have been carried out for volume-average fast-ion pressures of 0.33% and 0.66%. The number of used particles is 4×10^6 . Fast ion losses induced by the TAE modes are observed. For the higher-pressure case, 1% of fast ions are lost in 10^3 Alfvén time. Lost particles can be classified into three types; 1) passing particles with negative parallel velocities (counter-passing particles) which cross the passing-trapped boundary just before an encounter with the wall, 2) trapped particles, 3) passing particles other than the first type. Percentages to the total number of lost particles are 63%, 26%, and 11%, respectively. It is clear that counter-passing particles are the major part of lost particles and the main loss mechanism is crossing the passing-trapped boundary in the phase space. It is consistent with the results of test particle simulation in Ref. [10]. It must be noted, however, that the trapped particles near this boundary are also lost appreciably in addition to the counter-passing particles.

3. FOKKER-PLANCK-MHD SIMULATION

3.1. Model

Past simulation studies of TAE mode [11-14] started each simulation runs from arbitrarily chosen fast-ion distributions, and could not explain the recursive bursts at NBI heating experiments and steady saturation levels at ICRF heating experiments. Berk and Breizman [15] pointed out the significance of distribution forming processes such as particle source, slowing down, pitch-angle scattering, and wave heating. What seems to be lacking in the past simulation studies is to incorporate the distribution forming processes of fast ions.

As a first step towards a comprehensive simulation of TAE modes, we incorporate particle source and slowing down for fast ions in the Vlasov-MHD simulation code [12]. The time evolution of the fast-ion distribution function in a four-dimensional phase space (three-dimensional configuration space and one-dimensional velocity space for the parallel velocity) is followed by a finite difference method of fourth-order accuracy in space and time. The pitch-angle scattering and wave heating are not considered in the present study. We consider a four-dimensional phase space (R, ϕ, z, v) , where v is the parallel velocity and (R, ϕ, z) are the cylindrical coordinates. Although only the parallel velocity component is taken into account for simplicity, the Jacobian for the three-dimensional velocity space is employed to be consistent with the slowing-down term. The critical velocity in the slowing down term and the birth velocity of fast ions are chosen to be equal to 0.3 and 1.5 of the Alfvén velocity, respectively. The spatial profile of the particle source is a Gaussian with the scale-length of $0.4a$, where a is the minor radius. We introduce two parameters, P_0 , which is the fast-ion pressure at the magnetic axis when any MHD disturbance is absent, and the slowing down time τ_s . The initial condition is a tokamak MHD equilibrium where the aspect ratio is 3. The minor radius is 16 times larger than the parallel Larmor radius of a fast ion with the Alfvén velocity. In this paper a finite viscosity of $2 \times 10^{-3} R_0 v_A$ is considered to damp TAE modes. It yields, for example, an e -folding damping time of $130\tau_A$ for an $n=2$ TAE mode which is the most unstable TAE mode in the results described below.

3.2. Results

3.2.1. Case A: $P_0 = 2\%$ (normalized by the magnetic pressure) and $\tau_s = 100\tau_A$

In this case the slowing-down time is comparable to the damping time. Destabilized are TAE modes with the toroidal mode numbers from 1 to 3, and another Alfvén eigenmode with the toroidal mode number of 4, which has a doubled frequency to that of a TAE mode. Time history of $m/n = 2/1, 3/2, 4/3, 6/4$ harmonics is shown in Fig. 1(a). The $n=2-4$ eigenmodes are localized in the core region. It can be seen that $n=2-4$ modes reach roughly steady saturation levels after $500\tau_A$.

3.2.2. Case B: $P_0 = 4\%$ (normalized by the magnetic pressure) and $\tau_s = 1000\tau_A$

In this case the slowing-down time is much longer than the damping time and the fast-ion pressure is relatively high. TAE modes with toroidal mode numbers from 1 to 4 are destabilized. The spatial profiles of $n=3$ and 4 eigenmodes peak at outer locations and are more broadened in the radial direction than those in case A. Time history of $m/n = 2/1, 3/2, 4/3, 6/4$ harmonics is shown in Fig. 1(b). It can be seen that they have a pulsating behavior. The spatial profile of the fast-ion source destabilizes the $n=2$ TAE mode first. Later than this precursory growth of the $n=2$ TAE mode, the other TAE modes grow to levels comparable to that of the $n=2$ TAE mode. At the second and third bursts, all of the TAE modes behave coherently. This pulsating bursts can be naturally interpreted as a cycle of reformation of fast-ion distribution, growth of TAE modes due to increasing fast-ion drive, global flattening of fast-ion distribution, and damping of TAE modes due to reduced fast-ion drive.




FIG. 1. Time evolution of $n=1-4$ TAE mode amplitudes for (a) $P_0=2\%$ and $\tau_s = 100\tau_A$, and (b) $P_0=4\%$ and $\tau_s = 1000\tau_A$.

3.2.3 Fast ion distribution at nonlinear phase

The fast-ion distribution is strongly affected by TAE modes. We show the fast-ion distribution averaged in the toroidal angle with $v = v_A$ in Fig. 2. For case A, the distribution is locally flattened in the core region where the TAE activity is strong. On the other hand, for case B, the $n=2$ and 3 TAE modes, which spatially peak at outer locations as mentioned above, are destabilized, and the fast-ion distribution is globally flattened due to the overlapped many TAE modes. It is interesting to note that the distribution is spread close to the wall boundary at the mid-plane. It explains the fast-ion losses at the NBI experiments [3], which take place mainly at the mid-plane.

4. SUMMARY

In this paper we described the results of particle-MHD and Fokker-Planck-MHD simulations. With particle-MHD simulations, it is confirmed that the major part of lost alpha particles induced by TAE mode at the present large tokamaks is the counter-passing particles close to the passing-trapped boundary, and it is found that trapped particles are also lost appreciably. Furthermore, we have investigated the time evolution of fast ions and the toroidal Alfvén eigenmodes with the Fokker-Planck-MHD simulations. We have demonstrated that the consideration of the distribution forming processes of fast ions is crucial to understand the time evolution of TAE modes and fast ions.

Simulation results are summarized as follows:

- [1] When the time scale of the distribution forming process [the slowing-down time in the present study] is comparable to the damping time, TAE modes persist after saturation with steady amplitudes.
- [2] When the time scale of the distribution forming process is longer than the damping time and the fast-ion pressure is sufficiently high, many TAE modes have a coherent pulsating behavior.




FIG. 2. Fast ion distribution as a function of R and z averaged in the toroidal angle with $v = v_A$ for (a) $P_o=2\%$, $\tau_s = 100\tau_A$, $t=900\tau_A$, and (b) $P_o=4\%$, $\tau_s = 1000\tau_A$, $t=2800\tau_A$.

The pulsating behavior is similar to that observed in NBI heating experiments. In this case, the fast-ion distribution function is globally flattened due to the overlapped many TAE modes. This global flattening explains the fast-ion losses observed at the NBI heating experiments. The steady saturation found for the short slowing-down time is also interesting when we compare it with the ICRF-driven and alpha-driven TAE modes, though a more realistic simulation is required for a detailed discussion.

ACKNOWLEDGMENTS

The authors would like to express their thanks to all members of the Complexity Simulation Group for useful discussions. Numerical computations were performed at the Advanced Computing System for Complexity Simulation of National Institute for Fusion Science.

REFERENCES

- [1] CHENG, C. Z., CHANCE, M. S., Phys.Fluids 29 (1986) 3659.
- [2] WONG, K. L., et al., Phys. Rev. Lett. 66 (1991) 1874.
- [3] DUONG, H. H., et al., Nucl. Fusion 33 (1993) 749.
- [4] WONG, K. L., et al., Plasma Phys. Control. Fusion 36 (1994) 879.
- [5] KIMURA, H., et al., J. Plasma Fusion Res. 71 (1995) 1147.
- [6] NAZIKIAN, R., et al., Phys. Rev. Lett. 78 (1997) 2976.
- [7] DARROW, D. S., et al., Nucl. Fusion 37 (1997) 939.
- [8] TODO, Y., SATO, T., Phys. Plasmas 5 (1998) 1321.
- [9] TODO, Y., SATO, T., "TAE Instability and Fast Ion Losses", in the Proceedings of the Joint Varenna-Lausanne International Workshop on Theory of Fusion Plasmas (Varenna, 1998), in press.
- [10] SIGMAR, D. J., et al., Phys. Fluids B 4 (1992) 1506.
- [11] FU, G. Y., PARK, W., Phys. Rev. Lett. 74 (1995) 1594.
- [12] TODO, Y., et al., Phys. Plasmas 2 (1995) 2711.
- [13] WU, Y., et al., Phys. Plasmas 2 (1995) 4555.
- [14] CANDY, J., et al, Phys. Plasmas 4 (1997) 2597.
- [15] BERK, H. L., BREIZMAN, B. N., Phys. Fluids B 2 (1990) 2226, 2235, 2246.

Recent Issues of NIFS Series

- NIFS-529 J. Uramoto,
Pair Creation of Negative and Positive Pionlike (Muonlike) Particle by Interaction between an Electron Bunch and a Positive Ion Bunch; Dec 1997
- NIFS-530 J. Uramoto,
Measuring Method of Decay Time of Negative Muonlike Particle by Beam Collector Applied RF Bias Voltage, Dec 1997
- NIFS-531 J. Uramoto,
Confirmation Method for Metal Plate Penetration of Low Energy Negative Pionlike or Muonlike Particle Beam under Positive Ions, Dec. 1997
- NIFS-532 J. Uramoto,
Pair Creations of Negative and Positive Pionlike (Muonlike) Particle or K Mesonlike (Muonlike) Particle in H₂ or D₂ Gas Discharge in Magnetic Field; Dec 1997
- NIFS-533 S. Kawata, C. Boonmee, T. Teramoto, L. Drska, J. Limpouch, R. Liska, M. Sinor,
Computer-Assisted Particle-in-Cell Code Development, Dec 1997
- NIFS-534 Y. Matsukawa, T. Suda, S. Ohnuki and C. Namba,
Microstructure and Mechanical Property of Neutron Irradiated TiNi Shape Memory Alloy; Jan. 1998
- NIFS-535 A. Fujisawa, H. Iguchi, H. Idei, S. Kubo, K. Matsuoka, S. Okamura, K. Tanaka, T. Minami, S. Ohdachi, S. Monta, H. Zushi, S. Lee, M. Osakabe, R. Akiyama, Y. Yoshimura, K. Toi, H. Sanuki, K. Itoh, A. Shimizu, S. Takagi, A. Ejiri, C. Takahashi, M. Kojima, S. Hidekuma, K. Ida, S. Nishimura, N. Inoue, R. Sakamoto, S.-I. Itoh, Y. Hamada, M. Fujiwara,
Discovery of Electric Pulsation in a Toroidal Helical Plasma; Jan 1998
- NIFS-536 Lj.R. Hadzievski, M.M. Skoric, M. Kono and T. Sato,
Simulation of Weak and Strong Langmuir Collapse Regimes, Jan. 1998
- NIFS-537 H. Sugama, W. Horton,
Nonlinear Electromagnetic Gyrokinetic Equation for Plasmas with Large Mean Flows; Feb 1998
- NIFS-538 H. Iguchi, T.P. Crowley, A. Fujisawa, S. Lee, K. Tanaka, T. Minami, S. Nishimura, K. Ida, R. Akiyama, Y. Hamada, H., Idei M. Isobe, M. Kojima, S. Kubo, S. Morita, S. Ohdachi, S. Okamura, M. Osakabe, K. Matsuoka, C. Takahashi and K. Toi,
Space Potential Fluctuations during MHD Activities in the Compact Helical System (CHS); Feb. 1998
- NIFS-539 Takashi Yabe and Yan Zhang,
Effect of Ambient Gas on Three-Dimensional Breakup in Coronet Formation Process; Feb. 1998
- NIFS-540 H. Nakamura, K. Ikeda and S. Yamaguchi,
Transport Coefficients of InSb in a Strong Magnetic Field; Feb. 1998
- NIFS-541 J. Uramoto,
Development of ν_{μ} Beam Detector and Large Area ν_{μ} Beam Source by H₂ Gas Discharge (I); Mar. 1998
- NIFS-542 J. Uramoto,
Development of $\bar{\nu}_{\mu}$ Beam Detector and Large Area $\bar{\nu}_{\mu}$ Beam Source by H₂ Gas Discharge (II); Mar 1998
- NIFS-543 J. Uramoto,
Some Problems inside a Mass Analyzer for Pions Extracted from a H₂ Gas Discharge; Mar 1998
- NIFS-544 J. Uramoto,
Simplified ν_{μ} $\bar{\nu}_{\mu}$ Beam Detector and ν_{μ} $\bar{\nu}_{\mu}$ Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch; Mar. 1998
- NIFS-545 J. Uramoto,
Various Neutrino Beams Generated by D₂ Gas Discharge; Mar 1998

- NIFS-546 R. Kanno, N. Nakajima, T. Hayashi and M. Okamoto,
Computational Study of Three Dimensional Equilibria with the Bootstrap Current; Mar. 1998
- NIFS-547 R. Kanno, N. Nakajima and M. Okamoto,
Electron Heat Transport in a Self-Similar Structure of Magnetic Islands; Apr. 1998
- NIFS-548 J.E. Rice,
Simulated Impurity Transport in LHD from MIST; May 1998
- NIFS-549 M.M. Skonc, T. Sato, A.M. Maluckov and M.S. Jovanovic,
On Kinetic Complexity in a Three-Wave Interaction; June 1998
- NIFS-550 S. Goto and S. Kida,
Passive Saclar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation; June 1998
- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,
Initial Value Problem of the Toroidal Ion Temperature Gradient Mode ; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watan,
Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device ; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,
oronaization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-; July 1998
- NIFS-554 S. Murakami and H. Saleem,
Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations; July 1998
- NIFS-555 H. Nakamura, K. Ikeda and S. Yamaguchi,
Physical Model of Nernst Element; Aug. 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,
Numerical Computation of Thermoelectric and Thermomagnetic Effects; Aug. 1998
- NIFS-557 T. Yasuhiko, O. Masaki, T. Katsuyoshi, O. Yoshihide, K. Osamu, A. Eiji, K. Toshikazu, A. Ryuichi and T. Masanobu,
Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System; Aug. 1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Structure Organization of Polyampholytes; Sep. 1998
- NIFS-559 R. Horiuchi, K. Nishimura and T. Watanabe,
Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations; Sep. 1998
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Kholopenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,
Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection; Oct. 1998
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,
Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron; Oct. 1998
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,
Next Generation Toroidal Devices; Oct. 1998
- NIFS-563 Y. Todo and T. Sato,
Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes; Oct. 1998
(IAEA-CN-69/THP2/22)