



INTERNATIONAL ATOMIC ENERGY AGENCY

17th IAEA Fusion Energy Conference  
Yokohama, Japan, 19 - 24 October 1998

IAEA-CN-69/EX2/3

## NATIONAL INSTITUTE FOR FUSION SCIENCE

### Plasma Confinement Studies in LHD

M. Fujiwara, H. Yamada, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B. J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimoizuma, M. Shoji, H. Sazuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, M. Iima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iiyoshi

(Received - Oct. 7, 1998 )

NIFS-572

Oct. 1998

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

## RESEARCH REPORT NIFS Series

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The view expressed and the statements made remain the responsibility of the named author(s). The views do not necessarily reflect those of the government of the designated Member State(s). In particular, the organizations sponsoring this meeting cannot be held responsible for any material reproduced in this preprint.

NAGOYA, JAPAN

## PLASMA CONFINEMENT STUDIES IN LHD

M.FUJIWARA, H.YAMADA, A.EJIRI, M.EMOTO, H.FUNABA, M.GOTO,  
 K.IDA, H.IDEI, S.INAGAKI, S.KADO, O.KANEKO, K.KAWAHATA,  
 A.KOMORI, S.KUBO, R.KUMAZAWA, S.MASUZAKI, T.MINAMI,  
 J.MIYAZAWA, T.MORISAKI, S.MORITA, S.MURAKAMI, S.MUTO,  
 T.MUTOH, Y.NAGAYAMA, Y.NAKAMURA, H.NAKANISHI, K.NARIHARA,  
 K.NISHIMURA, N.NODA, T.KOBUCHI, S.OHDACHI, N.OHYABU, Y.OKA,  
 M.OSAKABE, T.OZAKI, B.J.PETERSON, A.SAGARA, S.SAKAKIBARA,  
 R.SAKAMOTO, H.SASAO, M.SASAO, K.SATO, M.SATO, T.SEKI,  
 T.SHIMOZUMA, M.SHOJI, H.SUZUKI, Y.TAKEIRI, K.TANAKA, K.TOI,  
 T.TOKUZAWA, K.TSUMORI, K.TSUZUKI, I.YAMADA, S.YAMAGUCHI,  
 M.YOKOYAMA, K.Y.WATANABE, T.WATARI, R.AKIYAMA, H.CHIKARAISHI,  
 K.HABA, S.HAMAGUCHI, M.IIMA, S.IMAGAWA, N.INOUE, K.IWAMOTO,  
 S.KITAGAWA, Y.KUBOTA, J.KODAIRA, R.MAEKAWA, T.MITO,  
 T.NAGASAKA, A.NISHIMURA, Y.TAKITA, C.TAKAHASHI, K.TAKAHATA,  
 K.YAMAUCHI, H.TAMURA, T.TSUZUKI, S.YAMADA, N.YANAGI, H.YONEZU,  
 Y.HAMADA, K.MATSUOKA, K.MURAI, K.OHKUBO, I.OHTAKE,  
 M.OKAMOTO, S.SATOH, T.SATOW, S.SUDO, S.TANAHASHI, K.YAMAZAKI,  
 O.MOTOJIMA, A.IIYOSHI

National Institute for Fusion Science,  
 Toki, Gifu,  
 Japan

### Abstract

The initial experiments of the Large Helical Device (LHD) have extended confinement studies on currentless plasmas to a large scale ( $R=3.9$  m,  $a = 0.6$  m). Heating by NBI of 3 MW has produced plasmas with a fusion triple product of  $8 \times 10^{18}$  keV m<sup>-3</sup>s at a magnetic field of 1.5T. An electron temperature of 1.5 keV and an ion temperature of 1.1 keV have been achieved simultaneously at the line-averaged electron density of  $1.5 \times 10^{19}$  m<sup>-3</sup>. The maximum stored energy has reached 0.22 MJ with neither unexpected confinement deterioration nor visible MHD instabilities, which corresponds to  $\langle \beta \rangle = 0.7\%$ . Energy confinement times, reaching 0.17 s at the maximum, have shown a manner similar to the present scaling law derived from the existing medium sized helical devices, but improve on it by 50%. A distinguishing feature of a favorable dependence of energy confinement time on density remains in the present power density ( $\sim 40$  kW/m<sup>3</sup>) and the electron density ( $3 \times 10^{19}$  m<sup>-3</sup>) regimes unlike L-mode in tokamaks. Temperatures of both electrons and ions as high as 200 eV have been observed at the outermost flux surface, which indicates a qualitative jump in performance from the helical devices to date. Spontaneously generated toroidal currents agree with the physical picture of neoclassical bootstrap currents. Change of magnetic configuration due to finite- $\beta$  effect has been well described by the 3-D MHD equilibrium analysis. An escape of particles from the core region leading to a hollow density profile has been observed in hydrogen plasmas, which is mitigated through core fueling with a pellet injection or in helium discharges.

### 1. INTRODUCTION

The physics research goal of the Large Helical Device (LHD) [1,2] is to achieve and study currentless plasmas which can be extrapolated to the reactor condition in a heliotron configuration. The minor radius of 0.6 m is prerequisite to separate the core region confining high temperature plasmas from the boundary where atomic processes play an essential role. This paper describes the first experiments with regard to confinement studies under conditions that the helical devices had not satisfied to date. LHD has the major radius of 3.9 m and the operated magnetic field in this study is 1.5 T although superconducting coils have capability of 3 T operation [3]. Progress in experiments

of helical devices (CHS, Heliotron E, Wendelstein VII-AS, and ATF) in the last decade [4] has demonstrated performance equivalent to L-mode plasmas in tokamaks with similar dimensions and has indicated clear clues for further confinement improvement [5-7]. The first primary subject in this study lies in the clarification of the extrapolation of prior experience to large-scale plasmas which are sufficiently collisionless.

The magnetic geometry in this study is set at the optimized one from MHD stability, neoclassical transport and particle confinement. The magnetic axis is 3.75 m which is inward shifted from the machine center and the averaged elongation is 1. The plasma vacuum vessel which also functions as the first wall is stainless steel and is conditioned by discharge cleanings of He glow or ECR (2.45GHz) DC plasmas, and by Ti gettering with a coverage area of 20 %. The 2nd harmonic 84-GHz ECH up to 400 kW has been employed to generate plasmas, which are targets for NBI. The ray of ECH is well focused with waist sizes of 50 mm in the toroidal direction and 15 mm in the radial direction. Since the focal point is controllable by movable mirrors, the deposition profile effect on transport and kinetic effect of power deposition with relation to the magnetic geometry can be examined. Two negative-ion-based neutral beam injectors are aligned for tangential injection in opposing directions relative to the helical coil current (co- and counter-injections) to operate plasmas free from beam driven toroidal currents. Each injector is designed to inject a 7.5 MW hydrogen beams at an energy of 180keV for 10 s. One of the injectors is also designed to inject 100 keV beams of 0.5 MW up to 90 s. In the initial experiments, the ion sources are optimized for a lower energy up to 100 keV considering that the target plasma density is low under the 1.5 T operation. In this study the available injecting beam power is 3 MW in total. Profile measurements are available by YAG-laser Thomson scattering for the electron temperature, charge-exchange-recombination spectroscopy (CXRS) for the ion temperature and multi-chord FIR interferometer for the electron density.

## 2. CHARACTERISTICS OF PLASMA DISCHARGE

ECH produces target plasmas with a density of more than  $1 \times 10^{19} \text{ m}^{-3}$  so as to have the shine-through power of NBI less than 40%. Waveforms of a typical hydrogen discharge are shown in Fig.1(a). The plasma expands in the radial direction with increasing temperature and density during the initial 0.5 s. When the high-temperature plasma reaches the magnetic boundary, a divertor channel is formed. The events of a rapid pump-out of density, a decrease of radiation and bursts of divertor flux start with regard to this timing. The decay time of the decrease in density is enhanced by an increase of the heating power. Density does not build up efficiently despite strong gas puffing in the pump-out phase for several hundreds ms. A pellet injection is effective in increasing density as well as the stored energy, however, the decay constant of the density is not improved. Typical electron density, temperature and particle flux on the divertor plates in this phase are  $5 \times 10^{17} \text{ m}^{-3}$ , 10 to 20 eV and  $10^{22} \text{ m}^{-2}/\text{s}$ , respectively. The amplitude of the oscillation in the particle flux is about 50% and no obvious peaks in the power spectrum have been observed. In the discharges which are longer than 1 s, density clamping is terminated in the later phase. Since the stainless steel becomes occluded at around  $1 \times 10^{22}$  hydrogenic ions per  $\text{m}^2$ , this phenomena can be explained by degradation of pumping efficiency. The pump-out phenomena is much milder and density can be maintained by recycling instead of auxiliary gas puffing in helium discharges since the metallic wall does not pump helium (see Fig.1 (b)). Helium plasmas also mitigate the shine-through losses of NBI in the low density regime.

The present operational density limits bounded by radiation collapses are  $4.5 \times 10^{19} \text{ m}^{-3}$  in hydrogen plasmas with a gas puff,  $6.0 \times 10^{19} \text{ m}^{-3}$  in hydrogen plasmas with a pellet injection, and  $6.3 \times 10^{19} \text{ m}^{-3}$  for helium plasmas. The radiation collapse phenomena can be classified into two types. The first type is characterized by a hollow radiation profile that increased symmetrically up until the radiation collapse. In these instances the peak total radiated power is at or below the beam input power. In these discharges the stored energy drops steadily to zero in a time frame of 100 ms or more. The




FIG.1. Typical discharges of NBI heated plasmas. (a) Hydrogen. The pellet is injected at  $t = 0.81s$ . (b) Helium.




FIG.2. Evolution of electron density and temperature profiles. Density profiles are derived from Abel inversion from FIR chordal data. Temperatures are measured by the YAG laser (50Hz)-Thomson scattering system. (a) Density profile in a hydrogen discharge. (b) Temperature profile in a hydrogen discharge. (c) Density in a helium discharge. (d) Temperature in a helium discharge.

second type is characterized by a rapid increase in the total radiated power over a 100 ms time frame resulting in an instantaneous peak total radiated power, which is well in excess of the input heating power. The detector profile shows a sharp peaking on the inboard side of the plasma which increases and moves towards the center of the plasma in 30 ms.

Figure 2 illustrates the evolution of density and temperature profiles in the discharges shown in Fig.1. In a hydrogen discharge, a broad density profile becomes hollow when the pump-out starts. Pellet injection (3 mm in diameter and 3 mm in length, particle inventory of  $10^{21}$ ) has mitigated the hollow density profile. Broad density profile is maintained throughout a helium discharge and the edge density is higher than that in a hydrogen discharge, which is due to a larger neutral source from recycling. The temperature profile is more peaked in a hydrogen discharge than a helium one.

While hydrogen discharges are accompanied by the density pump-out, the low recycling condition which has been strongly related to confinement improvements has been realized. The temperatures of both electrons and ions have reached 200 eV at the outermost flux surface and the density is sharply bounded at the outside of the last closed flux surface (see Fig.3). Although a clear significant improvement of confinement has not been observed yet, the above mentioned density and temperature characteristics have not been observed in the helical devices to date and the increase of




FIG3. Typical radial profiles of ion and electron temperatures and electron density in the pump-out low-recycling phase.




FIG4. Change of poloidal rotation before ( $t = 0.4s$ ), at ( $t=0.6s$ ) and during ( $t=0.8s$ ) the event of the density clamping.

edge temperatures which is a key of confinement improvement would be expected by increased input power in further experiments.

The poloidal rotation which is closely connected to the radial electric field  $E_r$  has been measured by CXRS. A change in the direction of the rotation has been observed before and after the start of pump-out in a hydrogen discharge. Figure 4 indicates that the plasma rotates initially in the electron diamagnetic direction and then changes to the ion diamagnetic direction. This may reflect a change in the sign of  $E_r$  from negative to positive. Hollow density profiles are enhanced by the positive electric field [8] and a neoclassical theory has pointed out that this problem is pronounced in large-scale plasmas [9].

### 3. GLOBAL CONFINEMENT

Four major inter-machine scalings of energy confinement time for helical devices (LHD scaling [10], Lackner-Gottardi scaling [11], gyro-reduced Bohm [12] and International Stellarator Scaling 95 (ISS95) [13]) have been proposed so far and all describe similar manners of power deterioration and positive density dependences. The international stellarator scaling 95 (ISS95) expresses,

$$\tau_E = 0.079 a^{2.21} R^{0.65} P^{-0.59} \bar{n}_e^{0.51} B_i^{0.83} t_{23}^{0.4},$$

where  $\tau_E$  in s,  $a$  and  $R$  in m,  $P$  in MW,  $\bar{n}_e$  in  $10^{19} \text{m}^{-3}$ , and  $B_i$  in T. While this scaling describes L-mode plasmas in large tokamaks as well, density dependence is pronounced unlike in the tokamak L-mode scaling. From experience in tokamaks, larger devices which have less heating power density make the transition to saturation from linear ohmic confinement in the lower density regime, which is connected to an L-mode with auxiliary heating. This issue is crucial for the prospects of helical devices since the present reactor design relies on the favorable density dependence on confinement.

Energy confinement times are estimated by the stored energy measured by diamagnetic loops. The response of diamagnetic loops has been calculated by the 3-D magnetic field analysis with 3-D finite- $\beta$  equilibria. The NBI power deposition is modeled by the database from a number of runs of a




FIG.5. (a) Comparison of experimental energy confinement times with those predicted by ISS95. Data from helical device are adapted from the international stellarator database and data from tokamaks are adapted from the ITER L-mode database. (b) Deviation of experimental data from the scalings. LHD: the LHD scaling [10], L-G: the Lackner-Gottardi scaling [11], GRB: the gyro-reduced Bohm scaling [12] and ISS95: the international stellarator scaling [13].

3-D Monte-Carlo simulation code, MCNBI [14]. In heliotrons a large Shafranov shift occurs at finite  $\beta$  to change the beam birth profile and the drift orbit of beam particles. Therefore, in the code, the 3-D magnetic configuration including the finite- $\beta$  effect is used for the evaluation of the deposition profile. The computational results suggest that heating efficiency and deposition profiles are not different for the cases of co- and counter- injections in the current experimental condition. The stored energy derived from kinetic measurements agrees with the diamagnetic measurement, which is consistent with the fact that results from the NBI power deposition model suggest a negligible effect from the beam pressure (at most a few %).

Figure 5 (a) shows the comparison of experimental data with the prediction from the ISS95 expression. The density and power absorption ranges cover  $1.0 - 5.0 \times 10^{19} \text{ m}^{-3}$  and  $0.5 - 2.3 \text{ MW}$ . The data in hydrogen discharges have been adopted here, since the estimate of power deposition in helium discharges cannot be discussed with the same accuracy as in hydrogen now. While the trend of energy confinement times of NBI heated plasmas in LHD has a manner similar to ISS95, experimental data is systematically better than ISS95 by a factor of 1.5. Deviation from the scalings indicates an enhancement of a factor of 2 from the LHD scaling and is comparable to the L-G scaling (see Fig. 5 (b)).

High density plasmas with a pellet injection and one NBI have almost the same absorption power as low density ones with two NBIs. Comparison of those sequences indicates that energy confinement is improved with the square root of the density in the accessible the density range ( $3 \times 10^{19} \text{ m}^{-3}$ ) with a power density of  $40 \text{ kW/m}^3$ .

#### 4. MHD CHARACTERISTICS

The MHD equilibrium properties have been well described by 3-D finite- $\beta$  equilibrium calculation in helical devices [15]. Since the volume averaged beta  $\langle \beta \rangle$  has reached 0.7%, The theory predicts the Shafranov shift of 14 cm on the prolate cross-section and 23 cm on the oblate one. Figure 6 shows the shift of the peak of the soft X-ray profile which corresponds to the magnetic axis. Since the increase of  $\beta$  is primarily attributed to the increase in density, the pressure profile is flattened in the higher  $\beta$  region in accordance with broadening of the density profile. The experimental estimates agree with the finite- $\beta$  calculation very well. Also, a similar change of plasma profiles has been observed in the density and the electron temperature.




FIG6. Shift in peak of the soft-X ray profile. Lines indicate estimates from the 3-D MHD computation. Solid line : assume a standard profile from  $p=p_0(1-\rho^2)^2$ . Broken line : assume a broad profile from  $p=p_0(1-\rho^2)(1-\rho^8)$ .

No internal relaxation phenomena have been observed in the core region so far. Spontaneous generation of a magnetic well due to the Shafranov shift realizes Mercier (ideal interchange mode) stable configuration leading to the equilibrium limit.

## 5. GENERATION OF TOROIDAL CURRENTS

Research in both tokamaks and helical devices [16,17] has confirmed neoclassical bootstrap currents. The magnetic field of LHD is excited in a real steady state with using superconducting coils. Therefore, the applied voltage is extremely low and ohmic currents are absolutely excluded. The capability for long-pulse discharges can also eliminate the ambiguity of time dependent behaviors. A precise investigation will progress in LHD making full use of these advantages. A toroidal current of up to 50 kA has been observed in a discharge with co-injected NBI of 1.5MW while the equivalent currents to generate the rotational transform in the vacuum magnetic field of LHD is 700 kA at 1.5 T. It takes more than 2 s for currents to reach a steady state, which is comparable to the L/R time of the plasma. Figure 7 (a) shows the dependence of the toroidal currents on density. These currents result from the synergetic effect of bootstrap currents and Ohkawa currents. All currents flow in the paramagnetic direction which increase the rotational transform. Since the bootstrap current is expected to flow in the paramagnetic direction, the total currents shift to this side. Observed currents with co-injection decrease in the high density regime which is explained by the degradation of efficiency of the Ohkawa current. The Ohkawa current is cancelled out in the case of balanced injection and bootstrap current is pronounced. Figure 7 (b) indicates that the toroidal current increases in proportion with the stored energy which reflects pressure gradient. A neoclassical estimate [18] bounds the upper limit of the experimental observation, which is explained by a discharge duration which is too short to ramp the full current.




FIG7. (a) Toroidal currents as a function of density. (b) Toroidal currents as a function of stored energy. Only the case of balanced injection is illustrated here for NBI. Since the ray of ECH is injected perpendicularly to the magnetic field, no momentum input exists. Lines: prediction from the neoclassical theory by scanning the density profile  $n_e = n_{e0}(1-\rho^{-8})$  with the fixed temperature profile  $T = (1-\rho^{-2})$  (keV).

## 6. DISCUSSIONS AND SUMMARY

The initial experiments on the Large Helical Device (LHD) have extended confinement studies on currentless plasmas to a large scale. An outlook of confinement of NBI heated plasmas has shown



a trend similar to the prior scaling laws, however, a significant improvement compared with the ISS95 expression by a factor of 1.5 and comparable data to the Lackner-Gottardi scaling have been obtained. In the present experimental condition, an absorbed power of NBI depends on the operational density. Therefore, it needs a careful attention and further systematic scans to separate the power degradation from a favorable density dependence. Comparison with two kinds of experimental set-up with the same absorbed power, i.e., high density discharges heated by 1 beam with a pellet injection and low density discharges heated by 2 beams is consistent with the prediction that confinement is improved with the square root of density.

Fueling of particle is reconfirmed as an important operational issue in the experiment. An escape of particles from the core of the plasma has been highlighted in hydrogen discharges. Low recycling condition is achieved simultaneously, which leads to high temperatures at the edge. This observation is explained from the concept of the high temperature divertor [19]. Combination with central fueling by pellet injection has mitigated a hollow density profile temporally. Helium discharges facilitates particle control through high recycling. Here it should be noted that the high edge temperature is not specific to hydrogen discharges. A careful control of helium gas puffing can realize a similar high edge temperature.

Heating by NBI of 3 MW has realized plasmas with a fusion triple product of  $8 \times 10^{18}$  keV m<sup>3</sup> s at the magnetic field of 1.5T. An electron temperature of 1.5 keV and an ion temperature of 1.1 keV have achieved simultaneously at the line-averaged electron density of  $1.5 \times 10^{19}$  m<sup>-3</sup>. At the density as low as  $4 \times 10^{18}$  m<sup>-3</sup>, the central ion temperature reaches 1.4 keV.

Spontaneous generation of toroidal currents has been observed, characteristics of which agree with neoclassical bootstrap currents. Changes of the plasma profile have been well described by the 3-D MHD equilibrium analysis.

## ACKNOWLEDGEMENTS

The authors are grateful to all collaboration with universities and institutes. Efforts by the industrial companies to complete the LHD construction are greatly appreciated.

## REFERENCES

- [1] IYOSHI, A., et al., these Proceedings.
- [2] FUJIWARA, M., et al., *J. Fusion Energy* **15** (1996)7.
- [3] MOTOJIMA, O., et al., these Proceedings.
- [4] IYOSHI, A., in *Plasma Physics and Controlled Nuclear Fusion Research 1996* (Proc.16th Conf. Montreal, 1996), Vol.1, IAEA, Vienna (1997) 113.
- [5] TOI, K., et al., in *Plasma Physics and Controlled Nuclear Fusion Research 1992* (Proc.14th Conf. Würzburg, 1992), Vol.2, IAEA, Vienna (1993) 461.
- [6] ERCKMANN, V., et al., *Phys. Rev. Lett.* **70** (1993) 2086
- [7] IDA, K., et al., *Phys. Rev. Lett.* **76** (1996) 1268.
- [8] IDEL, H., et al., *Phys. Plasmas* **1** (1994) 3400.
- [9] MAASSBERG, H., BEIDLER, C.D., SIMMET, E.E., *J. Plasma Fusion Res. SERIES*, Vol.1 (1998) 230.
- [10] SUDO, S., et al., *Nucl. Fusion* **30** (1990)11.
- [11] LACKNER, K., GOTTARDI, N.A.O., *Nucl. Fusion* **30** (1990) 767.
- [12] MURAKAMI, M., et al., in *Plasma Physics and Controlled Nuclear Fusion Research 1992* (Proc.14th Conf. Würzburg, 1992), Vol.2, IAEA, Vienna (1993) 391.
- [13] STROTH, U., et al., *Nucl. Fusion* **36** (1996) 1063.
- [14] MURAKAMI, S., et al., *Trans. Fusion Tech.* **27** (1995) 256.
- [15] YAMADA, H., et al., *Nucl. Fusion* **32** (1992) 25.
- [16] MURAKAMI, M., et al., *Phys. Rev. Lett.* **66**(1991)707.
- [17] YAMADA, H., et al., *Nucl. Fusion* **34** (1994) 641.
- [18] WATANABE, K.Y., et al., *Nucl. Fusion* **35** (1995) 335.
- [19] OHYABU, N., et al., *Nucl. Fusion* **34** (1994) 387.

## Recent Issues of NIFS Series

- NIFS-506 H. Kitauchi and S. Kida,  
*Intensification of Magnetic Field by Concentrate-and-Stretch of Magnetic Flux Lines*, Sep. 1997
- NIFS-507 R.L. Dewar,  
*Reduced form of MHD Lagrangian for Ballooning Modes*, Sep. 1997
- NIFS-508 Y.-N. Nejoh,  
*Dynamics of the Dust Charging on Electrostatic Waves in a Dusty Plasma with Trapped Electrons*, Sep. 1997
- NIFS-509 E. Matsunaga, T. Yabe and M. Tajima,  
*Baroclinic Vortex Generation by a Comet Shoemaker-Levy 9 Impact*, Sep. 1997
- NIFS-510 C.C. Hegna and N. Nakajima,  
*On the Stability of Mercier and Ballooning Modes in Stellarator Configurations*, Oct. 1997
- NIFS-511 K. Ono and T. Hatan, ~~Hatan~~,  
*Rotation and Oscillation of Nonlinear Dipole Vortex in the Drift-Unstable Plasma*, Oct. 1997
- NIFS-512 J. Uramoto,  
*Clear Detection of Negative Pionlike Particles from  $H_2$  Gas Discharge in Magnetic Field*, Oct. 1997
- NIFS-513 T. Shimozuma, M. Sato, Y. Takita, S. Ito, S. Kubo, H. Idei, K. Ohkubo, T. Watari, T.S. Chu, K. Felch, P. Cahalan and C.M. Loring, Jr.,  
*The First Preliminary Experiments on an 84 GHz Gyrotron with a Single-Stage Depressed Collector*, Oct. 1997
- NIFS-514 T. Shimozuma, S. Morimoto, M. Sato, Y. Takita, S. Ito, S. Kubo, H. Idei, K. Ohkubo and T. Watan, ~~Watan~~,  
*A Forced Gas-Cooled Single-Disk Window Using Silicon Nitride Composite for High Power CW Millimeter Waves*, Oct. 1997
- NIFS-515 K. Akaishi,  
*On the Solution of the Outgassing Equation for the Pump-down of an Unbaked Vacuum System*, Oct. 1997
- NIFS-516 *Papers Presented at the 6th H-mode Workshop (Seeon, Germany)*, Oct. 1997
- NIFS-517 John L. Johnson,  
*The Quest for Fusion Energy*, Oct. 1997
- NIFS-518 J. Chen, N. Nakajima and M. Okamoto,  
*Shift-and-Inverse Lanczos Algorithm for Ideal MHD Stability Analysis*, Nov. 1997
- NIFS-519 M. Yokoyama, N. Nakajima and M. Okamoto,  
*Nonlinear Incompressible Poloidal Viscosity in  $L=2$  Heliotron and Quasi-Symmetric Stellarators*, Nov. 1997
- NIFS-520 S. Kida and H. Miura,  
*Identification and Analysis of Vortical Structures*, Nov. 1997
- NIFS-521 K. Ida, S. Nishimura, T. Minami, K. Tanaka, S. Okamura, M. Osakabe, H. Idei, S. Kubo, C. Takahashi and K. Matsuoka,  
*High Ion Temperature Mode in CHS Heliotron/torsatron Plasmas*, Nov. 1997
- NIFS-522 M. Yokoyama, N. Nakajima and M. Okamoto,  
*Realization and Classification of Symmetric Stellarator Configurations through Plasma Boundary Modulations*, Dec. 1997
- NIFS-523 H. Kitauchi,  
*Topological Structure of Magnetic Flux Lines Generated by Thermal Convection in a Rotating Spherical Shell*, Dec. 1997
- NIFS-524 T. Ohkawa,  
*Tunneling Electron Trap*, Dec. 1997

- NIFS-525 K. Itoh, S.-I. Itoh, M. Yagi, A. Fukuyama,  
*Solitary Radial Electric Field Structure in Tokamak Plasmas*; Dec. 1997
- NIFS-526 Andrey N. Lyakhov,  
*Alfven Instabilities in FRC Plasma*; Dec. 1997
- NIFS-527 J. Uramoto,  
*Net Current Increment of negative Muonlike Particle Produced by the Electron and Positive Ion Bunch-method*; Dec. 1997
- NIFS-528 Andrey N. Lyakhov,  
*Comments on Electrostatic Drift Instabilities in Field Reversed Configuration*; Dec. 1997
- NIFS-529 J. Uramoto,  
*Pair Creation of Negative and Positive Pionlike (Muonlike) Particle by Interaction between an Electron Bunch and a Positive Ion Bunch*; Dec. 1997
- NIFS-530 J. Uramoto,  
*Measuring Method of Decay Time of Negative Muonlike Particle by Beam Collector Applied RF Bias Voltage*; Dec. 1997
- NIFS-531 J. Uramoto,  
*Confirmation Method for Metal Plate Penetration of Low Energy Negative Pionlike or Muonlike Particle Beam under Positive Ions*; Dec. 1997
- NIFS-532 J. Uramoto,  
*Pair Creations of Negative and Positive Pionlike (Muonlike) Particle or K Mesonlike (Muonlike) Particle in H<sub>2</sub> or D<sub>2</sub> Gas Discharge in Magnetic Field*; Dec. 1997
- NIFS-533 S. Kawata, C. Boonmee, T. Teramoto, L. Drska, J. Limpouch, R. Liska, M. Sinor,  
*Computer-Assisted Particle-in-Cell Code Development*; Dec. 1997
- NIFS-534 Y. Matsukawa, T. Suda, S. Ohnuki and C. Namba,  
*Microstructure and Mechanical Property of Neutron Irradiated TiNi Shape Memory Alloy*; Jan. 1998
- NIFS-535 A. Fujisawa, H. Iguchi, H. Idei, S. Kubo, K. Matsuoka, S. Okamura, K. Tanaka, T. Minami, S. Ohdachi, S. Monta, H. Zushi, S. Lee, M. Osakabe, R. Akiyama, Y. Yoshimura, K. Toi, H. Sanuki, K. Itoh, A. Shimizu, S. Takagi, A. Ejiri, C. Takahashi, M. Kojima, S. Hidekuma, K. Ida, S. Nishimura, N. Inoue, R. Sakamoto, S.-I. Itoh, Y. Hamada, M. Fujiwara,  
*Discovery of Electric Pulsation in a Toroidal Helical Plasma*; Jan. 1998
- NIFS-536 Lj.R. Hadzievski, M.M. Skoric, M. Kono and T. Sato,  
*Simulation of Weak and Strong Langmuir Collapse Regimes*; Jan. 1998
- NIFS-537 H. Sugama, W. Horton,  
*Nonlinear Electromagnetic Gyrokinetic Equation for Plasmas with Large Mean Flows*; Feb. 1998
- NIFS-538 H. Iguchi, T.P. Crowley, A. Fujisawa, S. Lee, K. Tanaka, T. Minami, S. Nishimura, K. Ida, R. Akiyama, Y. Hamada, H. Idei, M. Isobe, M. Kojima, S. Kubo, S. Monta, S. Ohdachi, S. Okamura, M. Osakabe, K. Matsuoka, C. Takahashi and K. Toi,  
*Space Potential Fluctuations during MHD Activities in the Compact Helical System (CHS)*; Feb. 1998
- NIFS-539 Takashi Yabe and Yan Zhang,  
*Effect of Ambient Gas on Three-Dimensional Breakup in Coronet Formation Process*; Feb. 1998
- NIFS-540 H. Nakamura, K. Ikeda and S. Yamaguchi,  
*Transport Coefficients of InSb in a Strong Magnetic Field*; Feb. 1998
- NIFS-541 J. Uramoto,  
*Development of  $v_{\mu}$  Beam Detector and Large Area  $v_{\mu}$  Beam Source by H<sub>2</sub> Gas Discharge (I)*; Mar. 1998
- NIFS-542 J. Uramoto,  
*Development of  $\bar{v}_{\mu}$  Beam Detector and Large Area  $\bar{v}_{\mu}$  Beam Source by H<sub>2</sub> Gas Discharge (II)*; Mar. 1998
- NIFS-543 J. Uramoto,

*Some Problems inside a Mass Analyzer for Pions Extracted from a H<sub>2</sub> Gas Discharge*, Mar 1998

- NIFS-544 J Uramoto,  
*Simplified  $\nu_{\mu}$  bar[ $\nu$ ] <sub>$\mu$</sub>  Beam Detector and  $\nu_{\mu}$  bar[ $\nu$ ] <sub>$\mu$</sub>  Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch*, Mar. 1998
- NIFS-545 J Uramoto,  
*Various Neutrino Beams Generated by D<sub>2</sub> Gas Discharge*, Mar 1998
- NIFS-546 R Kanno, N Nakajima, T Hayashi and M Okamoto,  
*Computational Study of Three Dimensional Equilibria with the Bootstrap Current*, Mar 1998
- NIFS-547 R Kanno, N. Nakajima and M. Okamoto,  
*Electron Heat Transport in a Self-Similar Structure of Magnetic Islands*; Apr 1998
- NIFS-548 J.E. Rice,  
*Simulated Impurity Transport in LHD from MIST*, May 1998
- NIFS-549 M M Skoric, T. Sato, A.M. Maluckov and M S Jovanovic,  
*On Kinetic Complexity in a Three-Wave Interaction*; June 1998
- NIFS-550 S Goto and S Kida,  
*Passive Saclar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation*; June 1998
- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,  
*Initial Value Problem of the Toroidal Ion Temperature Gradient Mode* ; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watan,  
*Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device* , June 1998
- NIFS-553 N. Noda, K. Tsuzuku, A. Sagara, N. Inoue, T. Muroga,  
*oronaization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-*. July 1998
- NIFS-554 S. Murakami and H. Saleem,  
*Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations*; July 1998
- NIFS-555 H. Nakamura , K. Ikeda and S. Yamaguchi,  
*Physical Model of Nernst Element*, Aug 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,  
*Numerical Computation of Thermoelectric and Thermomagnetic Effects*, Aug 1998
- NIFS-557 Y. Takeiri, M. Osakabe, K. Tsumon, Y. Oka, O. Kaneko, E. Asano, T. Kawamoto, R. Akiyama and M. Tanaka,  
*Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System*; Aug 1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,  
*Molecular Dynamics of Structure Organization of Polyampholytes*; Sep 1998
- NIFS-559 R. Horiuchi, K. Nishimura and T. Watanabe,  
*Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations*, Sep 1998  
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Kholopenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,  
*Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection*; Oct 1998  
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,  
*Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron*, Oct 1998

(IAEA-CN-69/EX5/1)

- NIFS-562 S. Yoshikawa,  
*Next Generation Toroidal Devices*; Oct 1998
- NIFS-563 Y. Todo and T. Sato,  
*Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes*; Oct. 1998  
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watari, T. Shimozuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Idei, Y. Oka, M. Osakabe, T. Seki, K. Tsumori, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Shimpō, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Morimoto, T. Bigelow and Y.P. Zhao.  
*Steady State Heating Technology Development for LHD*; Oct. 1998  
(IAEA-CN-69/FTP/21)
- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyabu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda,  
*LHD-Type Compact Helical Reactors*; Oct 1998  
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,  
*Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems*; Oct. 1998  
(IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D.S. Darrow, H. Idei, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tsumori, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watari,  
*Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas*; Oct 1998  
(IAEA-CN-69/EX2/2)
- NIFS-568 S. Okamura, K. Matsuoka, R. Akiyama, D.S. Darrow, A. Ejiri, A. Fujisawa, M. Fujiwara, M. Goto, K. Ida, H. Idei, H. Iguchi, N. Inoue, M. Isobe, K. Itoh, S. Kado, K. Khlopenkov, T. Kondo, S. Kubo, A. Lazaros, S. Lee, G. Matsunaga, T. Minami, S. Morita, S. Murakami, N. Nakajima, N. Nikai, S. Nishimura, I. Nomura, S. Ohdachi, K. Ohkuni, M. Osakabe, R. Pavlichenko, B. Peterson, R. Sakamoto, H. Sanuki, M. Sasao, A. Shimizu, Y. Shirai, S. Sudo, S. Takagi, C. Takahashi, S. Takayama, M. Takechi, K. Tanaka, K. Toi, K. Yamazaki, Y. Yoshimura and T. Watari,  
*Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS*; Oct 1998  
(IAEA-CN-69/OV4/5)
- NIFS-569 M.M. Skoric, T. Sato, A. Maluckov, M.S. Jovanovic,  
*Micro- and Macro-scale Self-organization in a Dissipative Plasma*; Oct 1998
- NIFS-570 T. Hayashi, N. Mizuguchi, T.-H. Watanabe, T. Sato and the Complexity Simulation Group,  
*Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak*; Oct 1998  
(IAEA-CN-69/TH3/3)
- NIFS-571 A. Iiyoshi, A. Komori, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Monsaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, H. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, S. Ima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, M. Fujiwara and O. Motojima,  
*An Overview of the Large Helical Device Project*; Oct 1998  
(IAEA-CN-69/OV1/4)
- NIFS-572 M. Fujiwara, H. Yamada, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Monsaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B. J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, M. Ima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iiyoshi,  
*Plasma Confinement Studies in LHD*; Oct. 1998  
(IAEA-CN-69/EX2/3)