



INTERNATIONAL ATOMIC ENERGY AGENCY

17th IAEA Fusion Energy Conference  
Yokohama, Japan, 19 - 24 October 1998

IAEA-CN-69/EXP1/19

## NATIONAL INSTITUTE FOR FUSION SCIENCE

### Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas

K. Toi, M. Takechi, S. Takagi, G. Matsunaga, M. Isobe, T. Kondo, M. Sasao, D.S. Darrow, K. Ohkuni, S. Ohdachi, R. Akiyama, A. Fujisawa, M. Gotoh, H. Idei, K. Ida, H. Iguchi, S. Kado, M. Kojima, S. Kubo, S. Lee, K. Matsuoka, T. Minami, S. Morita, N. Nikai, S. Nishimura, S. Okamura, M. Osakabe, A. Shimizu, Y. Shirai, C. Takahashi, K. Tanaka, T. Watari and Y. Yoshimura

(Received - Oct. 8, 1998 )

NIFS-574

Oct. 1998

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

## RESEARCH REPORT NIFS Series

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of this document, changes of substance or detail may have to be made before publication. The preprint is made available for the understanding that it will not be cited in the literature and that any use of the report must be done with the view expressed and the statements made remain the responsibility of the author(s). The views do not necessarily reflect those of the government of the host state. However, the views do not reflect those of the organizations sponsoring this meeting cannot be held responsible for any errors introduced in this preprint.

NAGOYA, JAPAN



INTERNATIONAL ATOMIC ENERGY AGENCY

17th IAEA Fusion Energy Conference  
Yokohama, Japan, 19 - 24 October 1998

---

IAEA-CN-69/

IAEA-F1-CN-69/EXP1/19

## GLOBAL MHD MODES EXCITED BY ENERGETIC IONS IN HELITRON/TORSATRON PLASMAS

K. TOI, M. TAKECHI<sup>1)</sup>, S. TAKAGI<sup>1)</sup>, G. MATSUNAGA<sup>1)</sup>, M. ISOBE,  
T. KONDO<sup>2)</sup>, M. SASAO, D.S. DARROW<sup>3)</sup>, K. OHKUNI<sup>1)</sup>, S. OHDACHI,  
R. AKIYAMA, A. FUJISAWA, M. GOTOH, H. IDEI, K. IDA,  
H. IGUCHI, S. KADO, M. KOJIMA, S. KUBO, S. LEE, K. MATSUOKA,  
T. MINAMI, S. MORITA, N. NIKAI<sup>1)</sup>, S. NISHIMURA, S. OKAMURA,  
M. OSAKABE, A. SHIMIZU<sup>1)</sup>, Y. SHIRAI<sup>2)</sup>, C. TAKAHASHI,  
K. TANAKA, T. WATARI, Y. YOSHIMURA

National Institute for Fusion Science, Toki-shi 509-5292, Japan

1) Dep. Energy Eng. Science, Nagoya Univ., Nagoya 464-8603, Japan

2) The Graduate Univ. for Advanced Studies, Toki-shi, 509-5292, Japan

3) Princeton Plasma Physics Laboratory, Princeton, N.J., U.S.A.

---

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The view expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designated Member State(s). In particular, the organizations sponsoring this meeting cannot be held responsible for any material reproduced in this preprint.

## GLOBAL MHD MODES EXCITED BY ENERGETIC IONS IN HELITRON/TORSATRON PLASMAS

### Abstract

In NBI heated plasmas of the CHS heliotron/torsatron where the beam driven current with a peaked profile is induced,  $m=2/n=1$  and  $m=3/n=2$  ( $m, n$  : poloidal and toroidal mode numbers) fishbone instabilities are observed for the first time. Pulsed increase in energetic ion loss flux is detected by an escaping ion probe during the  $m=3/n=2$  mode. The sawtooth crash is often induced by the  $m=2/n=1$  mode. The current driven internal kink mode is thought to be destabilized by the presence of energetic ions, having a character of pressure driven interchange mode. Toroidal Alfvén eigenmodes (TAEs) with  $n=1$  and  $n=2$  are also identified for the first time in NBI heated plasmas. TAEs localize near the plasma core region with fairly low magnetic shear realized by the small net plasma current, and do not lead to enhanced loss of energetic ions because of low fluctuation level.

### 1. INTRODUCTION

MHD instabilities such as fishbone instabilities [1] and Alfvén eigenmodes [2] driven by the presence of energetic ions are observed in a tokamak plasma heated by intense neutral beam and/or ICRF heating, and often lead to enhanced loss of energetic ions before their thermalization. In a tokamak fusion reactor, these kinetically driven modes, in particular, toroidal Alfvén eigenmodes (TAEs) might be excited by energetic alpha particles and considerably degrade their confinement [3,4]. Therefore, excitation and suppression of TAEs is intensively studied in large tokamaks. This issue is also very important for helical systems [5]. In particular, energetic ion loss might be considerably enhanced by these MHD modes in a helical system with appreciable magnetic field ripple. Recently, we observed fishbone instabilities (FBs) [6] and toroidal Alfvén eigenmodes (TAEs) [7] excited by energetic ions in the CHS heliotron/torsatron [8].

FBs in a tokamak plasma are excited by resonant interaction of  $m=1/n=1$  internal kink mode with the precessional motion of trapped ions. In contrast to the tokamak plasma, the CHS plasma is basically current-free and is confined in the three dimensional magnetic configuration. Therefore, main purpose of this research is to clarify the relevant MHD mode and the resonant orbit of energetic ions. On the other hand, TAEs can be destabilized by resonant interaction of Alfvén waves with circulating motion of passing ions or bounce motion of trapped ions in the toroidicity induced Alfvén spectrum gap. In a tokamak, the radial variation of the TAE frequency  $f_{TAE} = V_A / (4\pi R q)$  is fairly small ( $V_A$ : Alfvén velocity,  $q$ : safety factor), and then continuum damping of TAEs near the plasma edge might not act efficiently. On the other hand, the TAE frequency in CHS rapidly increases towards the edge, because  $q$  decreases towards the edge. This means that strong continuum damping of TAEs could take place near the edge. However, TAEs are obviously excited on the certain experimental conditions of CHS. In this paper we describe characteristics of FBs and TAEs observed in CHS.

### 2. EXPERIMENTAL SET-UP AND RESULTS

Main diagnostics for this study are a movable magnetic probe array, a soft X-ray detector (SX-) array, an escaping ion probe (EIP) [9] and heavy ion beam probe (HIBP) [10]. To enlarge contribution of energetic ions to the bulk plasma, the electron density is kept low ( $\leq 3 \times 10^{19} \text{m}^{-3}$ ). Typically, averaged (parallel) beta of energetic beam ions is in  $\sim 0.1-0.3\%$  and is comparable to the bulk plasma beta. Neutral beams are injected nearly parallel to the toroidal field and induce appreciable net plasma current dominated by beam driven current.

#### 2.1 Fishbone instabilities

Characteristics of FBs in CHS depend upon the position of magnetic axis in the vacuum field  $R_{ax}$ , where the plasma of  $R_{ax} \leq 0.92 \text{m}$  is referred as the "inward-shifted" plasma and  $R_{ax} \geq 0.95 \text{m}$  as the "outward-shifted" one. In the outward-shifted plasmas,  $m=3/n=2$  FBs are observed inducing enhanced loss of energetic ions. In the inward-shifted plasmas,  $m=2/n=1$  FBs are observed. In the initial experiments,  $m=2/n=1$  FBs were called "burst modes" [11].

Figure 1(a) shows time evolution of magnetic fluctuations of  $m=3/n=2$  FBs and energetic ion loss flux in the outward-shifted plasma (i.e.,  $R_{ax} = 0.95 \text{m}$ ) [6]. As seen from Fig. 1(a), magnetic fluctuations exhibit the fishbone like amplitude modulation and quickly suppressed after turn-off of the NBI pulse. Energetic ion loss is transiently enhanced by FB-burst. Figure 1(b)

shows the expanded time trace of magnetic fluctuations and energetic ion loss flux shown in Fig.1(a). The ion loss is suddenly enhanced as soon as the magnetic fluctuations reach the peak level. Recently, it has been clarified by changing the aperture direction of EIP that energetic ions with the pitch angle of  $\chi \sim 90^\circ$  and the energy  $E_p \sim 25$  keV are preferentially expelled by  $m=3/n=2$  FBs, while in the previous experiment ion loss flux at  $\chi \sim 45^\circ$  and  $E_p \sim 30$  keV was clearly enhanced by the FBs[12]. Note that these energetic ions are slightly slowed down from the injection energy  $E \sim 35-40$  keV. On the hand, time behaviours of  $m=2/n=1$  FBs in  $R_{ax}=0.92$  m are shown in Fig.2. In this case, obvious response in EIP signal to FBs is not observed, while floating potential measured by Langmuir probe near the last closed flux surface(LCFS) exhibits the pulsed increase. The amplitude of magnetic fluctuations grows in the rising phase of the net plasma current and then the sawtooth oscillations characterized by the annular crash at the  $q=2$  surface are induced in the latter half of the discharge. Both of  $m=3/n=2$  and  $m=2/n=1$  FBs show rapid frequency sweep by about factor of two in each burst(Fig.3). This frequency sweep is not caused by the change of the plasma potential near the rational surface. Appearance of FBs is very sensitive for the net plasma current. Radial profiles of SX-fluctuations related to FBs have two peaks in the region of  $\rho < 2/3$ , exhibiting ballooning nature(Fig.4). The largest peak is located near the inner rational surface. This suggests that two rational surfaces related to FBs are produced by NBI driven current with a peaked profile. It is thought that the current driven internal kink is destabilized preferentially at the inner rational surface in the  $q$ -profile with double rational surface, having a character of pressure driven interchange mode at the outer rational surface[13]. If the orbits of toroidally and/or helically trapped ions are assumed to be equivalent for the banana orbit in a tokamak, the toroidal precession frequency of trapped ions may be approximated by  $f_{pre} \sim (m/n)E[1-(R_t/R)^2]/(2\pi Z r_s R B_t)$ , where  $E$ ,  $R_t$ ,  $r_s$  and  $Z$  are the injection beam energy, the tangency radius of the beam line, the size of the rational surface and the charge of injected beam, respectively. If the size of the inner rational surface is adopted, the resonance frequency  $n f_{pre}$  is estimated to be  $\sim 90$  kHz for  $m=3/n=2$  FBs (Fig.3) and  $\sim 40$  kHz for  $m=2/n=1$  FBs. These values are close to the observed frequencies at the beginning of each burst.

## 2.2. Toroidal Alfvén Eigenmodes

In TAE experiments, hydrogen beam is injected into a hydrogen plasma, where the toroidal field, line averaged electron density and beam energy are scanned in the range of  $B_t=0.7 - 1.5$  T,  $n_e=0.5 - 3 \times 10^{19} \text{m}^{-3}$  and  $E=28-40$  keV, respectively. In the condition,  $n=1$  and  $n=2$  TAEs having very narrow frequency band width ( $\leq 1$  kHz) are observed in inward-shifted( $R_{ax}=0.92\text{m}$ ) and outward-shifted( $R_{ax}=0.95\text{m}$ ) plasmas[7]. Excitation of TAEs are sensitive for a small net plasma current which decreases the magnetic shear near the plasma core region. Figure 5 summarizes the observed frequencies of TAEs as a function of the calculated TAE frequency for pure hydrogen plasma  $f_{TAE}(0)$ , where the electron density at the plasma center is employed and  $q = (m+1/2)/n$ . As seen from Fig.5, the observed frequencies are in proportion to the TAE frequency. However, the values are by about 30% lower than the TAE frequency. In these plasmas,  $Z_{eff}=2-3$  and then impurity ions decrease  $f_{TAE}(0)$  by about 15%. On the other hand, since the TAE gap position is expected to be  $r=0.3-0.4$  typically, this effect increases  $f_{TAE}(0)$  by  $\sim 5\%$ . Therefore, the observed frequencies are still by  $\sim 20\%$  lower than  $f_{TAE}(0)$ . The TAE frequency and TAE gap position are approximately predicted by simple calculation of the Alfvén continua in a cylindrical configuration. Figure 6(a) shows the (uncoupled) Alfvén continua for  $n=1$  and  $n=2$  TAEs observed in CHS, where the above-mentioned impurity effect is taken into account in the calculation. This figure also shows the TAE gap structure calculated by the approximate expression for large-aspect-ratio and low beta tokamak [3]. Here, the rotational transform( $1/q$ ) is assumed to be the sum of the rotational transform due to the net plasma current and the external rotational transform in three dimensional current-free equilibrium with 0.2% averaged total beta. Here, the current density profile is assumed to be a plausible shape of  $j_\phi = j_0(1-\rho^2)^2$ . As seen from Fig.6, the observed frequency ( $\sim 97$  kHz) lies near the lower boundary of the innermost TAE gap. Internal structure of TAEs is measured by the SX array. Figure 6(b) shows the radial profiles of coherence  $\gamma$  between SX signal and magnetic probe signal for the observed  $n=1$  TAEs. This figure shows that high coherence region is in the core region( $\rho \leq 0.6$ ), where  $\gamma$  for noise is  $\sim 0.3$ . Note that high coherence observed around the plasma center is caused by the integral

effect of SX emission along the line of sight. Therefore, TAEs are predicted to be localized around  $\rho \sim 0.2-0.6$ , which is also confirmed by measurement of plasma potential fluctuations measured by HIBP[7]. This result is consistent with Fig.6(a). The core localization of TAEs seems to be similar to the core localized TAEs in a tokamak[14]. In addition, TAEs in CHS are excited when the ratio of the beam velocity to the central Alfvén velocity  $V_b/V_A$  exceeds about 0.5. This suggests the side-band excitation of TAEs[15].


### 3. SUMMARY

In the CHS heliotron/torsatron,  $m=2/n=1$  and  $m=3/n=2$  fishbone instabilities (FBs) are observed in NBI heated plasmas where the beam driven current is induced with a peaked profile. Energetic ion loss flux is transiently enhanced by FBs. In each FB-burst the frequency of the magnetic fluctuations is considerably swept down. Radial profiles of the fluctuations suggest that the current driven internal kink mode is destabilized by the presence of energetic ions, and has a character of pressure driven interchange mode with increase in the plasma beta. However, these FBs are easily suppressed by balanced or counter NBI heating. In addition,  $n=1$  and  $n=2$  TAEs are also identified for the first time in NBI heated plasmas. The modes are only observed when the ratio of beam velocity to the Alfvén one exceeds 0.5 and a net plasma current induced by co-NBI is in the required level depending on  $B_t$ . The observed TAEs localize near the plasma core region with fairly low magnetic shear realized by the net plasma current. TAEs in CHS do not enhance energetic ion loss, because the fluctuation level is low ( $b_\theta/B_t < 10^{-5}$  at LCFS).

The authors wish to acknowledge N. Nakajima (NIFS), Y. Nakamura (Kyoto Univ.), M. Wakatani (Kyoto Univ.), and J.W. Van Dam (Univ. of Texas) for fruitful discussions.

### REFERENCES

- [1] MCGUIRE, K., et al., Phys. Rev. Lett. **50**(1983) 891.
- [2] WONG, K.L., et al., Phys. Rev. Lett. **66**(1991) 1874.
- [3] CHENG, C.Z., CHANCE, M.S., Phys. Fluids **29**(1986)3695.
- [4] FU, G.Y., VAN, DAM, J.W., Phys. Fluids **B1**(1989)1949.
- [5] WELLER, A., et al., Phys. Rev. Lett. **72**(1994)1220.
- [6] TOI, K., et al., J. Plasma Fusion Res. SERIES **1**(1998)267.
- [7] TAKECHI, M., TOI, K. et al., submitted to Phys. Rev. Lett.
- [8] MATSUOKA, K., et al., Proc. 12th IAEA Conf. on Plasma Phys. and Controlled Nucl. Fusion Research, Nice, 1988, Vol.2, IAEA, Vienna (1989)411.
- [9] DARROW, D., ISOBE, M., et al., J. Plasma Fusion Res. SERIES, **1**(1998)362.
- [10] FUJISWA, A., IGUCHI, H., et al., Rev. Sci. Instrum. **63** (1992 ) 3694.
- [11] SAKAKIBARA, S., et al., Jpn. J. Appl. Phys. **63**(1994 )4406.
- [12] KONDO, T., ISOBE, M., et al., Proc. 24th EPS conf. on Plasma Phys. and Contr. Fusion, Prague, 1998.
- [13] MATSUMOTO, T., NAKAMURA, Y., WAKATANI, M., Nucl. Fusion **36**(1996)1571.
- [14] FU, G.Y., et al., Phys. Rev. Lett. **75**(1995)2336.
- [15] BIGLARI, H., ZONCA, F., CHEN, L., Phys. Fluids. **B4**(1992)2385.



**Fig.1** (a) Time evolution of magnetic fluctuations of  $m=3/n=2$  FBs and energetic ion loss flux in the outward-shifted plasma of  $R_{ax}=0.97$  m, where  $B_t=0.9$  T, line averaged electron density  $\sim 0.8 \times 10^{19} \text{ m}^{-3}$  and peak plasma current  $\sim 6$  kA. (b) expanded time trace of Fig.(a).




Fig.2(a)




Fig.3




Fig.2(b)




Fig.4

Fig.2 (a) Time evolution of magnetic fluctuations of  $m=2/n=1$  FBs in the inward-shifted plasma of  $R_{ax}=0.92$  m at  $B_t=0.9$  T. (b) sawtooth crashes induced by FBs are observed in the SX-signals.

Fig.3 Detailed behaviours of magnetic fluctuations of  $m=3/n=2$  FBs and temporal frequency . Fig.4 Radial profiles of soft X-ray fluctuations during two bursts of  $m=2/n=1$  FBs. Shaded zones indicate the predicted rational surfaces of  $q=2/1$ .




Fig.5





Fig.6(a) and (b)

Fig.5 Comparison of observed frequencies of TAEs with the calculated TAE frequencies.

Fig.6 (a) Alfvén continua calculated in the cylindrical configuration for  $n=1$  TAE observed in CHS, and TAE gap structure calculated from the approximate expression in large-aspect-ratio and low beta tokamak. The horizontal line indicates the observed frequency, (b) Radial variation of coherence between SX signals and magnetic fluctuations of  $n=1$  TAE, where the high coherence region around the center is due to the pass-integral effect in SX-signals.

## Recent Issues of NIFS Series

- NIFS-510 C.C. Hegna and N. Nakajima,  
*On the Stability of Mercier and Ballooning Modes in Stellarator Configurations*, Oct 1997
- NIFS-511 K. Ono and T. Hatori,  
*Rotation and Oscillation of Nonlinear Dipole Vortex in the Drift-Unstable Plasma*, Oct 1997
- NIFS-512 J. Uramoto,  
*Clear Detection of Negative Muonlike Particles from H<sub>2</sub> Gas Discharge in Magnetic Field*, Oct 1997
- NIFS-513 T. Shimozuma, M. Sato, Y. Takita, S. Ito, S. Kubo, H. Idei, K. Ohkubo, T. Watan, T.S. Chu, K. Felch, P. Cahalan and C.M. Lonng, Jr.,  
*The First Preliminary Experiments on an 84 GHz Gyrotron with a Single-Stage Depressed Collector*, Oct 1997
- NIFS-514 T. Shimozuma, S. Morimoto, M. Sato, Y. Takita, S. Ito, S. Kubo, H. Idei, K. Ohkubo and T. Watan,  
*A Forced Gas-Cooled Single-Disk Window Using Silicon Nitride Composite for High Power CW Millimeter Waves*, Oct 1997
- NIFS-515 K. Akaishi,  
*On the Solution of the Outgassing Equation for the Pump-down of an Unbaked Vacuum System*, Oct 1997
- NIFS-516 *Papers Presented at the 6th H-mode Workshop (Seeon, Germany)*, Oct 1997
- NIFS-517 John L. Johnson,  
*The Quest for Fusion Energy*, Oct 1997
- NIFS-518 J. Chen, N. Nakajima and M. Okamoto,  
*Shift-and-Inverse Lanczos Algorithm for Ideal MHD Stability Analysis*, Nov 1997
- NIFS-519 M. Yokoyama, N. Nakajima and M. Okamoto,  
*Nonlinear Incompressible Poloidal Viscosity in L=2 Heliotron and Quasi-Symmetric Stellarators*, Nov 1997
- NIFS-520 S. Kida and H. Miura,  
*Identification and Analysis of Vortical Structures*, Nov. 1997
- NIFS-521 K. Ida, S. Nishimura, T. Minami, K. Tanaka, S. Okamura, M. Osakabe, H. Idei, S. Kubo, C. Takahashi and K. Matsuoka,  
*High Ion Temperature Mode in CHS Heliotron/torsatron Plasmas*, Nov 1997
- NIFS-522 M. Yokoyama, N. Nakajima and M. Okamoto,  
*Realization and Classification of Symmetric Stellarator Configurations through Plasma Boundary Modulations*, Dec. 1997
- NIFS-523 H. Kitauchi,  
*Topological Structure of Magnetic Flux Lines Generated by Thermal Convection in a Rotating Spherical Shell*, Dec. 1997
- NIFS-524 T. Ohkawa,  
*Tunneling Electron Trap*, Dec 1997
- NIFS-525 K. Itoh, S.-I. Itoh, M. Yagi, A. Fukuyama,  
*Solitary Radial Electric Field Structure in Tokamak Plasmas*, Dec 1997
- NIFS-526 Andrey N. Lyakhov,  
*Alfven Instabilities in FRC Plasma*, Dec 1997
- NIFS-527 J. Uramoto,  
*Net Current Increment of negative Muonlike Particle Produced by the Electron and Positive Ion Bunch-method*, Dec. 1997
- NIFS-528 Andrey N. Lyakhov,  
*Comments on Electrostatic Drift Instabilities in Field Reversed Configuration*, Dec. 1997

- NIFS-529 J. Uramoto,  
*Pair Creation of Negative and Positive Pionlike (Muonlike) Particle by Interaction between an Electron Bunch and a Positive Ion Bunch*; Dec 1997
- NIFS-530 J. Uramoto,  
*Measuring Method of Decay Time of Negative Muonlike Particle by Beam Collector Applied RF Bias Voltage*; Dec. 1997
- NIFS-531 J. Uramoto,  
*Confirmation Method for Metal Plate Penetration of Low Energy Negative Pionlike or Muonlike Particle Beam under Positive Ions*. Dec. 1997
- NIFS-532 J. Uramoto,  
*Pair Creations of Negative and Positive Pionlike (Muonlike) Particle or K Mesonlike (Muonlike) Particle in H<sub>2</sub> or D<sub>2</sub> Gas Discharge in Magnetic Field*; Dec 1997
- NIFS-533 S. Kawata, C. Boonmee, T. Teramoto, L. Drska, J. Limpouch, R. Liska, M. Sinor,  
*Computer-Assisted Particle-in-Cell Code Development*; Dec. 1997
- NIFS-534 Y. Matsukawa, T. Suda, S. Ohnuki and C. Namba,  
*Microstructure and Mechanical Property of Neutron Irradiated TiNi Shape Memory Alloy*; Jan 1998
- NIFS-535 A. Fujisawa, H. Iguchi, H. Idei, S. Kubo, K. Matsuoka, S. Okamura, K. Tanaka, T. Minami, S. Ohdachi, S. Morita, H. Zushi, S. Lee, M. Osakabe, R. Akiyama, Y. Yoshimura, K. Toi, H. Sanuki, K. Itoh, A. Shimizu, S. Takagi, A. Ejiri, C. Takahashi, M. Kojima, S. Hidekuma, K. Ida, S. Nishimura, N. Inoue, R. Sakamoto, S.-I. Itoh, Y. Hamada, M. Fujiwara,  
*Discovery of Electric Pulsation in a Toroidal Helical Plasma*. Jan. 1998
- NIFS-536 Lj R. Hadzievski, M.M. Skoric, M. Kono and T. Sato,  
*Simulation of Weak and Strong Langmuir Collapse Regimes*; Jan. 1998
- NIFS-537 H. Sugama, W. Horton,  
*Nonlinear Electromagnetic Gyrokinetic Equation for Plasmas with Large Mean Flows*, Feb. 1998
- NIFS-538 H. Iguchi, T.P. Crowley, A. Fujisawa, S. Lee, K. Tanaka, T. Minami, S. Nishimura, K. Ida, R. Akiyama, Y. Hamada, H. Idei, M. Isobe, M. Kojima, S. Kubo, S. Morita, S. Ohdachi, S. Okamura, M. Osakabe, K. Matsuoka, C. Takahashi and K. Toi,  
*Space Potential Fluctuations during MHD Activities in the Compact Helical System (CHS)*; Feb. 1998
- NIFS-539 Takashi Yabe and Yan Zhang,  
*Effect of Ambient Gas on Three-Dimensional Breakup in Coronet Formation Process*, Feb. 1998
- NIFS-540 H. Nakamura, K. Ikeda and S. Yamaguchi,  
*Transport Coefficients of InSb in a Strong Magnetic Field*; Feb. 1998
- NIFS-541 J. Uramoto,  
*Development of  $\nu_{\mu}$  Beam Detector and Large Area  $\nu_{\mu}$  Beam Source by H<sub>2</sub> Gas Discharge (I)*; Mar. 1998
- NIFS-542 J. Uramoto,  
*Development of  $\bar{\nu}_{\mu}$  Beam Detector and Large Area  $\bar{\nu}_{\mu}$  Beam Source by H<sub>2</sub> Gas Discharge (II)*, Mar. 1998
- NIFS-543 J. Uramoto,  
*Some Problems inside a Mass Analyzer for Pions Extracted from a H<sub>2</sub> Gas Discharge*; Mar 1998
- NIFS-544 J. Uramoto,  
*Simplified  $\nu_{\mu}$   $\bar{\nu}_{\mu}$  Beam Detector and  $\nu_{\mu}$   $\bar{\nu}_{\mu}$  Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch*; Mar. 1998
- NIFS-545 J. Uramoto,  
*Various Neutrino Beams Generated by D<sub>2</sub> Gas Discharge*; Mar.1998
- NIFS-546 R. Kanno, N. Nakajima, T. Hayashi and M. Okamoto,  
*Computational Study of Three Dimensional Equilibria with the Bootstrap Current*; Mar. 1998



- NIFS-547 R Kanno, N Nakajima and M Okamoto,  
*Electron Heat Transport in a Self-Similar Structure of Magnetic Islands*, Apr 1998
- NIFS-548 J.E Rice,  
*Simulated Impurity Transport in LHD from MIST*; May 1998
- NIFS-549 M.M. Skonc, T Sato, A.M Maluckov and M S Jovanovic,  
*On Kinetic Complexity in a Three-Wave Interaction*, June 1998
- NIFS-550 S Goto and S Kida,  
*Passive Saclar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation*; June 1998
- NIFS-551 T Kuroda, H. Sugama, R Kanno, M Okamoto and W Horton,  
*Initial Value Problem of the Toroidal Ion Temperature Gradient Mode* : June 1998
- NIFS-552 T Mutoh, R Kumazawa, T Seki, F Simpo, G Nomura, T Ido and T Watan,  
*Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device* , June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T Muroga,  
*oronaization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-* July 1998
- NIFS-554 S Murakami and H Saleem,  
*Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations*, July 1998
- NIFS-555 H Nakamura , K. Ikeda and S Yamaguchi,  
*Physical Model of Nernst Element*, Aug 1998
- NIFS-556 H Okumura, S Yamaguchi, H. Nakamura, K Ikeda and K Sawada,  
*Numerical Computation of Thermoelectric and Thermomagnetic Effects*, Aug 1998
- NIFS-557 Y Takeiri, M. Osakabe, K. Tsumori, Y Oka, O Kaneko, E. Asano, T Kawamoto, R. Akiyama and M. Tanaka,  
*Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System* , Aug.1998
- NIFS-558 M. Tanaka, A Yu Grosberg and T Tanaka,  
*Molecular Dynamics of Structure Organization of Polyampholytes*, Sep 1998
- NIFS-559 R Honuchi, K. Nishimura and T. Watanabe,  
*Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations*, Sep 1998  
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Kholopenkov, K Matsuoka, S Okamura, C Takahashi, R Akiyama, A. Fujisawa, K Ida, H Idei, H Iguchi, M Isobe, S Kado, K Kondo, S Kubo, H. Kuramoto, T. Minami, S Morita, S. Nishimura, M. Osakabe, M Sasao, B. Peterson, K Tanaka, K. Toi and Y. Yoshimura,  
*Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection*, Oct 1998  
(IAEA-CN-69/EXP1/18)
- NIFS-561 A Fujisawa, H. Iguchi, S Lee, K Tanaka, T Minami, Y. Yoshimura, M Osakabe, K. Matsuoka, S Okamura, H Idei, S Kubo, S Ohdachi, S. Morita, R Akiyama, K Toi, H. Sanuki, K Itoh, K Ida, A. Shimizu, S Takagi, C Takahashi, M Kojima, S Hidekuma, S Nishimura, M. Isobe, A Ejri, N. Inoue, R. Sakamoto, Y. Hamada and M Fujiwara,  
*Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron*, Oct 1998  
(IAEA-CN-69/EX5/1)
- NIFS-562 S Yoshikawa,  
*Next Generation Toroidal Devices*; Oct 1998
- NIFS-563 Y. Todo and T Sato,  
*Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes*; Oct 1998  
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watan, T Shimozuma, Y Takeiri, R Kumazawa, T Mutoh, M Sato, O Kaneko, K Ohkubo, S Kubo, H Idei, Y Oka, M Osakabe, T Seki, K Tsumori, Y. Yoshimura, R Akiyama, T Kawamoto, S Kobayashi, F Shimpo, Y Takita, E Asano, S Itoh, G Nomura, T Ido, M. Hamabe, M. Fujiwara, A Iiyoshi, S Morimoto, T Bigelow and Y.P Zhao,  
*Steady State Heating Technology Development for LHD* : Oct 1998  
(IAEA-CN-69/FTP/21)

- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyaibu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda, *LHD-Type Compact Helical Reactors*, Oct 1998 (IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto, *Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems*; Oct 1998 (IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D.S. Darrow, H. Idei, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tsumon, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watari, *Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas*; Oct. 1998 (IAEA-CN-69/EX2/2)
- NIFS-568 S. Okamura, K. Matsuoka, R. Akiyama, D.S. Darrow, A. Ejiri, A. Fujisawa, M. Fujiwara, M. Goto, K. Ida, H. Idei, H. Iguchi, N. Inoue, M. Isobe, K. Itoh, S. Kado, K. Khlopenkov, T. Kondo, S. Kubo, A. Lazaros, S. Lee, G. Matsunaga, T. Minami, S. Morita, S. Murakami, N. Nakajima, N. Nikai, S. Nishimura, I. Nomura, S. Ohdachi, K. Ohkuni, M. Osakabe, R. Pavlichenko, B. Peterson, R. Sakamoto, H. Sanuki, M. Sasao, A. Shimizu, Y. Shirai, S. Sudo, S. Takagi, C. Takahashi, S. Takayama, M. Takechi, K. Tanaka, K. Toi, K. Yamazaki, Y. Yoshimura and T. Watari, *Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS*, Oct. 1998 (IAEA-CN-69/OV4/5)
- NIFS-569 M.M. Skoric, T. Sato, A. Maluckov, M.S. Jovanovic, *Micro- and Macro-scale Self-organization in a Dissipative Plasma*, Oct 1998
- NIFS-570 T. Hayashi, N. Mizuguchi, T-H. Watanabe, T. Sato and the Complexity Simulation Group, *Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak*, Oct 1998 (IAEA-CN-69/TH3/3)
- NIFS-571 A. Iyoshi, A. Komori, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Monsaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Nanbara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyaibu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumon, I. Yamada, H. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, S. Ima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, M. Fujiwara and O. Motojima, *An Overview of the Large Helical Device Project*; Oct 1998 (IAEA-CN-69/OV1/4)
- NIFS-572 M. Fujiwara, H. Yamada, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Monsaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Nanbara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyaibu, Y. Oka, M. Osakabe, T. Ozaki, B. J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumon, I. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, M. Ima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iyoshi, *Plasma Confinement Studies in LHD*; Oct. 1998 (IAEA-CN-69/EX2/3)
- NIFS-573 O. Motojima, K. Akaishi, H. Chikaraishi, H. Funaba, S. Hamaguchi, S. Imagawa, S. Inagaki, N. Inoue, A. Iwamoto, S. Kitagawa, A. Komori, Y. Kubota, R. Maekawa, S. Masuzaki, T. Mito, J. Miyazawa, T. Monsaki, T. Muroga, T. Nagasaka, Y. Nakamura, A. Nishimura, K. Nishimura, N. Noda, N. Ohyaibu, S. Sagara, S. Sakakibara, R. Sakamoto, S. Sato, T. Satow, M. Shoji, H. Suzuki, K. Takahata, H. Tamura, K. Watanabe, H. Yamada, S. Yamada, S. Yamaguchi, K. Yamazaki, N. Yanagi, T. Baba, H. Hayashi, M. Ima, T. Inoue, S. Kato, T. Kato, T. Kondo, S. Moriuchi, H. Ogawa, I. Ohtake, K. Ooba, H. Sekiguchi, N. Suzuki, S. Takami, Y. Taniguchi, T. Tsuzuki, N. Yamamoto, K. Yasui, H. Yonezu, M. Fujiwara and A. Iyoshi, *Progress Summary of LHD Engineering Design and Construction*; Oct. 1998 (IAEA-CN-69/FT2/1)
- NIFS-574 K. Toi, M. Takechi, S. Takagi, G. Matsunaga, M. Isobe, T. Kondo, M. Sasao, D.S. Darrow, K. Ohkuni, S. Ohdachi, R. Akiyama, A. Fujisawa, M. Goto, H. Idei, K. Ida, H. Iguchi, S. Kado, M. Kojima, S. Kubo, S. Lee, K. Matsuoka, T. Minami, S. Morita, N. Nikai, S. Nishimura, S. Okamura, M. Osakabe, A. Shimizu, Y. Shirai, C. Takahashi, K. Tanaka, T. Watari and Y. Yoshimura, *Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas*; Oct. 1998 (IAEA-CN-69/EXP1/19)