

# NATIONAL INSTITUTE FOR FUSION SCIENCE

## Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma

H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida,  
S. Toda, S.-I. Itoh, M. Yagi and A. Fukuyama

(Received - May 21, 1999 )

NIFS-597

May 1999

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

**RESEARCH REPORT**  
**NIFS Series**

# Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma

H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida, S. Toda,  
S.-I. Itoh<sup>†</sup>, M. Yagi<sup>†</sup>, A. Fukuyama\*

National Institute for Fusion Science, Toki, Gifu, 509-5292

<sup>†</sup>Research Institute for Applied Mechanics, Kyushu University, Kasuga 816-8580

\*Department of Nuclear Engineering, Kyoto University, Kyoto, 606-8501

## Abstract

Theoretical analysis of the electric field bifurcation is made for the LHD plasma. For given shapes of plasma profiles, a region of bifurcation is obtained in a space of the plasma parameters. In this region of plasma parameters, the electric field domain interface is predicted to appear in the plasma column. The reduction of turbulent transport is expected to occur in the vicinity of the interface, inducing an internal transport barrier. Within this simple model, the plasma with internal barriers is predicted to be realized for the parameters of  $T_e(0) \sim 2\text{keV}$  and  $n(0) \sim 10^{18}\text{m}^{-3}$ .

Keywords: Internal Transport Barrier, Electric Field Bifurcation, LHD plasma

## 1. Introduction

In the last two decades, various types of improved confinement states have been identified in tokamaks [1]. Among the variety of improved confinement modes, the H-mode [2] has been studied most thoroughly. The electric field bifurcation model [3] has been proposed to explain the physics of the H-mode, and the role of radial electric field structure on various improved confinement modes has been widely recognized [4, 5].

In high temperature plasmas which are confined in CHS and LHD, anomalous energy transport dominates over neoclassical energy

transport. The improved energy confinement, if it exists, could be realized when the anomalous transport is suppressed. Based on the turbulent transport model, the improved confinement has been analyzed in a geometry of Heliotron configuration [6]. A possibility of existence of internal transport barrier has been theoretically predicted; the electric field domain interface could be established, and transport reduction at the interface takes place [7]. The electric field domain interface has been identified in CHS plasma [8], and the internal transport barrier has been found in the CHS plasma [9]. These developments of confinement theory and experiment strongly suggest the

possibility of existence of internal transport barrier in the LHD plasma.

In this article, we first study a condition for the electric field domain interface to appear in the LHD plasma. On the parameter space of temperatures and density, the region is identified for the presence of electric field domain interface. The electric field gradient is estimated; it is strong enough to cause the reduced turbulent transport.

## 2. Model

### 2.1 Electric domain interface and internal transport barrier

For the transparency of an argument, we take the following simplification: The radial current is dominated by the neoclassical transport, while the energy transport is governed by the anomalous transport. Although progress has been made in the theory of turbulence in helical systems [10], the deductive theory has not been completed to predict the profiles of plasma density and temperature. Therefore the radial electric field is solved for a given profiles of the density and temperature. This simplification has been tested on CHS plasma, and was shown to be sufficient as a zeroth-order estimate of the radial electric field [11].

The radial electric field in the stationary state is governed by the charge neutrality equation

$$\sum_{c,j} e_j \Gamma_j = 0 \quad (1)$$

where  $e_j$  is the charge and  $\Gamma_j$  is the radial flux of the  $j$ -th particle. In this equation,  $\Gamma_j$  consists of two parts  $\Gamma_j^{\text{local}}$  and  $\Gamma_j^{\text{diffusive}}$ . The former,  $\Gamma_j^{\text{local}}$ , is expressed by the local plasma parameters alone, while the latter,  $\Gamma_j^{\text{diffusive}}$ , depends on the curvature of the radial electric field [4, 5]. As a result, Eq.(1) is a nonlinear diffusion equation. Analyses of

equation (1) have been performed. The equation of the balance of local current

$$\sum_{c,j} \Gamma_j^{\text{local}} = 0 \quad (2)$$

can have three solutions for one set of plasma parameters, two of which are stable solutions [12, 13]. More positive electric field solution is usually called 'electron root', and the other, less positive solution, is called 'ion root'. It was shown that the electric field interface appears in the plasma when Eq.(2) has multiple solutions in some region, say  $r_1 < r < r_2$  [4, 5]. There could exist a surface  $r = r_c$  ( $r_1 < r_c < r_2$ ) across which the domain of electron root and that of ion root touches each other.

When the electric field interface exist in the plasma, the transport barrier could be realized. The turbulent transport coefficient  $\chi$  for the Heliotron plasma has been derived as [6]

$$\chi = \left( 1 + \frac{\omega_E^2}{2G_0} \right)^{-1} \chi_L \quad (3)$$

where  $\omega_E = E_r \tau_{Ap} / B$  is the normalized gradient of electric field,  $\tau_{Ap}$  is the poloidal-Alfven-transit time, and  $G_0$  is the normalized pressure gradient multiplied by the magnetic field gradient.  $\chi_L$  is the transport coefficient in the absence of the electric field gradient, and the explicit formula is given in [10]. One sees that the electric field gradient  $E_r \approx 10^6 \text{Vm}^{-2}$  is the level of gradient which causes the reduction of transport for the parameters of interest. The difference of radial electric field across the domain interface is a few times of  $|\nabla T_e|$ . The layer thickness of the interface is dictated by the anomalous viscosity and collisional viscosity, and can be of the order of cm or less. The gradient

at the interface can exceed the critical level of gradient to drive the turbulence suppression [7]

Based on these theoretical analyses, it has been predicted that the internal transport barrier can exist at the electric domain interface, i.e., under the circumstance that Eq.(2) has multiple solutions. This theoretical prediction has been tested on CHS plasmas [9]. It has been confirmed by experiments that the internal transport barrier is established at the electric domain interface. Figure 1 illustrates the parameter space in which the multiple solutions of Eq.(2) exist in the plasma [7]. The region depends on the ratio of the electron and ion temperatures. The experimental region of electric domain interface is also shown by the hatched region. Plasma discharge with internal transport barrier exists in this hatched region [9]. One can conclude that the theoretical prediction for the existence of the internal transport barrier has been semi-quantitatively confirmed.

## 2.2 Analytic Model

The analysis of [7, 11] is extended to the LHD plasma. The objective of the present study is to provide an analytic insight, i.e., the level of comparison with CHS plasmas, so that a simplified theoretical model is used. The model Heliotron/torsatron configuration is employed, in which the magnetic field is characterized by the helical ripple and toroidal ripple as

$$\mathbf{B} = B_0 \left( 1 - \varepsilon_t \cos \theta - \varepsilon_h \cos (m\theta - \ell\varphi) \right). \quad (4)$$

In this expression, quasi-toroidal coordinates  $(r, \theta, \varphi)$  are used ( $r$  is the minor radius,  $\theta$  is the poloidal angle,  $\varphi$  is the toroidal angle),  $\varepsilon_t = r/R$  is the toroidal ripple,  $\varepsilon_h$  is the helical ripple,  $\ell$  is the polarity of the field, and  $m$  is the toroidal pitch

number. Limit of high aspect ratio,  $R/a \gg 1$ , is employed ( $R$  and  $a$  being the major and minor radius, respectively). The radial profile of the ripple is approximated as  $\varepsilon_h(r) = 2\varepsilon_0 I_2(mr/R)$  with  $\varepsilon_0 = \sqrt{1 - (2\iota(0)/m - 1)^2}$ , where  $I_2$  is the second order modified Bessel function and  $\iota(0)$  is the central rotational transform divided by  $2\pi$ .

The local radial current due to the neoclassical diffusion is calculated by use of the formula

$$\Gamma_j^{\text{local}} = \Gamma_j^s + \Gamma_j^{\text{as}} \quad (5)$$

where  $\Gamma_j^s$  is the particle flux associated with transit or toroidally trapped particles, corresponding to the one in axisymmetric systems, and  $\Gamma_j^{\text{as}}$  indicates the flux due to helically trapped particles. The explicit expressions for  $\Gamma_j^s$  and  $\Gamma_j^{\text{as}}$  are Eqs.(6)-(13) of Ref.[12]. This formula is simple, but was useful in understanding the Heliotron experiments.

## 3. Parameter Region of Domain Interface

Equation (2) with Eq.(5) is solved for a given plasma profiles. We choose the profiles as parabolic,

$$n/n(0) = T_e/T_e(0) = T_i/T_i(0) = 1 - r^2/a^2 \quad (6)$$

Within this choice, the plasma profiles are characterized by a set of parameters  $n(0), T_e(0), T_i(0)$ . Other parameters are chosen as  $R = 3.75$  m,  $a = 0.576$  m, and  $B_0 = 3.0$  T.

The region of multiple solutions for Eq.(2) is searched for in the parameter space of  $(n(0), T_e(0), T_i(0))$ . Figure 2 illustrates regions of radial electric field structures. There are three characteristic regions in this diagram; the regions of electron root, multiple solutions, and ion root.

The electron root dominates over the whole plasma column, if the temperature is high enough. In the high density region, the ion root dominates. In between these two regions, the region of multiple solutions exists. The region of multiple solution is shown on the  $(n(0), T_e(0))$  plane for various values of the ratio of electron temperature to ion temperature. For the ratio of  $T_e(0)/T_i(0) = 1$ , the multiple solutions are possible in the temperature region

$$1\text{keV} < T_e(0) < 2\text{keV} \quad (7)$$

at the low plasma density  $n_e(0) \approx 10^{18}\text{m}^{-3}$ . The condition for the interface to exist in the LHD plasma is that the plasma parameter lies in the hatched region, which is bounded like Eq.(7). When the temperature is in this region and high (for fixed density), the interface exists near the plasma boundary. When the temperature is lower, the interface appears near the center. For the fixed temperature, the interface approaches to the edge when the density is lower.

As the density increases, the region of interests moves to the higher temperature region. In this region, the electric domain interface is predicted to appear. On the boundary between these regions, the bifurcation occurs from one radial structure to the other.

The appearance of the multiple solution, i.e., the electric domain interface, is influenced by the ion temperature. Figure 3 shows the electric field structure on the  $(T_e(0), T_i(0))$  plane for the fixed density,  $n(0) = 5 \times 10^{18}\text{m}^{-3}$ . The lower boundary of the region of our interests weakly depends on the ion temperature. An empirical fit of the lower boundary of the electron temperature for transition,  $T_{e,c}(0)$ , is observed as

$$T_{e,c\{[\text{keV}]\}}(0) \sim 2 T_{i\{[\text{keV}]\}}(0)^{1/4} \quad (8)$$

for this density. On the other hand, the upper boundary is modified by ion temperature more strongly. In the low ion temperature limit,  $T_i(0)/T_e(0) < 1/5$ , or in the high ion temperature limit,  $T_i(0)/T_e(0) > 7/3$ , the upper boundary merges to the lower boundary: When two boundaries merge, the whole plasma column is dominated by either the electron root or ion root solutions, so that the electric domain interface is difficult to appear.

From these analyses, one can conclude that, when the electron temperature reaches as  $T_e(0) \approx 2\text{keV}$ , the electric domain interface is sustained in the plasma column. If the domain interface appears in the plasma column, the gradient of the electric field at the domain is possible to reach the level of  $E_r' \sim 10^6\text{Vm}^{-2}$ . Therefore, the internal transport barrier is predicted to appear under such circumstance, based on the study of the turbulent transport and comparison with CHS experiments.

#### 4. Summary and discussion

In this article, we studied the region of plasma parameters for which the electric field domain interface appears in the LHD plasma. If the domain interface exists, the gradient of the field at this interface is so strong as to cause the reduction of turbulent transport. Based on this theoretical hypothesis, which has been successfully tested on CHS plasma, the possibility to sustain the internal transport barrier is examined. By employing a model of profiles, the parameter survey is performed. The condition for the possible existence of internal transport barrier is identified.

Within this simple model, the plasma with electric domain interface could be realized for the parameters of  $T_e(0) \sim 2\text{keV}$  and  $n(0) = 10^{18}\text{m}^{-3}$ . This parameter range is not far from the experimentally-accessible parameters. More quantitative analysis must be performed by employing profiles which are observed in experiments. The objective of this article is to point out the possible existence of the internal transport barrier; more quantitative estimate shall be reported in future.

Before closing this article, a role of energetic electron is discussed. In the presence of intense electron heating by ECH, considerable amount of energetic electrons is confined in plasmas. The radial flux of such energetic electrons, which is not included in Eq.(2), could influence the radial electric field [14, 15]. There are two ways in influencing the electric field. One is the quantitative modification of the transition condition. An additional loss of energetic electrons modifies the boundaries in Fig.2 to a higher-density region. This is the quantitative shift, and it does not change the conclusion of this article. The second way of the influence of the energetic electrons is a possibility to induce a strong positive electric field in the edge region. The analysis has been developed in [15, 16], and recent qualitative computation is seen in [17]. This induced electron root was observed in the higher density region [18] in the CHS plasma. The improved confinement associated with the electric field interface has been observed experimentally under the circumstance that the stronger radial electric field is observed in the central region of the plasma. The electric interface, which is sustained without the contribution of energetic electrons, has been verified to provide an improved confinement.

### Acknowledgements

This work is partly supported by the Grant-in-Aid for Scientific Research and the Grant-in-Aid for International Scientific Research of Ministry of Education, Science, Sports and Culture of Japan. Authors wish to thank the collaboration programme of National Institute for Fusion Science and the collaboration programme of Advanced Fusion Research Center, RIAM of Kyushu University.

## References

- [1] A review of improved confinement is given in S.-I. Itoh, K. Itoh, A. Fukuyama; *Journal of Nuclear Materials* **220-222** (1995), pp.117-131,
- [2] Wagner F et al 1982 *Phys. Rev. Lett.* **49** 1408.
- [3] Itoh S-I, Itoh K 1988 *Phys. Rev. Lett.* **60** 2276.
- [4] S. -I. Itoh, et al., *Phys. Rev. Lett.* **67**, 2485 (1991). See a review, e.g., K. Itoh, S.-I. Itoh, *Plasma Phys. Contr. Fusion* **38** (1996) 1.
- [5] Itoh K, Itoh S-I, Fukuyama A; *Transport and Structural Formation in Plasmas* (IoP, Bristol, England, 1999).
- [6] K. Itoh, S.-I. Itoh, A. Fukuyama, H. Sanuki and M. Yagi.; *Plasma Physics and Controlled Fusion* **36** (1994), pp.123-129.
- [7] K. Itoh, et al.; *J. Plasma Fusion research Suppl.* **74** (1998) 282. [In Japanese]
- [8] A. Fujisawa, H. Iguchi, H. Sanuki, K. Itoh, et al.; *Phys. Rev. Lett.* **79** (1997) 1054-1057
- [9] A. Fujisawa, H. Iguchi, T. Minami, Y. Yoshimura, H. Sanuki, K. Itoh, et al.; *Phys. Rev. Lett.* **82** (1999) 2669
- [10] Itoh K, Itoh S-I, Fukuyama A, Yagi M, Azumi M; *Plasma Phys. Cont. Fusion* **36** (1994) 1501.
- [11] H. Sanuki, K. Itoh, K. Ida and S.-I. Itoh; *J. Phys. Soc. Japan* **60** (1991), pp.3698-3705. See also H. Sanuki; in *Frontier of Physics in Fusion-Relevant Plasmas* (ed., M. Fujiwara, Y. X. Wan, World Scientific, Singapore 1998) 199.
- [12] L. M. Kovrizhnykh; *Nucl. Fusion* **24** (1984) 435.
- [13] D. E. Hastings and T. Kamimura; *J. Computational Physics* **61** (1985) 286.
- [14] H. Sanuki, K. Itoh and S.-I. Itoh; *J. Phys. Soc. Jpn.* **62** (1993), pp.123-139.
- [15] H. Sanuki, K. Itoh, J. Todoroki, et al.; *Physica Scripta* **52** (1995) 461. See also H. Sanuki, K. Itoh, M. Yokoyama, et al; to be published in *Chinese Physics Letters* (1999).
- [16] H. Idei, K. Ida, H. Sanuki, et al.; *Phys. Plasmas* **1** (1994) 3400.
- [17] S. Murakami, N. Nakajima, U. Gasparino, M. Okamoto; *Plasma Physics Report* **23** (1997) 556.
- [18] H. Idei, K. Ida, H. Sanuki et al., *Phys. Rev. Lett.* **71** 2220 (1993).

## Figure Captions

**Fig.1** Boundary in the plasma parameter space for the positive radial electric field and negative one. Hatched region indicates the possibility of multiple solutions. Computations are made for the case of CHS plasma with  $B = 1T$ . Bifurcation is experimentally observed in the area which is denoted by a shaded ellipse. This comparison suggests that the present theoretical method provides a rough and qualitative explanation of radial electric field.

**Fig.2** Regions of the multiple electric solutions. electron root, and ion root are shown on the  $(n(0), T_e(0))$  plane. In the shaded area, the multiple solution is possible and the electric domain interface could be sustained in the plasma

column The cases of  $T_e(0)/T_i(0) = 1$  (a) and  $T_e(0)/T_i(0) = 3$  (b) are shown.

**Fig. 3** Regions of the multiple electric solutions, electron root, and ion root are shown on the  $(T_e(0), T_i(0))$  plane for the parameter of  $n(0) = 5 \times 10^{18} \text{m}^{-3}$ . In the shaded area, the multiple solution is possible and the electric domain interface could be sustained in the plasma column.



Fig.1




Fig.2






Fig.3



## Recent Issues of NIFS Series

- NIFS-533 S Kawata, C Boonmee, T Teramoto, L Drska, J Limpouch, R Liska, M Sinor.  
*Computer-Assisted Particle-in-Cell Code Development* Dec 1997
- NIFS-534 Y Matsukawa, T Suda, S Ohnuki and C Namba.  
*Microstructure and Mechanical Property of Neutron Irradiated TiNi Shape Memory Alloy* Jan 1998
- NIFS-535 A Fujisawa, H Iguchi, H Idei, S Kubo, K. Matsuoka, S Okamura, K Tanaka, T Minami, S Ohdachi, S Morita, H. Zushi, S Lee, M Osakabe, R Akiyama, Y Yoshimura, K Toi, H Sanuki, K Itoh, A Shimizu, S Takagi, A Ejiri, C Takahashi, M Kojima, S Hidekuma, K Ida, S Nishimura, N Inoue, R Sakamoto, S-I Itoh, Y Hamada, M Fujiwara.  
*Discovery of Electric Pulsation in a Toroidal Helical Plasma* Jan 1998
- NIFS-536 Lj R Hadzievski, M M Skoric, M Kono and T Sato.  
*Simulation of Weak and Strong Langmuir Collapse Regimes* Jan 1998
- NIFS-537 H Sugama, W Horton  
*Nonlinear Electromagnetic Gyrokinetic Equation for Plasmas with Large Mean Flows* Feb 1998
- NIFS-538 H. Iguchi, T.P Crowley, A Fujisawa, S Lee, K Tanaka, T Minami, S Nishimura, K. Ida, R Akiyama, Y Hamada, H. Idei, M Isobe, M. Kojima, S Kubo, S Morita, S Ohdachi, S Okamura, M Osakabe, K Matsuoka, C Takahashi and K Toi.  
*Space Potential Fluctuations during MHD Activities in the Compact Helical System (CHS)*: Feb 1998
- NIFS-539 Takashi Yabe and Yan Zhang.  
*Effect of Ambient Gas on Three-Dimensional Breakup in Coronet Formation Process*. Feb 1998
- NIFS-540 H Nakamura, K Ikeda and S Yamaguchi  
*Transport Coefficients of InSb in a Strong Magnetic Field*. Feb 1998
- NIFS-541 J Uramoto,  
*Development of  $v_{\mu}$  Beam Detector and Large Area  $v_{\mu}$  Beam Source by  $H_2$  Gas Discharge (I)*; Mar 1998
- NIFS-542 J. Uramoto,  
*Development of  $\bar{v}_{\mu}$  Beam Detector and Large Area  $\bar{v}_{\mu}$  Beam Source by  $H_2$  Gas Discharge (II)*.  
Mar. 1998
- NIFS-543 J Uramoto,  
*Some Problems inside a Mass Analyzer for Pions Extracted from a  $H_2$  Gas Discharge*. Mar 1998
- NIFS-544 J Uramoto,  
*Simplified  $v_{\mu}$ ,  $\bar{v}_{\mu}$  Beam Detector and  $v_{\mu}$ ,  $\bar{v}_{\mu}$  Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch*. Mar. 1998
- NIFS-545 J. Uramoto,  
*Various Neutrino Beams Generated by  $D_2$  Gas Discharge*, Mar.1998
- NIFS-546 R Kanno, N. Nakajima, T Hayashi and M. Okamoto,  
*Computational Study of Three Dimensional Equilibria with the Bootstrap Current*. Mar. 1998
- NIFS-547 R. Kanno, N. Nakajima and M. Okamoto.  
*Electron Heat Transport in a Self-Similar Structure of Magnetic Islands*: Apr 1998
- NIFS-548 J.E Rice,  
*Simulated Impurity Transport in LHD from MIST*: May 1998
- NIFS-549 M.M. Skoric, T Sato, A M Maluckov and M S. Jovanovic.  
*On Kinetic Complexity in a Three-Wave Interaction*: June 1998
- NIFS-550 S Goto and S Kida.  
*Passive Saclar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation*: June 1998

- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,  
*Initial Value Problem of the Toroidal Ion Temperature Gradient Mode* ; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watari,  
*Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device* ; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,  
*Ionization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-* July 1998
- NIFS-554 S. Murakami and H. Saleem,  
*Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations*; July 1998
- NIFS-555 H. Nakamura, K. Ikeda and S. Yamaguchi,  
*Physical Model of Nernst Element*, Aug 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,  
*Numerical Computation of Thermoelectric and Thermomagnetic Effects*; Aug 1998
- NIFS-557 Y. Takeiri, M. Osakabe, K. Tsumori, Y. Oka, O. Kaneko, E. Asano, T. Kawamoto, R. Akiyama and M. Tanaka,  
*Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System*; Aug. 1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,  
*Molecular Dynamics of Structure Organization of Polyampholytes*; Sep 1998
- NIFS-559 R. Horiuchi, K. Nishimura and T. Watanabe,  
*Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations*, Sep 1998  
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Kholopenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,  
*Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection*; Oct. 1998  
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,  
*Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron*; Oct 1998  
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,  
*Next Generation Toroidal Devices*; Oct. 1998
- NIFS-563 Y. Todo and T. Sato,  
*Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes*; Oct 1998  
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watari, T. Shimozuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Idei, Y. Oka, M. Osakabe, T. Seki, K. Tsumori, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Shimo, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Morimoto, T. Bigelow and Y. P. Zhao,  
*Steady State Heating Technology Development for LHD*; Oct. 1998  
(IAEA-CN-69/FTP/21)
- NIFS-565 A. Sagara, K. Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyaibu, T. Satow, A. A. Shishkin, S. Tanaka, T. Terai and T. Uda,  
*LHD-Type Compact Helical Reactors*. Oct. 1998  
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,  
*Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems*, Oct. 1998  
(IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D. S. Darrow, H. Idei, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tsumori, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watari,  
*Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas*; Oct. 1998  
(IAEA-CN-69/EX2/2)

- NIFS-568 S Okamura, K Matsuoka, R Akiyama, D S Darrow, A Ejiri, A Fujisawa, M Fujiwara, M Goto, K Ida, H Idei, H Iguchi, N Inoue, M Isobe, K Itoh, S Kado, K Khipopenkov, T Kondo, S Kubo, A Lazaros, S Lee, G Matsunaga, T Minami, S Morita, S Murakami, N Nakajima, N Nikai, S Nishimura, I Nomura, S Ohdachi, K Ohkuni, M Osakabe, R Pavlichenko, B Peterson, R Sakamoto, H Sanuki, M Sasao, A Shimizu, Y Shirai, S Sudo, S Takagi, C Takahashi, S Takayama, M Takechi, K Tanaka, K Toi, K Yamazaki, Y Yoshimura and T Watari  
*Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS*, Oct 1998  
(IAEA-CN-69/OV4/5)
- NIFS-569 M.M Skoric, T Sato, A Maluckov, M S Jovanovic,  
*Micro- and Macro-scale Self-organization in a Dissipative Plasma* Oct 1998
- NIFS-570 T Hayashi, N Mizuguchi, T-H Watanabe, T Sato and the Complexity Simulation Group,  
*Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak* Oct 1998  
(IAEA-CN-69/TH3/3)
- NIFS-571 A Iiyoshi, A Komori, A Ejiri, M Emoto, H Funaba, M Goto, K Ida, H Idei, S Inagaki, S Kado, O Kaneko, K Kawahata, S Kubo, R Kumazawa, S Masuzaki, T Minami, J Miyazawa, T Morisaki, S Morita, S Murakami, S Muto, T Muto, Y Nagayama, Y Nakamura, H Nakanishi, K Narihara, K Nishimura, N Noda, T Kobuchi, S Ohdachi, N Ohyabu, Y Oka, M Osakabe, T Ozaki, B J Peterson, A Sagara, S Sakakibara, R Sakamoto, H Sasao, M Sasao, K Sato, M Sato, T Seki, T Shimozuma, M Shoji, H Suzuki, Y Takeiri, K Tanaka, K Toi, T Tokuzawa, K Tsumori, I Yamada, H Yamada, S Yamaguchi, M Yokoyama, K Y Watanabe, T Watari, R Akiyama, H Chikaraishi, K Haba, S Hamaguchi, S Iima, S Imagawa, N Inoue, K Iwamoto, S Kitagawa, Y Kubota, J Kodaira, R Maekawa, T Mito, T Nagasaka, A Nishimura, Y Takita, C Takahashi, K Takahata, K Yamauchi, H Tamura, T Tsuzuki, S Yamada, N Yanagi, H Yonezu, Y Hamada, K Matsuoka, K Murai, K Ohkubo, I Ohtake, M Okamoto, S Sato, T Satow, S Sudo, S Tanahashi, K Yamazaki, M Fujiwara and O Motojima  
*An Overview of the Large Helical Device Project* Oct. 1998  
(IAEA-CN-69/OV1/4)
- NIFS-572 M. Fujiwara, H Yamada, A Ejiri, M Emoto, H Funaba, M Goto, K Ida, H Idei, S Inagaki, S Kado, O Kaneko, K Kawahata, A Komori, S Kubo, R Kumazawa, S Masuzaki, T Minami, J Miyazawa, T Morisaki, S Morita, S Murakami, S Muto, T Muto, Y Nagayama, Y Nakamura, H Nakanishi, K Narihara, K Nishimura, N Noda, T Kobuchi, S Ohdachi, N Ohyabu, Y Oka, M Osakabe, T Ozaki, B J Peterson, A Sagara, S Sakakibara, R Sakamoto, H Sasao, M Sasao, K Sato, M Sato, T Seki, T Shimozuma, M Shoji, H Suzuki, Y Takeiri, K Tanaka, K Toi, T Tokuzawa, K Tsumori, I Yamada, S Yamaguchi, M Yokoyama, K Y Watanabe, T Watari, R Akiyama, H Chikaraishi, K Haba, S Hamaguchi, M Iima, S Imagawa, N Inoue, K Iwamoto, S Kitagawa, Y Kubota, J Kodaira, R Maekawa, T Mito, T Nagasaka, A Nishimura, Y Takita, C Takahashi, K Takahata, K Yamauchi, H Tamura, T Tsuzuki, S Yamada, N Yanagi, H Yonezu, Y Hamada, K Matsuoka, K Murai, K Ohkubo, I Ohtake, M Okamoto, S Sato, T Satow, S Sudo, S Tanahashi, K Yamazaki, O Motojima and A Iiyoshi,  
*Plasma Confinement Studies in LHD*: Oct 1998  
(IAEA-CN-69/EX2/3)
- NIFS-573 O Motojima, K Akaishi, H Chikaraishi, H Funaba, S Hamaguchi, S Imagawa, S Inagaki, N Inoue, A Iwamoto, S Kitagawa, A Komori, Y Kubota, R Maekawa, S Masuzaki, T Mito, J Miyazawa, T Morisaki, T Muroga, T Nagasaka, Y Nakamura, A Nishimura, K Nishimura, N Noda, N Ohyabu, S Sagara, S Sakakibara, R Sakamoto, S Satoh, T Satow, M Shoji, H Suzuki, K Takahata, H Tamura, K Watanabe, H Yamada, S Yamada, S Yamaguchi, K Yamazaki, N Yanagi, T Baba, H Hayashi, M Iima, T Inoue, S Kato, T Kato, T Kondo, S Moriuchi, H Ogawa, I Ohtake, K Ooba, H Sekiguchi, N Suzuki, S Takami, Y Taniguchi, T Tsuzuki, N Yamamoto, K Yasui, H Yonezu, M Fujiwara and A Iiyoshi,  
*Progress Summary of LHD Engineering Design and Construction*: Oct 1998  
(IAEA-CN-69/FT2/1)
- NIFS-574 K Toi, M Takechi, S Takagi, G Matsunaga, M Isobe, T Kondo, M Sasao, D S Darrow, K Ohkuni, S Ohdachi, R Akiyama, A Fujisawa, M Gotoh, H Idei, K Ida, H Iguchi, S Kado, M Kojima, S Kubo, S Lee, K Matsuoka, T Minami, S Morita, N Nikai, S Nishimura, S Okamura, M Osakabe, A Shimizu, Y Shirai, C Takahashi, K Tanaka, T Watari and Y Yoshimura,  
*Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas*, Oct 1998  
(IAEA-CN-69/EXP1/19)
- NIFS-575 Y Hamada, A Nishizawa, Y Kawasumi, A Fujisawa, M Kojima, K Narihara, K Ida, A Ejiri, S Ohdachi, K Kawahata, K Toi, K Sato, T Seki, H Iguchi, K Adachi, S Hidekuma, S Hirokura, K Iwasaki, T Ido, R Kumazawa, H Kuramoto, T Minami, I Nomura, M Sasao, K N Sato, T Tsuzuki, I Yamada and T Watari,  
*Potential Turbulence in Tokamak Plasmas* Oct 1998  
(IAEA-CN-69/EXP2/14)
- NIFS-576 S Murakami, U Gasparino, H Idei, S Kubo, H Maassberg, N Marushchenko, N Nakajima, M Romé and M Okamoto,  
*5D Simulation Study of Suprathermal Electron Transport in Non-Axisymmetric Plasmas*, Oct 1998  
(IAEA-CN-69/THP1/01)
- NIFS-577 S Fujiwara and T Sato,  
*Molecular Dynamics Simulation of Structure Formation of Short Chain Molecules*, Nov 1998
- NIFS-578 T Yamagishi,  
*Eigenfunctions for Vlasov Equation in Multi-species Plasmas* Nov 1998
- NIFS-579 M Tanaka, A Yu Grosberg and T Tanaka,  
*Molecular Dynamics of Strongly-Coupled Multichain Coulomb Polymers in Pure and Salt Aqueous*

*Solutions*; Nov. 1998

- NIFS-580 J. Chen, N. Nakajima and M. Okamoto,  
*Global Mode Analysis of Ideal MHD Modes in a Heliotron/Torsatron System: I. Mercier-unstable Equilibria*; Dec. 1998
- NIFS-581 M. Tanaka, A. Yu Grosberg and T. Tanaka,  
*Comparison of Multichain Coulomb Polymers in Isolated and Periodic Systems: Molecular Dynamics Study*; Jan. 1999
- NIFS-582 V.S. Chan and S. Murakami,  
*Self-Consistent Electric Field Effect on Electron Transport of ECH Plasmas*; Feb. 1999
- NIFS-583 M. Yokoyama, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani,  
*Roles of Bumpy Field on Collisionless Particle Confinement in Helical-Axis Heliotrons*; Feb. 1999
- NIFS-584 T.-H. Watanabe, T. Hayashi, T. Sato, M. Yamada and H. Ji,  
*Modeling of Magnetic Island Formation in Magnetic Reconnection Experiment*; Feb. 1999
- NIFS-585 R. Kumazawa, T. Mutoh, T. Seki, F. Shinpo, G. Nomura, T. Ido, T. Watari, Jean-Marie Noterdaeme and Yangping Zhao,  
*Liquid Stub Tuner for Ion Cyclotron Heating*; Mar. 1999
- NIFS-586 A. Sagara, M. Ima, S. Inagaki, N. Inoue, H. Suzuki, K. Tsuzuki, S. Masuzaki, J. Miyazawa, S. Morita, Y. Nakamura, N. Noda, B. Peterson, S. Sakakibara, T. Shimojima, H. Yamada, K. Akaishi, H. Chikaraishi, H. Funaba, O. Kaneko, K. Kawahata, A. Komori, N. Ohyabu, O. Motojima, LHD Exp. Group 1, LHD Exp. Group 2,  
*Wall Conditioning at the Starting Phase of LHD*; Mar. 1999
- NIFS-587 T. Nakamura and T. Yabe,  
*Cubic Interpolated Propagation Scheme for Solving the Hyper-Dimensional Vlasov-Poisson Equation in Phase Space*; Mar. 1999
- NIFS-588 W.X. Wnag, N. Nakajima, S. Murakami and M. Okamoto,  
*An Accurate  $\delta f$  Method for Neoclassical Transport Calculation*; Mar. 1999
- NIFS-589 K. Kishida, K. Araki, S. Kishiba and K. Suzuki,  
*Local or Nonlocal? Orthonormal Divergence-free Wavelet Analysis of Nonlinear Interactions in Turbulence*; Mar. 1999
- NIFS-590 K. Araki, K. Suzuki, K. Kishida and S. Kishiba,  
*Multiresolution Approximation of the Vector Fields on  $T^3$* ; Mar. 1999
- NIFS-591 K. Yamazaki, H. Yamada, K.Y. Watanabe, K. Nishimura, S. Yamaguchi, H. Nakanishi, A. Komori, H. Suzuki, T. Mito, H. Chikaraishi, K. Murai, O. Motojima and the LHD Group,  
*Overview of the Large Helical Device (LHD) Control System and Its First Operation*; Apr. 1999
- NIFS-592 T. Takahashi and Y. Nakao,  
*Thermonuclear Reactivity of D-T Fusion Plasma with Spin-Polarized Fuel*; Apr. 1999
- NIFS-593 H. Sugama,  
*Damping of Toroidal Ion Temperature Gradient Modes*; Apr. 1999
- NIFS-594 Xiaodong Li,  
*Analysis of Crowbar Action of High Voltage DC Power Supply in the LHD ICRF System*; Apr. 1999
- NIFS-595 K. Nishimura, R. Horiuchi and T. Sato,  
*Drift-kink Instability Induced by Beam Ions in Field-reversed Configurations*; Apr. 1999
- NIFS-596 Y. Suzuki, T.-H. Watanabe, T. Sato and T. Hayashi,  
*Three-dimensional Simulation Study of Compact Toroid Plasmoid Injection into Magnetized Plasmas*; Apr. 1999
- NIFS-597 H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida, S. Toda, S.-I. Itoh, M. Yagi and A. Fukuyama,  
*Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma*; May 1999