

NATIONAL INSTITUTE FOR FUSION SCIENCE

First ICRF Heating Experiment in the Large Helical Device

T. Mutoh, R. Kumazawa, T. Seki, K. Saito, F. Simpo, G. Nomura, T. Watari, X. Jikang, G. Cattanei, H. Okada, K. Ohkubo, M. Sato, S. Kubo, T. Shimoizuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B.J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Narihara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohya, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Sato, T. Morisaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K. Ida, N. Noda, K. Yamazaki, K. Akaishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iiyoshi, LHD Exp. Group 1 and 2

(Received - June 10, 1999)

NIFS-602

July 1999

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

RESEARCH REPORT
NIFS Series

First ICRF Heating Experiment in the Large Helical Device

T. Mutoh, R. Kumazawa, T. Seki, K. Saito¹, F. Shimpō, G. Nomura, T. Watari, X. Jikang², G. Cattanei³, H. Okada⁴, K. Ohkubo, M. Sato, S. Kubo, T. Shimozuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B. J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Narihara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohyabu, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Sato, T. Morisaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K. Ida, N. Noda, K. Yamazaki, K. Akaishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iiyoshi, LHD Exp.Group1 and 2

National Institute for Fusion Science, 322-6 Oroshi-cho, Toki, 509-5292 Japan

¹ Faculty of Engineering, Nagoya University, Furo-cho, Nagoya, Japan

² Institute of Plasma Physics, Academia Sinica, 230031, Hefei, Anhui, China

³ Max-planck-Institut fuer Plasmaphysik, Euratom Association, D-85748, Garching, Germany

⁴ Institute of Advanced Energy, Kyoto University, Gokasho, Uji, Kyoto, Japan

Keywords: LHD, ICRF heating, loop antenna, coupling resistance, wave damping

INTRODUCTION

First ICRF heating experiment in the Large Helical Device (LHD) was carried out in the 2nd experimental campaign[1],[2] in the end of 1998. During the experimental series, maximum 300kW/0.2sec of ICRF power was injected to the LHD plasma by using a pair of loop antennas. This paper reports the installation of antennas and the results of coupling and heating experiments.

Target plasma was produced by 2nd harmonic ECH heating of 84GHz gyrotrons at a magnetic field of 1.5 Tesla. Target plasma density was quite low of less than $1 \times 10^{19} \text{m}^{-3}$.



Fig.1 Photograph of ICRF loop antennas in LHD vacuum chamber

Therefore the maximum injected power was limited by low coupling resistance of plasma. Center current strap is

one strap type, therefore the parallel wave number is not regulated. Usually antenna of one strap type from low field side excitation has an impurity problem in tokamaks. However in helical device as LHD, this antenna location is the high field side excitation even if antenna is located at the outward side. Faraday shield is single layer and each pipe is directed to the external DC magnetic field. The heating efficiency and heating species were changed by the minority ion gas-puffing rate. For the optimum condition of the minority ion ratio (H/He=0.3), the plasma internal energy was increased from 13kJ to 26kJ. The heating power of ECH and ICRF were about 300kW in each. The diamagnetic measurement showed that ICRF heating has good efficiency comparable to the ECH heating in this power range.

The position of the launcher section of the loop antenna can be changed. By changing the distance between the last closed flux surface (LCFS) and the launcher front from 9cm to 5cm, the antenna coupling resistance was almost doubled. In the ICRF heating experiment, the antenna front was usually placed at around 6cm from the last closed flux surface.

ANTENNA INSTALLATION AND PLASMA COUPLING

One pair of the loop antennas was inserted from the top and the bottom vertical ports of the LHD. These antennas are designed to launch the fast magnetosonic wave in the ion cyclotron range of frequency. Photograph of the loop antennas in the LHD vacuum chamber is shown in Fig.1. The launcher section in the figure is divided to two separate parts. Upper section is fed through the top port and lower section is fed through the bottom port. The antenna launching section is just in front of the helical winding which is seen as ridged

helical structures and is placed on the outward side of toroid. The shape of the launcher is designed and fabricated to fit the plasma scrape off layer. The shape is 3 dimensional twisted form as shown in the figure. The total sizes of the antenna launching section are 120 cm in length (include upper and lower antenna), 46 cm in width and 17 cm in depth.

The antennas have strong cooling channels for long pulse operation. The steady state technologies obtained in the R&D tests are fully incorporated in the design of the loop antenna [3]. The swing motion with the pivot point near the feedthrough section realizes position shift of the launcher section. The shift length is 15cm and elapsed time is around 10 sec. Throughout the experiment, the antenna position was changed to fit the experimental requests of the LHD.




Fig.2 Increment of coupling resistance due to plasma is shown by changing the antenna positions (Faraday shield surface) between plasma last closed flux surface (LCFS) and vacuum vessel wall.




Fig.3 Calculation of antenna loading resistance by slab plasma model. Antenna structures of length, width, depth and shorted end are included in the model. Calculation parameters are same with in Fig.2.

Plasma coupling resistance was measured by changing the distance between the antenna and the plasma surface. In Fig.2, the coupling increments due to the plasma are shown. On the equatorial plane, last closed flux surface (LCFS) is at 417cm and vacuum

vessel wall is at 451.5cm. Experimental conditions are as follows; $B=1.5\text{ Tesla}$, $n_{e\text{-average}}=8 \times 10^{19} \text{ m}^{-3}$, $H/(H+He)=0.3$, frequency= 25.6MHz. In both antenna loops, coupling resistances were varied around twice by changing the positions. The vacuum loading level is almost half value of the nearest ones. Therefore the coupled power was 60-70% of the transmitter power. The coupling resistance values of two antenna loops were different. The reason of this difference is not clear. One of the candidates is the difference of antenna grounding structure. The lower antenna has three thin copper straps to ground the antenna back-plate to the vacuum vessel wall. On the other hand, there is no strap for grounding at the upper antenna.

The antenna resistance was calculated analytically to compare the experimental data. In Fig.3, theoretically calculated resistance was shown by changing the antenna positions. The model is two dimensional slab model and basic formulation is same with that of Reference [4]. The antenna current strap model includes length, width and shorted end and feeder section. And plasma is one dimensional slab structure. The range of the experimental condition in Fig.2 is between from 7cm to 11cm on horizontal axis. The absolute value of the calculation is larger than the experimental value about factor two. It mainly depends on the boundary parameters on the calculation. But the decay curve is




Fig.4 Time evolution of plasma parameters of ICRF heated plasma and target ECH plasma. ($B=1.5\text{ Tesla}$, $n_{e\text{-average}}=8 \times 10^{19} \text{ m}^{-3}$, $H/(H+He)=0.3$, Frequency = 25.6MHz)

almost same with the experimental ones. This agreement suggests that the antenna coupling on the experiment was mainly due to the fast magnetosonic wave.

HEATING EXPERIMENTS

Heating experiment was carried out when the magnetic field of LHD was 1.5 Tesla and the RF frequency was 25.6 MHz. In this condition, the cyclotron layer and the mode conversion layer were located at the 0.5-0.7 of radius. Time evolutions of the plasma parameters are shown in Fig.4. The stored energy, line averaged electron density, impurity line (OV) and radiation power are shown. Dotted line is of the ECH target plasma and solid line is of the ICRF heated plasma. ICRF pulse was applied at 0.3 sec to 0.5sec and about one third of the duration was overlapped with the ECH pulse. Plasma energy was raised from 13kJ to 23.5kJ. During the ICRF operation, the plasma density is sustained at constant level.

Electron temperature profiles of target plasma and




Fig.5 Electron temperature profiles of target ECH plasma and ICRF heated plasma at 0.38sec. of Fig.4.

ICRF heated plasma are shown in Fig.5. In all radius, electron temperatures were increased. However the increasing rate in peripheral region was higher than in central region. The mode conversion region from the fast wave to the IBW was located at 0.65-0.8 radius and it agreed that the peripheral region was apparently heated. Radiation power and some impurity lines are increased during the ICRF pulse. The stored energy began to decrease during the ICRF pulse and this is consistent with the increase of the radiation power. Oxygen and iron impurity lines were increased during ICRF pulse. These increases depended on the minority ion ratio, which is directly related with the wave damping mechanism.

Heating properties were largely affected by the minority ion ratio $H/(H+He)$. In Fig.6, the increments of




Fig.6 Increments of total stored energy W_p , and electron stored energy W_e versus input gas puff ratio $H/(He+H)$. (W_p ; diamagnetic loop, W_e ; Thomson $T_e(r)$ and FIR $n_e(r)$)

total stored energy W_p , and the incremental electron stored energy W_e , due to ICRF heating are shown by changing the minority ion ratio. This experiment was conducted by increasing the hydrogen gas-puff amount. On the other hand the majority helium gas was constant. Therefore the electron density was slightly increased as increasing the minority ion ratio. W_p is diamagnetic signal and electron stored energy W_e is calculated from the temperature of Thomson scattering and the density of FIR interferometer. To achieve the high W_p discharges, it was necessary to add the minority ion of over than 20 %. If the H ratio was less than 15%, no electron heating was observed. W_p - W_e seems to be the increment of ion component.

To understand the wave heating mechanism in the LHD, one dimensional full wave analysis code [5] was used. This code includes the effects of the wave propagation, damping process, mode conversion, eigen mode formation in one dimension and current drive. Figure 7 shows the power partitions to plasma species from the launched wave. The calculation was done for the parameters along the line on the equatorial plane of LHD. On the low H ratio region, there is small electron heating and hydrogen cyclotron damping mainly occurred. This is so-called minority heating region. On the medium H ratio region, there is electron heating by mode conversion. In the higher H ratio region, fast wave evanescent layer becomes thick, then excited fast wave is reflected at a left-hand cut-off. In this range, minority ion cyclotron damping is dominant.

The experimental results of Fig.6 agree qualitatively with this calculation. The gas puff ratio of $H/(H+He)$ was not exactly appeared on plasma species. Due to the high recycling rate of helium particles, the proton ratio in plasma was seemed to be lower than the input gas puff ratio.




Fig.7 Full wave analysis on a slab plasma model shows that electron absorption occurs at minority hydrogen content of 5 to 15 % of electron.

SUMMARY

Initial ICRF heating experiment in the LHD was carried out in 1998. One pair of the movable loop antennas was used and the coupling resistance was around one ohm for the low density ECH plasma. The loading characteristics were consistent with the fast wave excitation. By applying the ICRF heating of 300 kW to the ECH target plasma, the diamagnetic energy was increased from 13 kJ to 26 kJ. The heating performance was decided by hydrogen mixture rate on puffing gas. Efficient electron heating was observed at higher hydrogen gas ratio. These results can be explained by the one dimensional wave analysis calculation on slab plasma model. On the experiment of 1999, the magnetic field of the LHD will be increased to 2.8 Tesla and the RF frequency will be also raised. The higher frequency and higher density plasma should lead to the better plasma coupling and higher heating power. In addition, another new type antenna (Folded Wave-Guide antenna) is installed and will be operated in 1999.

REFERENCES:

- [1] O. Motojima, et al., Physics of Plasmas, Vol.6, No5, (1999) 1843
- [2] M. Fujiwara, et al., this conference

- [3] T. Mutoh, et al., Journal of Plasma and Fusion Research, SERIES, Vol.1, (1998) 334
- [4] K.Theilhaber, J.Jacquinot, Nuclear Fusion, Vol.24, (1984) 541
- [5] A. Fukuyama, et al., Computer Physics Report 4 (1986) 137

Recent Issues of NIFS Series

- NIFS-539 Takashi Yabe and Yan Zhang,
Effect of Ambient Gas on Three-Dimensional Breakup in Coronet Formation Process, Feb. 1998
- NIFS-540 H. Nakamura, K. Ikeda and S. Yamaguchi,
Transport Coefficients of InSb in a Strong Magnetic Field, Feb 1998
- NIFS-541 J. Uramoto,
Development of v_μ Beam Detector and Large Area v_μ Beam Source by H_2 Gas Discharge (I); Mar. 1998
- NIFS-542 J. Uramoto,
Development of $\bar{v}_\mu$ Beam Detector and Large Area $\bar{v}_\mu$ Beam Source by H_2 Gas Discharge (II); Mar. 1998
- NIFS-543 J. Uramoto,
Some Problems inside a Mass Analyzer for Pions Extracted from a H_2 Gas Discharge; Mar 1998
- NIFS-544 J. Uramoto,
Simplified v_μ $\bar{v}_\mu$ Beam Detector and v_μ $\bar{v}_\mu$ Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch; Mar 1998
- NIFS-545 J. Uramoto,
Various Neutrino Beams Generated by D_2 Gas Discharge; Mar.1998
- NIFS-546 R. Kanno, N. Nakajima, T. Hayashi and M. Okamoto,
Computational Study of Three Dimensional Equilibria with the Bootstrap Current; Mar. 1998
- NIFS-547 R. Kanno, N. Nakajima and M. Okamoto,
Electron Heat Transport in a Self-Similar Structure of Magnetic Islands; Apr. 1998
- NIFS-548 J.E. Rice,
Simulated Impurity Transport in LHD from MIST; May 1998
- NIFS-549 M.M. Skoric, T. Sato, A.M. Maluckov and M.S. Jovanovic,
On Kinetic Complexity in a Three-Wave Interaction; June 1998
- NIFS-550 S. Goto and S. Kida,
Passive Saclar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation; June 1998
- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,
Initial Value Problem of the Toroidal Ion Temperature Gradient Mode ; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watan,
Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device ; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,
Oronization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-; July 1998
- NIFS-554 S. Murakami and H. Saleem,
Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations, July 1998
- NIFS-555 H. Nakamura, K. Ikeda and S. Yamaguchi,
Physical Model of Nernst Element; Aug 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,
Numerical Computation of Thermoelectric and Thermomagnetic Effects; Aug. 1998
- NIFS-557 Y. Takeiri, M. Osakabe, K. Tsumon, Y. Oka, O. Kaneko, E. Asano, T. Kawamoto, R. Akiyama and M. Tanaka,
Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System; Aug 1998

- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Structure Organization of Polyampholytes; Sep. 1998
- NIFS-559 R. Horiuchi, K. Nishimura and T. Watanabe,
Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations; Sep. 1998
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Kholopenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,
Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection; Oct. 1998
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,
Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron; Oct. 1998
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,
Next Generation Toroidal Devices; Oct. 1998
- NIFS-563 Y. Todo and T. Sato,
Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes; Oct. 1998
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watari, T. Shimoizuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Idei, Y. Oka, M. Osakabe, T. Seki, K. Tsumori, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Shimo, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Morimoto, T. Bigelow and Y.P. Zhao,
Steady State Heating Technology Development for LHD; Oct. 1998
(IAEA-CN-69/FTP/21)
- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyabu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda,
LHD-Type Compact Helical Reactors; Oct. 1998
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems; Oct. 1998
(IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D.S. Darrow, H. Idei, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tsumori, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watari,
Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EX2/2)
- NIFS-568 S. Okamura, K. Matsuoka, R. Akiyama, D.S. Darrow, A. Ejiri, A. Fujisawa, M. Fujiwara, M. Goto, K. Ida, H. Idei, H. Iguchi, N. Inoue, M. Isobe, K. Itoh, S. Kado, K. Kholopenkov, T. Kondo, S. Kubo, A. Lazaros, S. Lee, G. Matsunaga, T. Minami, S. Morita, S. Murakami, N. Nakajima, N. Nikai, S. Nishimura, I. Nomura, S. Ohdachi, K. Ohkuni, M. Osakabe, R. Pavlichenko, B. Peterson, R. Sakamoto, H. Sanuki, M. Sasao, A. Shimizu, Y. Shirai, S. Sudo, S. Takagi, C. Takahashi, S. Takayama, M. Takechi, K. Tanaka, K. Toi, K. Yamazaki, Y. Yoshimura and T. Watari,
Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS; Oct. 1998
(IAEA-CN-69/OV4/5)
- NIFS-569 M.M. Skoric, T. Sato, A. Maluckov, M.S. Jovanovic,
Micro- and Macro-scale Self-organization in a Dissipative Plasma; Oct. 1998
- NIFS-570 T. Hayashi, N. Mizuguchi, T.-H. Watanabe, T. Sato and the Complexity Simulation Group,
Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak; Oct. 1998
(IAEA-CN-69/TH3/3)
- NIFS-571 A. Iiyoshi, A. Komori, A. Ejiri, M. Ernsto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimoizuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, H. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, S. Iima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, M. Fujiwara and O. Motojima,

An Overview of the Large Helical Device Project; Oct 1998
(IAEA-CN-69/OV1/4)

- NIFS-572 M Fujiwara, H Yamada, A Ejiri, M Emoto, H Funaba, M Goto, K Ida, H Idei, S Inagaki, S Kado, O Kaneko, K Kawahata, A Komori, S Kubo, R Kumazawa, S Masuzaki, T Minami, J Miyazawa, T Morisaki, S Morita, S Murakami, S Muto, T Muto, Y Nagayama, Y Nakamura, H Nakanishi, K Narihara, K Nishimura, N Noda, T Kobuchi, S Ohdachi, N Ohyaibu, Y Oka, M Osakabe, T Ozaki, B. J. Peterson, A Sagara, S. Sakakibara, R. Sakamoto, H Sasao, M Sasao, K Sato, M. Sato, T Seki, T Shimozuma, M Shoji, H Sazuki, Y Takein, K. Tanaka, K. Toi, T Tokuzawa, K Tsumon, I. Yamada, S. Yamaguchi, M. Yokoyama, K.Y. Watanabe, T Watan, R. Akiyama, H. Chikaraishi, K Haba, S. Hamaguchi, M Iima, S. Imagawa, N Inoue, K Iwamoto, S Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iiyoshi,
Plasma Confinement Studies in LHD; Oct 1998
(IAEA-CN-69/EX2/3)
- NIFS-573 O. Motojima, K. Akaishi, H. Chikaraishi, H. Funaba, S. Hamaguchi, S. Imagawa, S. Inagaki, N. Inoue, A. Iwamoto, S. Kitagawa, A. Komori, Y. Kubota, R. Maekawa, S. Masuzaki, T. Mito, J. Miyazawa, T. Monsaki, T. Muroga, T. Nagasaka, Y. Nakamura, A. Nishimura, K. Nishimura, N. Noda, N. Ohyaibu, S. Sagara, S. Sakakibara, R. Sakamoto, S. Satoh, T. Satow, M. Shoji, H. Suzuki, K. Takahata, H. Tamura, K. Watanabe, H. Yamada, S. Yamada, S. Yamaguchi, K. Yamazaki, N. Yanagi, T. Baba, H. Hayashi, M. Iima, T. Inoue, S. Kato, T. Kato, T. Kondo, S. Moriuchi, H. Ogawa, I. Ohtake, K. Ooba, H. Sekiguchi, N. Suzuki, S. Takami, Y. Taniguchi, T. Tsuzuki, N. Yamamoto, K. Yasui, H. Yonezu, M. Fujiwara and A. Iiyoshi,
Progress Summary of LHD Engineering Design and Construction; Oct. 1998
(IAEA-CN-69/FT2/1)
- NIFS-574 K. Toi, M. Takechi, S. Takagi, G. Matsunaga, M. Isobe, T. Kondo, M. Sasao, D.S. Darrow, K. Ohkuni, S. Ohdachi, R. Akiyama, A. Fujisawa, M. Gotoh, H. Idei, K. Ida, H. Iguchi, S. Kado, M. Kojima, S. Kubo, S. Lee, K. Matsuoka, T. Minami, S. Morita, N. Nikai, S. Nishimura, S. Okamura, M. Osakabe, A. Shimizu, Y. Shirai, C. Takahashi, K. Tanaka, T. Watari and Y. Yoshimura,
Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EXP1/19)
- NIFS-575 Y. Hamada, A. Nishizawa, Y. Kawasumi, A. Fujisawa, M. Kojima, K. Narihara, K. Ida, A. Ejiri, S. Ohdachi, K. Kawahata, K. Toi, K. Sato, T. Seki, H. Iguchi, K. Adachi, S. Hidekuma, S. Hirokura, K. Iwasaki, T. Ido, R. Kumazawa, H. Kuramoto, T. Minami, I. Nomura, M. Sasao, K.N. Sato, T. Tsuzuki, I. Yamada and T. Watan,
Potential Turbulence in Tokamak Plasmas; Oct. 1998
(IAEA-CN-69/EXP2/14)
- NIFS-576 S. Murakami, U. Gasparino, H. Idei, S. Kubo, H. Maassberg, N. Marushchenko, N. Nakajima, M. Romé and M. Okamoto,
5D Simulation Study of Suprathermal Electron Transport in Non-Axisymmetric Plasmas; Oct. 1998
(IAEA-CN-69/THP1/01)
- NIFS-577 S. Fujiwara and T. Sato,
Molecular Dynamics Simulation of Structure Formation of Short Chain Molecules; Nov. 1998
- NIFS-578 T. Yamagishi,
Eigenfunctions for Vlasov Equation in Multi-species Plasmas Nov. 1998
- NIFS-579 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Strongly-Coupled Multichain Coulomb Polymers in Pure and Salt Aqueous Solutions; Nov. 1998
- NIFS-580 J. Chen, N. Nakajima and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in a Heliotron/Torsatron System: I. Mercier-unstable Equilibria; Dec. 1998
- NIFS-581 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Comparison of Multichain Coulomb Polymers in Isolated and Periodic Systems: Molecular Dynamics Study; Jan. 1999
- NIFS-582 V.S. Chan and S. Murakami,
Self-Consistent Electric Field Effect on Electron Transport of ECH Plasmas, Feb. 1999
- NIFS-583 M. Yokoyama, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani,
Roles of Bumpy Field on Collisionless Particle Confinement in Helical-Axis Heliotrons; Feb. 1999
- NIFS-584 T.-H. Watanabe, T. Hayashi, T. Sato, M. Yamada and H. Ji,
Modeling of Magnetic Island Formation in Magnetic Reconnection Experiment; Feb. 1999
- NIFS-585 R. Kumazawa, T. Mutoh, T. Seki, F. Shinpo, G. Nomura, T. Ido, T. Watari, Jean-Marie Noterdaeme and Yangping Zhao,
Liquid Stub Tuner for Ion Cyclotron Heating, Mar. 1999

- NIFS-586 A. Sagara, M. Ima, S. Inagaki, N. Inoue, H. Suzuki, K. Tsuzuki, S. Masuzaki, J. Miyazawa, S. Morita, Y. Nakamura, N. Noda, B. Peterson, S. Sakakibara, T. Shimoizuma, H. Yamada, K. Akaishi, H. Chikaraishi, H. Funaba, O. Kaneko, K. Kawahata, A. Komori, N. Ohyabu, O. Motojima, LHD Exp. Group 1, LHD Exp. Group 2,
Wall Conditioning at the Starting Phase of LHD; Mar. 1999
- NIFS-587 T. Nakamura and T. Yabe,
Cubic Interpolated Propagation Scheme for Solving the Hyper-Dimensional Vlasov-Poisson Equation in Phase Space; Mar. 1999
- NIFS-588 W.X. Wnag, N. Nakajima, S. Murakami and M. Okamoto,
An Accurate δf Method for Neoclassical Transport Calculation; Mar. 1999
- NIFS-589 K. Kishida, K. Araki, S. Kishiba and K. Suzuki,
Local or Nonlocal? Orthonormal Divergence-free Wavelet Analysis of Nonlinear Interactions in Turbulence; Mar. 1999
- NIFS-590 K. Araki, K. Suzuki, K. Kishida and S. Kishiba,
Multiresolution Approximation of the Vector Fields on T^3 ; Mar. 1999
- NIFS-591 K. Yamazaki, H. Yamada, K.Y. Watanabe, K. Nishimura, S. Yamaguchi, H. Nakanishi, A. Komori, H. Suzuki, T. Mito, H. Chikaraishi, K. Murai, O. Motojima and the LHD Group,
Overview of the Large Helical Device (LHD) Control System and Its First Operation; Apr. 1999
- NIFS-592 T. Takahashi and Y. Nakao,
Thermonuclear Reactivity of D-T Fusion Plasma with Spin-Polarized Fuel; Apr. 1999
- NIFS-593 H. Sugama,
Damping of Toroidal Ion Temperature Gradient Modes, Apr. 1999
- NIFS-594 Xiaodong Li,
Analysis of Crowbar Action of High Voltage DC Power Supply in the LHD ICRF System; Apr. 1999
- NIFS-595 K. Nishimura, R. Horiuchi and T. Sato,
Drift-kink Instability Induced by Beam Ions in Field-reversed Configurations; Apr. 1999
- NIFS-596 Y. Suzuki, T.-H. Watanabe, T. Sato and T. Hayashi,
Three-dimensional Simulation Study of Compact Toroid Plasmoid Injection into Magnetized Plasmas; Apr. 1999
- NIFS-597 H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida, S. Toda, S.-I. Itoh, M. Yagi and A. Fukuyama,
Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma; May 1999
- NIFS-598 S. Nakazawa, N. Nakajima, M. Okamoto and N. Ohyabu,
One Dimensional Simulation on Stability of Detached Plasma in a Tokamak Divertor; June 1999
- NIFS-599 S. Murakami, N. Nakajima, M. Okamoto and J. Nhrnberg,
Effect of Energetic Ion Loss on ICRF Heating Efficiency and Energy Confinement Time in Heliotrons; June 1999
- NIFS-600 R. Horiuchi and T. Sato,
Three-Dimensional Particle Simulation of Plasma Instabilities and Collisionless Reconnection in a Current Sheet; June 1999
- NIFS-601 W. Wang, M. Okamoto, N. Nakajima and S. Murakami,
Collisional Transport in a Plasma with Steep Gradients; June 1999
- NIFS-602 T. Mutoh, R. Kumazawa, T. Seki, K. Sato, F. Simpo, G. Nomura, T. Watari, X. Jikang, G. Cattanei, H. Okada, K. Ohkubo, M. Sato, S. Kubo, T. Shimoizuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B.J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Narihara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohyabu, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Sato, T. Morisaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K. Ida, N. Noda, K. Yamazaki, K. Akaishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iiyoshi, LHD Exp. Group 1 and 2,
First ICRF Heating Experiment in the Large Helical Device; July 1999