

NATIONAL INSTITUTE FOR FUSION SCIENCE

δf Simulation of Ion Neoclassical Transport

W. Wang, N. Nakajima, M. Okamoto and S. Murakami

(Received - June 29, 1999)

NIFS-604

July 1999

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

**RESEARCH REPORT
NIFS Series**

NAGOYA, JAPAN

δf Simulation of Ion Neoclassical Transport*

W. Wang^{1,†}, N. Nakajima^{1,2}, M. Okamoto^{1,2,‡}, and S. Murakami^{1,2}

¹*National Institute for Fusion Science*

and

²*Graduate University for Advanced Studies*

322-6 Oroshir-cho, Toki, Gifu 509-5292, Japan

Abstract

Ion neoclassical transport with finite orbit width dynamics is calculated over whole poloidal cross section by using accurate δf method which employs an improved like-particle collision operator and an accurate weighting scheme to solve drift kinetic equation. Ion thermal transport near magnetic axis shows a great reduction from its conventional neoclassical level due to non-standard orbit topology, like that of previous δf simulation. On other hand, the direct particle loss from confinement region may strongly increase ion energy transport near the edge. It is found that ion parallel flow near the axis is also largely reduced due to non-standard orbit topology. In the presence of steep density gradient, ion thermal conductivity is significantly reduced, and an ion particle flux is driven by self-collision alone.

Keywords : neoclassical transport, finite orbit width dynamics, δf method,

The study of neoclassical transport phenomena is newly motivated by recent development in fusion experiments where plasmas are operated in parameter regimes beyond the valid scope of the conventional neoclassical theory [1]. The modifications to the neoclassical theory are required for taking into account finite orbit width dynamics, strong radial electric field with large radial gradient, and non-standard orbit topology near magnetic axis. The low noise δf particle simulation, solving drift kinetic equation, can be a powerful tool for neoclassical transport calculation [2,3].

We solve linearized drift kinetic equation (distribution function $f = f_0 + f_1$ with f_0 and f_1 being the zeroth and first order parts, respectively)

$$\frac{D}{Dt}f_1 = -\vec{v}_d \cdot \nabla f_0 + C(f_0, f_1), \quad (1)$$

from the dynamics of a finite number of particles (markers), where the notation D/Dt denotes, for an arbitral function f_A ,

$$\frac{D}{Dt}f_A = \frac{\partial f_A}{\partial t} + (\vec{v}_{||} + \vec{v}_d) \cdot \nabla f_A - C(f_A, f_0), \quad (2)$$

and C is the Coulomb collision operator. Here the drift term is retained to the leading order in order to take into account finite orbit effects. Each marker

is pushed in the extended phase space (\vec{r}, \vec{v}, w) along characteristic

$$\frac{d\vec{r}}{dt} = \dot{\vec{r}}, \quad \frac{d\vec{v}}{dt} = \dot{\vec{v}}, \quad \text{and} \quad \frac{dw}{dt} = \dot{w}. \quad (3)$$

The numerical representation of f_1 is

$$f_1(\vec{r}, \vec{v}, t) = \sum_j w_j \delta(\vec{r} - \vec{r}_j(t)) \delta(\vec{v} - \vec{v}_j(t)). \quad (4)$$

The particle weight w is calculated from

$$\dot{w} = \frac{1}{g} \left[- \int w S_M dw - \vec{v}_d \cdot \nabla f_0 + C(f_0, f_1) \right], \quad (5)$$

where S_M is the marker source used to control marker population and g is marker density [4].

Accurate implementation of Coulomb collisions is an important ingredient for neoclassical transport calculation. Collisions between ions and electrons can be simply modeled by employing the large mass ratio approximation. Here we present a linear like-particle collision model, accurately conserving all particle number, momentum, and energy for δf simulation. Our collision model reads

$$C^\ell(f_1) = C_0^\ell + [P_1 + Q_1 + P_2 + Q_2] f_M, \quad (6)$$

where

$$C_0^\ell(f_1) = C_{TP}(f_1) + Pf_M, \quad (7)$$

$$P(\vec{r}, \vec{v}) = \vec{A} \cdot \frac{d}{dt} \langle \vec{v} \rangle_{TP} + B \frac{d}{dt} \langle v^2 \rangle_{TP}, \quad (8)$$

$$P_1(\vec{r}, \vec{v}) = \vec{A} \cdot \frac{d}{dt} \langle \vec{v} \rangle_{TP} + B \frac{d}{dt} \langle v^2 \rangle_{TP+P}, \quad (9)$$

$$Q_1(\vec{r}, \vec{v}) = D \frac{d}{dt} \langle v^0 \rangle_P, \quad (10)$$

$$P_2 = \vec{A} \cdot \frac{d}{dt} \langle \vec{v} \rangle_{TP+P+P_1+Q_2} + B \frac{d}{dt} \langle v^2 \rangle_{TP+P+P_1+Q_2}, \quad (11)$$

$$Q_2(\vec{r}, \vec{v}) = D \frac{d}{dt} \langle v^0 \rangle_{P+P_1+Q_2}, \quad (12)$$

$$\vec{A} = -6 \sqrt{\frac{\pi}{2}} \phi \frac{v_{th} \vec{v}}{v^3}, \quad (13)$$

$$B = -2 \sqrt{\frac{\pi}{2}} \left[\phi - \frac{d\phi}{dy} \right] \frac{1}{vv_{th}}, \quad (14)$$

$$D(\vec{r}, \vec{v}) = 3 \sqrt{\frac{\pi}{2}} \left[\phi - \frac{d\phi}{dy} \right] \frac{v_{th}}{v} - 1, \quad (15)$$

and $\phi(y)$ is Maxwellian integral with $y = v^2/v_{th}^2$. The $C_{TP}(f_1)$ is drag and diffusion part that describes the test particle (f_1) collisions with Maxwellian field particles (f_M). The Pf_M originating from $C(f_M, f_1)$ works for compensating $\frac{d}{dt} \langle \vec{v} \rangle_{TP}$ and $\frac{d}{dt} \langle v^2 \rangle_{TP}$, the averaged momentum and energy losses due to C_{TP} , respectively. C_0^ℓ represents the previous collision operator [2,5]. It is shown that the accuracy in particle, momentum, and energy conservation is insufficient with using C_0^ℓ . There are still considerable non-vanishing $\frac{d}{dt} \langle \vec{v} \rangle_{TP+P}$, $\frac{d}{dt} \langle v^2 \rangle_{TP+P}$ and $\frac{d}{dt} \langle v^0 \rangle_P$ coming from the use of a finite number of particles, and the numerical errors may get so enlarged as to distort simulation results. The compensation terms are then developed to restore the conservations. For example, $P_1 f_M$ is introduced to compensate the momentum and energy loss, $\frac{d}{dt} \langle \vec{v} \rangle_{TP+P}$ and $\frac{d}{dt} \langle v^2 \rangle_{TP+P}$, and $Q_1 f_M$ is used to compensate $\frac{d}{dt} \langle v^0 \rangle_P$ (the particle loss coming from the implementation of Pf_M). With the suggested compensation terms, all the three quantities are conserved nearly perfectly. The calculations of neoclassical transport using the present collision scheme demonstrate the increased accuracy in results. A shifted Maxwellian solution of ion drift kinetic equation under $\nabla T = 0$, which gives zero particle and energy flux, is recovered (Fig.1) owing to the use of the improved operator.

The calculation of particle weight is crucial in a δf simulation. It is found that valid results essentially rely on the correct evaluation of marker density g in weight calculation. The function g satisfies

$$\frac{D}{Dt} g = \int S_M dw. \quad (16)$$

Previous weighting schemes including the nonlinear

weighting scheme [2,6] employ an assumed g in weight equation for advancing particle weights. Such a scheme is found to be ineffective or inaccurate for solving the drift kinetic equation because of a severe constraint that the real marker distribution must be consistent with the initially assumed g during the simulation. Otherwise, correct results can not be guaranteed. Instead of using an approximation of g for advancing particle weights, we solve g directly from its kinetic equation using the idea of δf method. To this end, we employ a new weight function ω and a marker source Ω_M . The accurate weight equation for each marker is derived as

$$\dot{\omega} = \frac{1-\omega}{f_0} \left[- \int w S_M dw - \vec{v}_d \cdot \nabla f_0 + C(f_0, f_1) \right], \quad (17)$$

$$\omega = \frac{1-\omega}{f_0} \left[- \int \omega \Omega_M d\omega - \vec{v}_d \cdot \nabla f_0 + \int S_M dw \right]. \quad (18)$$

Now each simulation particle is assigned two weights rather than one. A convenient choice for the sources is $S_M = \nu(t)s(\vec{r})f_0\delta(w)$, and $\Omega_M = \nu(t)s(\vec{r})f_0\delta(\omega)$. This choice means that new Maxwellian markers with $w = \omega = 0$ are added in terms of the rate $\nu(t)$ and spatial distribution $s(\vec{r})$.

The results of simulations (solving Eq.(1) with $\nabla T = 0$) using the previous nonlinear scheme and the present scheme are compared in Fig 2. Because $g = f$, which is assumed for advancing particle weights, is not recognized in the simulation, the nonlinear weighting scheme fails to give the correct particle flux and energy flux. While the accurate weighting scheme works for the problem effectively ; it leads to the correct results with vanishing particle and energy fluxes over the whole cross section

A δf simulation code named “FORTEC” (Finite ORbit Transport study by the Extensive Code) has

been developed based on the above new method. By using “FORTEC”, the modifications to ion neoclassical transport from the finite orbit width dynamics are investigated. The simulations which are performed over global poloidal cross section enable us to calculate the neoclassical transport in both the region near magnetic axis and that far from it (including the edge region of confined plasma). Moreover, the global simulations enable us to calculate nonlocal transport fluxes which depend not only on the local gradients but on the entire density and temperature profiles as well. Our calculations focus on the banana-plateau regime in which the present fusion plasmas are most operated.

A typical feature of ion thermal conductivity over the whole cross section is shown in Fig.3. The ion thermal conductivity in the region within inner half radius is greatly reduced from its conventional neoclassical level due to non-standard orbit topology near the axis. This modification due to finite banana width dynamics was previously proposed [3] to interpret the very low ion thermal transport observed in the enhanced reversed shear (ERS) core plasmas. The reduction of ion thermal conductivity is attributed to both the decrease of orbit size and the increase of the trapped particle fraction. This finite orbit effects commonly exist in the tokamak plasmas near the axis. However, the reduced ion neoclassical thermal transport is revealed only as turbulence transport is suppressed, as achieved in ERS plasmas. In contrast to the region near the axis, the ion energy transport near the edge is strongly enhanced by the direct particle loss from the confinement region.

Another important transport quantity that is modified by the finite orbit effect is ion parallel flow. A

typical simulation result of ion parallel flow is shown in Fig.4. It is found that the ion parallel flow near the axis is also largely reduced as the finite orbit width dynamics is taken into account. The modification of the parallel flow again results from the non-standard orbit topology. For trapped ions, a net parallel flow results from the noncancellation of the parallel momentum carried by co-moving ions coming from the high temperature and density inner region and antiparallel momentum carried by counter-moving ions coming from the low temperature and density outer region. Therefore, the net parallel flow is driven by the plasma gradients, and is proportional to the radial excursion of trapped particles, the banana width Δ_b . It is found that the real banana width near the axis is smaller than the conventional neoclassical estimation. Then the finite banana effect could help reduce the parallel flow near the axis. But this reduction is small because the ion parallel flow is mostly contributed by passing ions rather than trapped ions. The point is as follows. The collisions between trapped and passing ions force the passing ion distribution function to be continuous with the trapped distribution. A displacement δ of particles from their home flux surface Ψ is defined by [7] $f = f_M(\epsilon, \Psi) - \delta \partial f_M / \partial r$ (ϵ : energy). For trapped ions, δ is about the half-width of the banana orbits. The continuity of the distribution function across the trapped-passing boundary implies that $\delta_p \approx \delta_b$. That is, the relative displacement of co-moving passing ions from counter-moving passing ions is determined by the orbit width of barely trapped ions, which is, in the region near the axis, proportional to the local minor radius r and is significantly smaller than the conventional neoclassical estimate. The parallel flow arising from the relative displace-

ment between co- and counter-moving passing ions is therefore reduced. The ion parallel flow represents a friction on electrons. As an interesting consequence, the modification of ion parallel flow may modify the standard neoclassical bootstrap current near the axis.

Figure 5(a) shows the dependence of ion thermal conductivity on density gradient length. When the banana width and the gradient length become comparable, the ion thermal conductivity is significantly reduced from its conventional neoclassical result. It is found from Fig.5(b) that in the presence of steep density gradient ion particle flux can be driven by ion-ion collisions alone. The simulation results suggest that the reduction of the conductivity and the particle flux depend linearly on $(1/n)dn/dr$. Since the finite orbit effect on electrons is very weak, the electron transport is unchanged in the presence of steep density gradient. Then the neoclassical particle transport could be non-ambipolar, or, a radial electric field can be established within the neoclassical framework in order to maintain an ambipolar neoclassical transport. The neoclassical transport with steep gradients is under investigation.

References

- [1] F.L. Hinton and R.D. Hazeltine, *Rev. Mod. Phys.* **48**, 239 (1976).
- [2] Z. Lin, W.M. Tang, and W.W. Lee, *Phys. Plasmas* **2**, 2975 (1995).
- [3] Z. Lin, W.M. Tang, and W.W. Lee, *Phys. Plasmas* **4**, 1707 (1997).
- [4] Y. Chen and R.B. White, *Phys. Plasmas* **4**, 3591 (1997).

- [5] A.M. Dimits and B.I. Cohen, *Phys. Rev. E* **49**, 709 (1994).
- [6] S. Parker and W.W. Lee, *Phys. Fluids B* **5**, 77 (1993).
- [7] A.H. Boozer and H.J. Gardner, *Phys. Fluids B* **2**, 2048 (1990).
- [8] C.S. Chang and F.L. Hinton, *Phys. Fluids* **25**, 1493 (1982).

*Presented at The 9th International Toki Conference on Plasma Physics and Controlled Nuclear Fusion (ITC-9) – *New Frontiers in Plasma Physics*, December 7 - 11, 1998, Toki, Japan

†Present address . Institute of Applied Physics and Computational Mathematics, P.O. Box 8009, Beijing 100088, P.R. China

‡Corresponding author's e-mail : okamoto@nifs.ac.jp

Figure captions

Fig.1 Comparison of simulation results from using present and previous collision operators, with zero orbit width and $\nabla T = 0$. Γ_0 and Q_0 : fluxes computed from using non-conservation operator C_{TP} ; $u_{\parallel T}$: analytical parallel velocity.

Fig.2 Comparison of simulation results from using present weighting scheme and the nonlinear weighting scheme, with finite orbit width and $\nabla T = 0$.

Fig.3 Ion thermal conductivity χ_r vs. minor radius r (theory 1-Ref.[1], theory 2-Ref [8]). An uniform q -profile ($q = 1.1$) is used here as well as in Fig.4 below.

Fig.4 Ion parallel flow u_{\parallel} vs. r .

Fig.5 Ion thermal conductivity χ_r and particle flux Γ vs. Δ_b (banana width) / L_n (density gradient scale length).




Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5

Recent Issues of NIFS Series

- NIFS-541 J. Uramoto,
Development of ν_μ Beam Detector and Large Area ν_μ Beam Source by H₂ Gas Discharge (I); Mar. 1998
- NIFS-542 J. Uramoto,
Development of $\bar{\nu}_\mu$ Beam Detector and Large Area $\bar{\nu}_\mu$ Beam Source by H₂ Gas Discharge (II); Mar. 1998
- NIFS-543 J. Uramoto,
Some Problems inside a Mass Analyzer for Pions Extracted from a H₂ Gas Discharge; Mar. 1998
- NIFS-544 J. Uramoto,
Simplified ν_μ $\bar{\nu}_\mu$ Beam Detector and ν_μ $\bar{\nu}_\mu$ Beam Source by Interaction between an Electron Bunch and a Positive Ion Bunch; Mar. 1998
- NIFS-545 J. Uramoto,
Various Neutrino Beams Generated by D₂ Gas Discharge; Mar. 1998
- NIFS-546 R. Kanno, N. Nakajima, T. Hayashi and M. Okamoto,
Computational Study of Three Dimensional Equilibria with the Bootstrap Current; Mar. 1998
- NIFS-547 R. Kanno, N. Nakajima and M. Okamoto,
Electron Heat Transport in a Self-Similar Structure of Magnetic Islands; Apr. 1998
- NIFS-548 J.E. Rice,
Simulated Impurity Transport in LHD from MIST; May 1998
- NIFS-549 M.M. Skonc, T. Sato, A.M. Maluckov and M.S. Jovanovic,
On Kinetic Complexity in a Three-Wave Interaction; June 1998
- NIFS-550 S. Goto and S. Kida,
Passive Scalar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation; June 1998
- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,
Initial Value Problem of the Toroidal Ion Temperature Gradient Mode ; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watari,
Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device ; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,
oronaization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-; July 1998
- NIFS-554 S. Murakami and H. Saleem,
Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations; July 1998
- NIFS-555 H. Nakamura , K. Ikeda and S. Yamaguchi,
Physical Model of Nernst Element; Aug. 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,
Numerical Computation of Thermoelectric and Thermomagnetic Effects; Aug. 1998
- NIFS-557 Y. Takeiri, M. Osakabe, K. Tsumori, Y. Oka, O. Kaneko, E. Asano, T. Kawamoto, R. Akiyama and M. Tanaka,
Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System; Aug. 1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Structure Organization of Polyampholytes; Sep. 1998
- NIFS-559 R. Horiuchi, K. Nishimura and T. Watanabe,
Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations; Sep. 1998
(IAEA-CN-69/THP1/11)

- NIFS-560 S. Sudo, K. Kholopenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,
Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection; Oct. 1998
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,
Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron; Oct. 1998
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,
Next Generation Toroidal Devices; Oct. 1998
- NIFS-563 Y. Todo and T. Sato,
Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes; Oct. 1998
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watari, T. Shimozuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Idei, Y. Oka, M. Osakabe, T. Seki, K. Tsumori, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Shimpo, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Morimoto, T. Bigelow and Y.P. Zhao,
Steady State Heating Technology Development for LHD; Oct. 1998
(IAEA-CN-69/FTP/21)
- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyabu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda,
LHD-Type Compact Helical Reactors; Oct. 1998
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems; Oct. 1998
(IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D.S.Darrow, H. Idei, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tumori, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watari,
Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EX2/2)
- NIFS-568 S. Okamura, K. Matsuoka, R. Akiyama, D.S. Darrow, A. Ejiri, A. Fujisawa, M. Fujiwara, M. Goto, K. Ida, H. Idei, H. Iguchi, N. Inoue, M. Isobe, K. Itoh, S. Kado, K. Khlopenkov, T. Kondo, S. Kubo, A. Lazaros, S. Lee, G. Matsunaga, T. Minami, S. Morita, S. Murakami, N. Nakajima, N. Nikai, S. Nishimura, I. Nomura, S. Ohdachi, K. Ohkuni, M. Osakabe, R. Pavlichenko, B. Peterson, R. Sakamoto, H. Sanuki, M. Sasao, A. Shimizu, Y. Shirai, S. Sudo, S. Takagi, C. Takahashi, S. Takayama, M. Takechi, K. Tanaka, K. Toi, K. Yamazaki, Y. Yoshimura and T. Watari,
Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS; Oct. 1998
(IAEA-CN-69/OV4/5)
- NIFS-569 M.M. Skoric, T. Sato, A. Maluckov, M.S. Jovanovic,
Micro- and Macro-scale Self-organization in a Dissipative Plasma; Oct. 1998
- NIFS-570 T. Hayashi, N. Mizuguchi, T-H. Watanabe, T. Sato and the Complexity Simulation Group,
Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak; Oct. 1998
(IAEA-CN-69/TH3/3)
- NIFS-571 A. Iiyoshi, A. Komori, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakaniishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, H. Yamada, S. Yamaguchi, M. Yokoyama, K.Y.Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, S. Iima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamachi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, M. Fujiwara and O. Motojima,
An Overview of the Large Helical Device Project; Oct. 1998
(IAEA-CN-69/OV1/4)
- NIFS-572 M. Fujiwara, H. Yamada, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakaniishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B. J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, S. Yamaguchi,

M. Yokoyama, K.Y. Watanabe, T. Watanabe, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, M. Iima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Otake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iiyoshi.

Plasma Confinement Studies in LHD; Oct. 1998

(IAEA-CN-69/EX2/3)

- NIFS-573 O. Motojima, K. Akaishi, H. Chikaraishi, H. Funaba, S. Hamaguchi, S. Imagawa, S. Inagaki, N. Inoue, A. Iwamoto, S. Kitagawa, A. Komon, Y. Kubota, R. Maekawa, S. Masuzaki, T. Mito, J. Miyazawa, T. Monsaki, T. Muroga, T. Nagasaka, Y. Nakamura, A. Nishimura, K. Nishimura, N. Noda, N. Ohyabu, S. Sagara, S. Sakakibara, R. Sakamoto, S. Satoh, T. Satow, M. Shoji, H. Suzuki, K. Takahata, H. Tamura, K. Watanabe, H. Yamada, S. Yamada, S. Yamaguchi, K. Yamazaki, N. Yanagi, T. Baba, H. Hayashi, M. Iima, T. Inoue, S. Kato, T. Kato, T. Kondo, S. Moriuchi, H. Ogawa, I. Otake, K. Ooba, H. Sekiguchi, N. Suzuki, S. Takami, Y. Taniguchi, T. Tsuzuki, N. Yamamoto, K. Yasui, H. Yonezu, M. Fujiwara and A. Iiyoshi,
Progress Summary of LHD Engineering Design and Construction; Oct. 1998
(IAEA-CN-69/FT2/1)
- NIFS-574 K. Toi, M. Takechi, S. Takagi, G. Matsunaga, M. Isobe, T. Kondo, M. Sasao, D.S. Darrow, K. Ohkuni, S. Ohdachi, R. Akiyama, A. Fujisawa, M. Gotoh, H. Idei, K. Ida, H. Iguchi, S. Kado, M. Kojima, S. Kubo, S. Lee, K. Matsuoka, T. Minami, S. Morita, N. Nikai, S. Nishimura, S. Okamura, M. Osakabe, A. Shimizu, Y. Shirai, C. Takahashi, K. Tanaka, T. Watari and Y. Yoshimura,
Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EXP1/19)
- NIFS-575 Y. Hamada, A. Nishizawa, Y. Kawasumi, A. Fujisawa, M. Kojima, K. Narihara, K. Ida, A. Ejin, S. Ohdachi, K. Kawahata, K. Toi, K. Sato, T. Seki, H. Iguchi, K. Adachi, S. Hidekuma, S. Hirokura, K. Iwasaki, T. Ido, R. Kumazawa, H. Kuramoto, T. Minami, I. Nomura, M. Sasao, K.N. Sato, T. Tsuzuki, I. Yamada and T. Watari,
Potential Turbulence in Tokamak Plasmas; Oct. 1998
(IAEA-CN-69/EXP2/14)
- NIFS-576 S. Murakami, U. Gasparino, H. Idei, S. Kubo, H. Maassberg, N. Marushchenko, N. Nakajima, M. Romé and M. Okamoto,
3D Simulation Study of Suprathermal Electron Transport in Non-Axisymmetric Plasmas; Oct. 1998
(IAEA-CN-69/THP1/01)
- NIFS-577 S. Fujiwara and T. Sato,
Molecular Dynamics Simulation of Structure Formation of Short Chain Molecules; Nov. 1998
- NIFS-578 T. Yamagishi,
Eigenfunctions for Vlasov Equation in Multi-species Plasmas Nov. 1998
- NIFS-579 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Strongly-Coupled Multichain Coulomb Polymers in Pure and Salt Aqueous Solutions; Nov. 1998
- NIFS-580 J. Chen, N. Nakajima and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in a Heliotron/Torsatron System: I. Mercier-unstable Equilibria; Dec. 1998
- NIFS-581 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Comparison of Multichain Coulomb Polymers in Isolated and Periodic Systems: Molecular Dynamics Study; Jan. 1999
- NIFS-582 V.S. Chan and S. Murakami,
Self-Consistent Electric Field Effect on Electron Transport of ECH Plasmas; Feb. 1999
- NIFS-583 M. Yokoyama, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani,
Roles of Bumpy Field on Collisionless Particle Confinement in Helical-Axis Heliotrons; Feb. 1999
- NIFS-584 T.-H. Watanabe, T. Hayashi, T. Sato, M. Yamada and H. Ji,
Modeling of Magnetic Island Formation in Magnetic Reconnection Experiment; Feb. 1999
- NIFS-585 R. Kumazawa, T. Mutoh, T. Seki, F. Shinpo, G. Nomura, T. Ido, T. Watanabe, Jean-Marie Noterdaeme and Yangping Zhao,
Liquid Stub Tuner for Ion Cyclotron Heating; Mar. 1999
- NIFS-586 A. Sagara, M. Iima, S. Inagaki, N. Inoue, H. Suzuki, K. Tsuzuki, S. Masuzaki, J. Miyazawa, S. Monta, Y. Nakamura, N. Noda, B. Peterson, S. Sakakibara, T. Shimozuma, H. Yamada, K. Akaishi, H. Chikaraishi, H. Funaba, O. Kaneko, K. Kawahata, A. Komon, N. Ohyabu, O. Motojima, LHD Exp. Group 1, LHD Exp. Group 2,
Wall Conditioning at the Starting Phase of LHD; Mar. 1999
- NIFS-587 T. Nakamura and T. Yabe,

Cubic Interpolated Propagation Scheme for Solving the Hyper-Dimensional Vlasov-Poisson Equation in Phase Space; Mar. 1999

- NIFS-588 W.X. Wang, N. Nakajima, S. Murakami and M. Okamoto,
An Accurate δ Method for Neoclassical Transport Calculation; Mar. 1999
- NIFS-589 K. Kishida, K. Araki, S. Kishiba and K. Suzuki,
Local or Nonlocal? Orthonormal Divergence-free Wavelet Analysis of Nonlinear Interactions in Turbulence; Mar. 1999
- NIFS-590 K. Araki, K. Suzuki, K. Kishida and S. Kishiba,
Multiresolution Approximation of the Vector Fields on T^3 ; Mar. 1999
- NIFS-591 K. Yamazaki, H. Yamada, K.Y. Watanabe, K. Nishimura, S. Yamaguchi, H. Nakanishi, A. Komori, H. Suzuki, T. Mito, H. Chikaraishi, K. Murai, O. Motojima and the LHD Group,
Overview of the Large Helical Device (LHD) Control System and Its First Operation; Apr. 1999
- NIFS-592 T. Takahashi and Y. Nakao,
Thermonuclear Reactivity of D-T Fusion Plasma with Spin-Polarized Fuel; Apr. 1999
- NIFS-593 H. Sugama,
Damping of Toroidal Ion Temperature Gradient Modes; Apr. 1999
- NIFS-594 Xiaodong Li ,
Analysis of Crowbar Action of High Voltage DC Power Supply in the LHD ICRF System; Apr. 1999
- NIFS-595 K. Nishimura, R. Horiuchi and T. Sato,
Drift-kink Instability Induced by Beam Ions in Field-reversed Configurations; Apr. 1999
- NIFS-596 Y. Suzuki, T-H. Watanabe, T. Sato and T. Hayashi,
Three-dimensional Simulation Study of Compact Toroid Plasmoid Injection into Magnetized Plasmas; Apr. 1999
- NIFS-597 H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida, S. Toda, S.-I. Itoh, M. Yagi and A. Fukuyama,
Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma; May 1999
- NIFS-598 S. Nakazawa, N. Nakajima, M. Okamoto and N. Ohyabu,
One Dimensional Simulation on Stability of Detached Plasma in a Tokamak Divertor; June 1999
- NIFS-599 S. Murakami, N. Nakajima, M. Okamoto and J. Nhrnberg,
Effect of Energetic Ion Loss on ICRF Heating Efficiency and Energy Confinement Time in Heliotrons; June 1999
- NIFS-600 R. Horiuchi and T. Sato,
Three-Dimensional Particle Simulation of Plasma Instabilities and Collisionless Reconnection in a Current Sheet; June 1999
- NIFS-601 W. Wang, M. Okamoto, N. Nakajima and S. Murakami,
Collisional Transport in a Plasma with Steep Gradients; June 1999
- NIFS-602 T. Mutoh, R. Kumazawa, T. Saki, K. Saito, F. Simpo, G. Nomura, T. Watari, X. Jikang, G. Cattanei, H. Okada, K. Ohkubo, M. Sato, S. Kubo, T. Shimozuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B.J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Narihara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohyabu, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Saito, T. Morisaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K. Ida, N. Noda, K. Yamazaki, K. Akaishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iiyoshi and LHD Exp. Group 1 and 2,
First ICRF Heating Experiment in the Large Helical Device; July 1999
- NIFS-603 P.C. de Vries, Y. Nagayama, K. Kawahata, S. Inagaki, H. Sasao and K. Nagasaki,
Polarization of Electron Cyclotron Emission Spectra in LHD; July 1999
- NIFS-604 W. Wang, N. Nakajima, M. Okamoto and S. Murakami,
δ Simulation of Ion Neoclassical Transport; July 1999