

NATIONAL INSTITUTE FOR FUSION SCIENCE

Core Localized Toroidal Alfvén Eigenmodes Destabilized By Energetic Ions in the CHS Heliotron/Torsatron

M. Takechi, G. Matsunaga, S. Takagi, K. Ohkuni, K. Toi, M. Osakabe,
M. Isobe, S. Okamura, K. Matsuoka, A. Fujisawa, H. Iguchi, S. Lee,
T. Minami, K. Tanaka, Y. Yoshimura and CHS Group

(Received - Aug. 19, 1999)

NIFS-610

Sep. 1999

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

RESEARCH REPORT NIFS Series

Core Localized Toroidal Alfvén Eigenmodes Destabilized by Energetic Ions in the CHS Heliotron/Torsatron

M. Takechi¹, G. Matsunaga¹, S. Takagi¹, K. Ohkuni¹, K. Toi, M. Osakabe,
M. Isobe, S. Okamura, K. Matsuoka, A. Fujisawa, H. Iguchi, S. Lee²,
T. Minami, K. Tanaka, Y. Yoshimura and CHS Group

National Institute for Fusion Science, Toki 509-52, Japan

¹Department of Energy Engineering Science, Nagoya Univ., Nagoya 464-01, Japan

²Naka Fusion Research Establishments, Japan Atomic Energy Research Institute, Naka-machi, Naka-gun, Ibaraki-ken, 311-01, Japan

Keywords: CHS, TAE, NBI, energetic particle, magnetic probe, soft X-ray, internal structure, continuum damping

INTRODUCTION

In a fusion reactor, it is predicted that the toroidal Alfvén eigenmodes (TAE) destabilized by the pressure gradient of energetic alpha particles with super-Alfvénic velocity, could expel the alpha particles before thermalization [1]. In large tokamaks, therefore, the excitation of TAEs has been extensively studied by the use of energetic ions generated by neutral beam injection (NBI) and ion cyclotron heating and also by the use of alpha particles in a deuterium-tritium plasma. TAE research is also important for helical systems, with moderate or high magnetic shear, that is, a heliotron/torsatron device. In the CHS heliotron/torsatron, we investigate TAE excitation by energetic ions in NBI-heated plasma [2]. In CHS plasmas, the TAE frequency, estimated from the formula $f_{\text{TAE}} = V_A / (4\pi R q)$ where V_A is the Alfvén velocity, increases rapidly towards the plasma edge because q decreases towards the edge. As a result, the frequency of a TAE in the plasma core usually intersects the Alfvén continuum in the plasma edge where the magnetic shear is high, which can cause the corresponding TAE to be stabilized due to strong continuum damping. However, it is possible for TAEs to be destabilized in this configuration if the magnetic shear in the plasma core is appreciably reduced by net plasma current I_p or finite plasma

pressure. In this situation, TAEs localized in the plasma core region may be excited by energetic ions, as observed in a tokamak [3]

EXCITATION OF TAE

In neutral-beam-heated plasmas in CHS, coherent magnetic fluctuations excited by energetic ions are detected (Fig.1), where the energetic beam-ion beta value is comparable to the bulk plasma beta ($\langle \beta_{\text{beam}} \rangle \sim 0.1\text{-}0.3\%$). The observed frequency is




Fig. 1 Typical time evolution of magnetic fluctuations related to TAEs in NBI heated plasmas in CHS, where electron density is increased linearly in time. Toroidal and poloidal mode numbers are indicated in the figure.

proportional to the Alfvén frequency. Comparison of the observed frequencies and mode numbers to a simple TAE analysis based on a large-aspect ratio tokamak clearly indicates that the magnetic fluctuations are due to TAEs. The modes were excited only when the beam velocity exceeds about half the central Alfvén velocity and the net plasma current induced by co-injected neutral beams is in the required range. Figure 2 shows dependence of the net plasma current I_p required for TAE excitation on the toroidal magnetic field. This figure suggests that the reduced magnetic shear in the plasma core is necessary for excitation of TAEs in CHS. Note that data in the plasmas having the same beta value ($\langle\beta_{\text{bulk}}\rangle \sim 0.2\%$) are plotted in Fig. 2, since the bulk plasma beta value also influences the magnetic shear in the plasma core region. The required lower bound of I_p increases with increase in B_t . The plasma current I_p in inward shifted plasmas (magnetic axis position $R_{\text{ax}}=0.92\text{m}$) is larger than that in slightly outward shifted plasmas ($R_{\text{ax}}=0.95\text{m}$). This tendency is interpreted that the rotational transform in the plasma core region of the inner shifted plasmas ($R_{\text{ax}}=0.92\text{m}$) is smaller than that of the plasmas with $R_{\text{ax}}=0.95\text{m}$. On the other hand, the upper bound of I_p is determined by elimination of the relevant TAE gap. Figure 3 shows the radial profiles of the rotational transform for the plasmas obtained at various magnetic axis positions, the bulk plasma beta values and the toroidal magnetic fields. The solid curves in Fig. 3 indicate the rotational transforms without the net plasma current where TAEs are not observed. The broken curves indicate the rotational transform calculated taking account the lowest plasma current where TAEs are observed. The horizontal solid line in Fig. 3 indicates the rotational transform $\tau=0.4$ where poloidal mode coupling between $m=2$ and $m=3$ harmonics takes place in the $n=1$ TAE. When the plasma current is in the required range, the TAE gap position (for instance, $\tau=0.4$) moves toward the plasma center and the magnetic

shear decreases there. As seen from Fig. 3, predicted profiles of the rotational transform for excitation of TAEs are very similar in the plasma core region.




Fig. 2 Dependence of the net plasma current on the toroidal magnetic field where the TAEs are detected.




Fig. 3 .Comparison of the rotational transform profiles between the cases that TAEs are observed and those without TAEs.

INTERNAL STRUCTURE OF TAEs

The internal structure of TAEs was measured with a soft X-ray detector (SX) array. The soft X-ray fluctuation level for TAEs is too low to obtain the radial profile of the fluctuation intensity because of fairly low electron temperature ($T_{eo} \leq 0.3 \text{ keV}$). For this reason, the coherence between the soft X-ray signal of each channel and the magnetic probe signal was calculated in the TAE frequency range. The solid circles in Fig. 4 (a) shows the radial profiles of coherence for the TAE observed in the plasma with $R_{ax} = 0.92 \text{ m}$. This figure suggests that the high coherence region is localized within $\rho \approx 0.6$, where the coherence of noise is ~ 0.3 . This figure shows mode localization at $\rho \approx 0.2-0.6$. Note that the high coherence near the plasma center is due to path-integral effect of soft X-ray signal. Moreover, plasma potential fluctuations due to TAEs measured by heavy ion beam probe (HIBP) are also localized in $\rho \approx 0.2-0.6$ [2]. Therefore, the TAEs are estimated to be localized in the region $\rho \approx 0.2-0.6$. In order to compare the radial profile of TAE fluctuations to other MHD activity, we shows the radial profiles of the coherence for the fishbone-like burst modes (FBs) $m=2/n=1$ observed together with TAEs [4] in Fig. 4(b), when the frequencies of TAEs and FBs are $\sim 100 \text{ kHz}$ and $\sim 30 \text{ kHz}$, respectively. FBs are destabilized by the presence of energetic ions such as TAEs. In contrast to TAEs, the fluctuations of FB are localized in more outer region $\rho \approx 0.4-0.7$. Results in Fig. 4 clearly shows that TAEs observed in CHS have a character of the core localized mode.

CONCLUSION

In CHS, core-localized TAEs are detected when the plasma current induced by co-injected NBI is in the required range depending on the toroidal magnetic field and the magnetic axis position. The parallel velocity of beam ions is also required for

excitation of the TAEs to be large than half of the central Alfvén velocity. The TAE localized in the plasma core region was supported by the radial profiles of TAE fluctuations measured by the SX-array and HIBP. The observed fluctuation frequencies are near the lower boundary of the predicted TAE gap [2]. The observed TAEs exhibit ballooning nature, as can be seen in Fig. 4(a). These results seem to be consist with the core localized TAEs. So far enhanced loss of energetic ions due to TAEs was not observed.




Fig. 4. (a) Radial profiles of the coherence of TAEs between the soft X-ray signal of each channel and the magnetic probe signal (b) Radial profiles of the coherence for fishbone-like burst mode other than TAEs. $\rho > 0$ means the outboard side of the plasma.

Detailed numerical analysis in the three-dimensional configuration is necessary to understand the TAE excitation in heliotron/torsatron configuration more clearly.

REFERENCES:

- [1] G.Y. Fu and J. W. Van Dam, Phys. Fluids B **1**, 1949 (1989).
- [2] M. Takechi, K. Toi et al., to be published in Phys. Rev. Lett. (1999).
- [3] R. Nazikian et al., Phys. Rev. Lett. **78**, 2976 (1997).
- [4] K. Toi, M. Takechi et al., in *Proceedings of the 17th International Conference on Fusion Energy*, Yokohama, Japan, 1998 (IAEA, Vienna 1999).

Recent Issues of NIFS Series

- NIFS-547 R. Kanno, N. Nakajima and M. Okamoto,
Electron Heat Transport in a Self-Similar Structure of Magnetic Islands; Apr. 1998
- NIFS-548 J.E. Rice,
Simulated Impurity Transport in LHD from MIST; May 1998
- NIFS-549 M.M. Skonc, T. Sato, A.M. Maluckov and M.S. Jovanovic,
On Kinetic Complexity in a Three-Wave Interaction; June 1998
- NIFS-550 S. Goto and S. Kida,
Passive Scalar Spectrum in Isotropic Turbulence: Prediction by the Lagrangian Direct-interaction Approximation; June 1998
- NIFS-551 T. Kuroda, H. Sugama, R. Kanno, M. Okamoto and W. Horton,
Initial Value Problem of the Toroidal Ion Temperature Gradient Mode; June 1998
- NIFS-552 T. Mutoh, R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watanai,
Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,
oronization in Future Devices -Protecting Layer against Tritium and Energetic Neutrals-; July 1998
- NIFS-554 S. Murakami and H. Saleem,
Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations; July 1998
- NIFS-555 H. Nakamura, K. Ikeda and S. Yamaguchi,
Physical Model of Nernst Element; Aug 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,
Numerical Computation of Thermoelectric and Thermomagnetic Effects; Aug 1998
- NIFS-557 Y. Takeiri, M. Osakabe, K. Tsumori, Y. Oka, O. Kaneko, E. Asano, T. Kawamoto, R. Akiyama and M. Tanaka,
Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System, Aug. 1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Structure Organization of Polyampholytes; Sep. 1998
- NIFS-559 R. Horuchi, K. Nishimura and T. Watanabe,
Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations, Sep. 1998
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Khlopchenko, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Idei, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Monta, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,
Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection; Oct 1998
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Idei, S. Kubo, S. Ohtdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,
Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron, Oct 1998
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,
Next Generation Toroidal Devices; Oct. 1998
- NIFS-563 Y. Todo and T. Sato,
Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes; Oct 1998
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watanai, T. Shimozuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Idei, Y. Oka, M. Osakabe, T. Seki, K. Tsumori, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Simpo, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Monmoto, T. Bigelow and Y.P. Zhao,

Steady State Heating Technology Development for LHD; Oct. 1998
(IAEA-CN-69/FTP/21)

- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyabu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda,
LHD-Type Compact Helical Reactors; Oct. 1998
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems; Oct. 1998
(IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D.S.Darrow, H. Idei, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tumori, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watari,
Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EX2/2)
- NIFS-568 S. Okamura, K. Matsuoka, R. Akiyama, D.S. Darrow, A. Ejiri, A. Fujisawa, M. Fujiwara, M. Goto, K. Ida, H. Idei, H. Iguchi, N. Inoue, M. Isobe, K. Itoh, S. Kado, K. Khloopenkov, T. Kondo, S. Kubo, A. Lazaros, S. Lee, G. Matsunaga, T. Minami, S. Morita, S. Murakami, N. Nakajima, N. Nikai, S. Nishimura, I. Nomura, S. Ohdachi, K. Ohkuni, M. Osakabe, R. Pavlichenko, B. Peterson, R. Sakamoto, H. Sanuki, M. Sasao, A. Shimizu, Y. Shirai, S. Sudo, S. Takagi, C. Takahashi, S. Takayama, M. Takechi, K. Tanaka, K. Toi, K. Yamazaki, Y. Yoshimura and T. Watari,
Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS; Oct. 1998
(IAEA-CN-69/OV4/5)
- NIFS-569 M.M. Skoric, T. Sato, A. Maluckov, M.S. Jovanovic,
Micro- and Macro-scale Self-organization in a Dissipative Plasma; Oct. 1998
- NIFS-570 T. Hayashi, N. Mizuguchi, T-H. Watanabe, T. Sato and the Complexity Simulation Group,
Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak; Oct. 1998
(IAEA-CN-69/TH3/3)
- NIFS-571 A. Iiyoshi, A. Komori, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, H. Yamada, S. Yamaguchi, M. Yokoyama, K.Y.Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, S. Ima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, M. Fujiwara and O. Motojima,
An Overview of the Large Helical Device Project; Oct. 1998
(IAEA-CN-69/OV1/4)
- NIFS-572 M. Fujiwara, H. Yamada, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, B. J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, S. Yamaguchi, M. Yokoyama, K.Y.Watanabe, T. Watari, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, M. Ima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Murai, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iiyoshi,
Plasma Confinement Studies in LHD; Oct. 1998
(IAEA-CN-69/EX2/3)
- NIFS-573 O. Motojima, K. Akaishi, H. Chikaraishi, H. Funaba, S. Hamaguchi, S. Imagawa, S. Inagaki, N. Inoue, A. Iwamoto, S. Kitagawa, A. Komori, Y. Kubota, R. Maekawa, S. Masuzaki, T. Mito, J. Miyazawa, T. Morisaki, T. Muroga, T. Nagasaka, Y. Nakamura, A. Nishimura, K. Nishimura, N. Noda, N. Ohyabu, S. Sagara, S. Sakakibara, R. Sakamoto, S. Satoh, T. Satow, M. Shoji, H. Suzuki, K. Takahata, H. Tamura, K. Watanabe, H. Yamada, S. Yamada, S. Yamaguchi, K. Yamazaki, N. Yanagi, T. Baba, H. Hayashi, M. Ima, T. Inoue, S. Kato, T. Kato, T. Kondo, S. Moriuchi, H. Ogawa, I. Ohtake, K. Ooba, H. Sekiguchi, N. Suzuki, S. Takami, Y. Taniguchi, T. Tsuzuki, N. Yamamoto, K. Yasui, H. Yonezu, M. Fujiwara and A. Iiyoshi,
Progress Summary of LHD Engineering Design and Construction; Oct. 1998
(IAEA-CN-69/FT2/1)
- NIFS-574 K. Toi, M. Takechi, S. Takagi, G. Matsunaga, M. Isobe, T. Kondo, M. Sasao, D.S. Darrow, K. Ohkuni, S. Ohdachi, R. Akiyama, A. Fujisawa, M. Gotoh, H. Idei, K. Ida, H. Iguchi, S. Kado, M. Kojima, S. Kubo, S. Lee, K. Matsuoka, T. Minami, S. Morita, N. Nikai, S. Nishimura, S. Okamura, M. Osakabe, A. Shimizu, Y. Shirai, C. Takahashi, K. Tanaka, T. Watari and Y. Yoshimura,
Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EXP1/19)
- NIFS-575 Y. Hamada, A. Nishizawa, Y. Kawasumi, A. Fujisawa, M. Kojima, K. Narihara, K. Ida, A. Ejiri, S. Ohdachi, K. Kawahata, K. Toi,

K Sato, T Seki, H Iguchi, K Adachi, S Hidekuma, S Hirokura, K Iwasaki, T Ido, R Kumazawa, H Kuramoto, T Minami, I Nomura, M Sasao, K N. Sato, T Tsuzuki, I. Yamada and T Watan, *Potential Turbulence in Tokamak Plasmas*, Oct 1998
(IAEA-CN-69/EXP2/14)

NIFS-576 S Murakami, U Gasparino, H. Idei, S. Kubo, H Maassberg, N Marushchenko, N Nakajima, M Romi and M Okamoto, *5D Simulation Study of Suprathermal Electron Transport in Non-Axisymmetric Plasmas*, Oct 1998
(IAEA-CN-69/THP1/01)

NIFS-577 S Fujiwara and T Sato,
Molecular Dynamics Simulation of Structure Formation of Short Chain Molecules, Nov 1998

NIFS-578 T Yamagishi,
Eigenfunctions for Vlasov Equation in Multi-species Plasmas Nov 1998

NIFS-579 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Strongly-Coupled Multichain Coulomb Polymers in Pure and Salt Aqueous Solutions, Nov 1998

NIFS-580 J. Chen, N Nakajima and M Okamoto,
Global Mode Analysis of Ideal MHD Modes in a Heliotron/Torsatron System: I. Mercier-unstable Equilibria; Dec. 1998

NIFS-581 M Tanaka, A Yu Grosberg and T. Tanaka,
Comparison of Multichain Coulomb Polymers in Isolated and Periodic Systems: Molecular Dynamics Study; Jan 1999

NIFS-582 V S. Chan and S. Murakami,
Self-Consistent Electric Field Effect on Electron Transport of ECH Plasmas, Feb 1999

NIFS-583 M. Yokoyama, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani,
Roles of Bumpy Field on Collisionless Particle Confinement in Helical-Axis Heliotrons; Feb 1999

NIFS-584 T.-H. Watanabe, T. Hayashi, T. Sato, M. Yamada and H. Ji,
Modeling of Magnetic Island Formation in Magnetic Reconnection Experiment; Feb. 1999

NIFS-585 R. Kumazawa, T. Mutoh, T. Seki, F. Shinpo, G. Nomura, T. Ido, T. Watan, Jean-Marie Noterdaeme and Yangping Zhao,
Liquid Stub Tuner for Ion Cyclotron Heating; Mar 1999

NIFS-586 A Sagara, M. Iima, S. Inagaki, N. Inoue, H. Suzuki, K. Tsuzuki, S. Masuzaki, J. Miyazawa, S. Morita, Y. Nakamura, N. Noda, B. Peterson, S. Sakakibara, T. Shimozuma, H. Yamada, K. Akaishi, H. Chikaraishi, H. Funaba, O. Kaneko, K. Kawahata, A. Komori, N. Ohyabu, O. Motojima, LHD Exp. Group 1, LHD Exp. Group 2,
Wall Conditioning at the Starting Phase of LHD, Mar. 1999

NIFS-587 T Nakamura and T Yabe,
Cubic Interpolated Propagation Scheme for Solving the Hyper-Dimensional Vlasov-Poisson Equation in Phase Space, Mar. 1999

NIFS-588 W.X. Wang, N Nakajima, S Murakami and M Okamoto,
An Accurate δ Method for Neoclassical Transport Calculation, Mar 1999

NIFS-589 K. Kishida, K. Araki, S. Kishiba and K. Suzuki,
Local or Nonlocal? Orthonormal Divergence-free Wavelet Analysis of Nonlinear Interactions in Turbulence; Mar. 1999

NIFS-590 K. Araki, K. Suzuki, K. Kishida and S. Kishiba,
Multiresolution Approximation of the Vector Fields on T^3 , Mar 1999

NIFS-591 K Yamazaki, H. Yamada, K Y.Watanabe, K Nishimura, S. Yamaguchi, H. Nakanishi, A. Komori, H. Suzuki, T. Mito, H. Chikaraishi, K. Murai, O. Motojima and the LHD Group,
Overview of the Large Helical Device (LHD) Control System and Its First Operation; Apr 1999

NIFS-592 T Takahashi and Y Nakao,
Thermonuclear Reactivity of D-T Fusion Plasma with Spin-Polarized Fuel, Apr 1999

NIFS-593 H Sugama,

Damping of Toroidal Ion Temperature Gradient Modes; Apr. 1999

- NIFS-594 Xiaodong Li ,
Analysis of Crowbar Action of High Voltage DC Power Supply in the LHD ICRF System; Apr. 1999
- NIFS-595 K. Nishimura, R. Horiuchi and T. Sato,
Drift-kink Instability Induced by Beam Ions in Field-reversed Configurations; Apr. 1999
- NIFS-596 Y. Suzuki, T-H Watanabe, T. Sato and T. Hayashi,
Three-dimensional Simulation Study of Compact Toroid Plasmoid Injection into Magnetized Plasmas; Apr. 1999
- NIFS-597 H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida, S. Toda, S.-I. Itoh, M. Yagi and A. Fukuyama,
Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma; May 1999
- NIFS-598 S. Nakazawa, N. Nakajima, M. Okamoto and N. Ohyabu,
One Dimensional Simulation on Stability of Detached Plasma in a Tokamak Divertor; June 1999
- NIFS-599 S. Murakami, N. Nakajima, M. Okamoto and J. Nhrenberg,
Effect of Energetic Ion Loss on ICRF Heating Efficiency and Energy Confinement Time in Heliotrons; June 1999
- NIFS-600 R. Horiuchi and T. Sato,
Three-Dimensional Particle Simulation of Plasma Instabilities and Collisionless Reconnection in a Current Sheet; June 1999
- NIFS-601 W. Wang, M. Okamoto, N. Nakajima and S. Murakami,
Collisional Transport in a Plasma with Steep Gradients; June 1999
- NIFS-602 T. Mutoh, R. Kumazawa, T. Saki, K. Saito, F. Simpo, G. Nomura, T. Watan, X. Jikang, G. Cattanei, H. Okada, K. Ohkubo, M. Sato, S. Kubo, T. Shimozuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B.J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Nanbara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohyabu, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Sato, T. Morisaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K.. Ida, N. Noda, K. Yamazaki, K. Akaishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iiyoshi and LHD Exp. Group 1 and 2,
First ICRF Heating Experiment in the Large Helical Device; July 1999
- NIFS-603 P.C. de Vries, Y. Nagayama, K. Kawahata, S. Inagaki, H. Sasao and K. Nagasaki,
Polarization of Electron Cyclotron Emission Spectra in LHD; July 1999
- NIFS-604 W. Wang, N. Nakajima, M. Okamoto and S. Murakami,
 δf Simulation of Ion Neoclassical Transport; July 1999
- NIFS-605 T. Hayashi, N. Mizuguchi, T. Sato and the Complexity Simulation Group,
Numerical Simulation of Internal Reconnection Event in Spherical Tokamak; July 1999
- NIFS-606 M. Okamoto, N. Nakajima and W. Wang,
On the Two Weighting Scheme for δf Collisional Transport Simulation; Aug. 1999
- NIFS-607 O. Motojima, A.A. Shishkin, S. Inagaki, K. Y. Watanabe,
Possible Control Scenario of Radial Electric Field by Loss-Cone-Particle Injection into Helical Device; Aug. 1999
- NIFS-608 R. Tanaka, T. Nakamura and T. Yabe,
Constructing Exactly Conservative Scheme in Non-conservative Form; Aug. 1999
- NIFS-609 H. Sugama,
Gyrokinetic Field Theory; Aug. 1999
- NIFS-610 M. Takechi, G. Matsunaga, S. Takagi, K. Ohkuni, K. Toi, M. Osakabe, M. Isobe, S. Okamura, K. Matsuoka, A. Fujisawa, H. Iguchi, S. Lee, T. Minami, K. Tanaka, Y. Yoshimura and CHS Group,
Core Localized Toroidal Alfvén Eigenmodes Destabilized By Energetic Ions in the CHS Heliotron/Torsatron; Sep. 1999