

NATIONAL INSTITUTE FOR FUSION SCIENCE

Analysis of Current Diffusive Ballooning Mode in Tokamaks

M. Uchida, A. Fukuyama, K. Itoh, S.-I. Itoh and M. Yagi

(Received - Nov. 19, 1999)

NIFS-615

Dec. 1999

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

RESEARCH REPORT NIFS Series

NAGOYA, JAPAN

Analysis of Current Diffusive Ballooning Mode in Tokamaks

M. Uchida*, A. Fukuyama*, K. Itoh**, S.-I. Itoh*** and M. Yagi***

*Department of Nuclear Engineering, Kyoto University.

**National Institute for Fusion Science,

***Research Institute for Applied Mechanics, Kyushu University

Abstract

The effect of finite gyroradius on the current diffusive ballooning mode is examined. Starting from the reduced MHD equations including turbulent transports, coupling with drift motion and finite gyroradius effect of ions, we derive a ballooning mode equation with complex transport coefficients. The eigenfrequency, saturation level and thermal diffusivity are evaluated numerically from the marginal stability condition. Preliminary results of their parameter dependence is presented.

Keywords: ballooning mode, current diffusivity, turbulent transport, finite gyroradius effect

1 Introduction

The self-sustained turbulence model [1] of the current diffusive ballooning mode (CDBM) is one of the most promising candidates to explain the particle and energy transport in tokamaks. This transport model has successfully described the L mode transport and improved confinement associated with the formation of the internal transport barriers [2]. Since the previous analysis [3] based on the reduced MHD equations [4] neglected the finite ion gyroradius effect, we could not separate the transport coefficients of ions and electrons. In this paper, we describe the formulation and the results of numerical analysis of the ballooning mode equation which includes the coupling with the drift motion and the finite gyro radius effect.

2 Ballooning mode equation

We start from a set of reduced MHD equations, i.e. the equation of perpendicular motion of ions, the extended Ohm's law, the energy equations of electrons and ions [5],

$$\left(\frac{\partial}{\partial t} + i\omega_{*i} - \mu_i \nabla_{\perp}^2 \right) \nabla_{\perp}^2 \tilde{\phi} = \nabla_{\parallel} \tilde{J}_{\parallel} + (\mathbf{b} \times \boldsymbol{\kappa}) \cdot \nabla (\tilde{p}_i + \tilde{p}_e) \quad (1)$$

$$\frac{\partial}{\partial t} \tilde{A} - \nabla_{\parallel} \tilde{\phi} = -\frac{1}{\omega_{ci} \tau_A} \nabla_{\parallel} \tilde{p}_e - \lambda \nabla_{\perp}^2 \tilde{J}_{\parallel} \quad (2)$$

$$\frac{\partial}{\partial t} \tilde{p}_e - \chi_e \nabla_{\perp}^2 \tilde{p}_e + \mathbf{b} \cdot (\nabla_{\perp} \tilde{\phi} \times \nabla_{\perp} p_{e0}) = 0 \quad (3)$$

$$\frac{\partial}{\partial t} \tilde{p}_i - \chi_i \nabla_{\perp}^2 \tilde{p}_i + \mathbf{b} \cdot (\nabla_{\perp} \tilde{\phi} \times \nabla_{\perp} p_{i0}) = 0 \quad (4)$$

$$\nabla_{\perp}^2 \tilde{A} = \tilde{J}_{\parallel} \quad (5)$$

where $\tau_A = qR/v_A$, μ_i the ion viscosity, χ_i and χ_e the ion and electron thermal diffusivities, λ the current diffusivity due to the electron viscosity μ_e and other notations are standard. The streaming function for ions $\tilde{\phi}$ is reduced from the electrostatic potential ϕ by a factor of $\Lambda_0(b) = I_0(b)e^{-b}$ where I_0 is the modified Bessel function of the first kind and $b = k_{\perp}^2 \rho_i^2$. Length and time are normalized by the minor radius r and the poloidal Alfvén time τ_A , respectively. Using the ballooning transformation [6], we obtain

$$\begin{aligned} \frac{d}{d\eta} \frac{f}{-i\omega + \lambda n^4 q^4 f^2} \frac{d}{d\eta} \left(1 - \frac{-i\omega_{*e}}{-i\omega + \chi_e n^2 q^2 f} \right) \tilde{\phi} \\ + \left(\frac{\alpha_e}{-i\omega + \chi_e n^2 q^2 f} + \frac{\alpha_i \Lambda_0}{-i\omega + \chi_i n^2 q^2 f} \right) H(\eta) \tilde{\phi} \\ + (i\omega - i\omega_{*i} - \mu_i n^2 q^2 f) f \Lambda_0 \tilde{\phi} = 0 \end{aligned} \quad (6)$$

where $\alpha = -q^2 R (d\beta/dr)$ is the normalized pressure gradient, $s = (r/q)(dq/dr)$ the magnetic shear and η the ballooning coordinates. Taking account of the Shafranov shift of the magnetic surface, we take $f = 1 + (s\eta - \alpha \sin \eta)^2$ and $H(\eta) = \kappa + \cos \eta + (s\eta - \alpha \sin \eta) \sin \eta$ where $\kappa = (r/R)(1 - 1/q^2)$ is the average magnetic curvature. The first term in eq.(6) has

a stabilizing effect due to the bending of the magnetic field and is reduced by the current diffusivity λ . The second term coming from the pressure gradient destabilizes the mode in the bad curvature region and is reduced by the thermal diffusivity. The last term represents the stabilizing effect of ω_{*i} and the ion viscosity.

The transport coefficients are determined by the relation for the nonlinear transfer rate γ , e.g. $\chi_e = \langle \tilde{\phi}^2 \rangle / \gamma$ and $\gamma = -i\omega + k_\perp^2 \chi_e$, where the bracket $\langle \rangle$ denotes the average over the wave number and the η dependence of $\langle \tilde{\phi}^2 \rangle$ is neglected in the present analysis [1]. Since mode frequency has a real part, the transport coefficients are written as

$$\mu_e = \xi \lambda = \sqrt{\frac{\langle \tilde{\phi}^2 \rangle}{n^2 q^2} - \frac{1}{4} \left\{ \frac{\omega(\xi + n^2 q^2)}{n^4 q^4 f^2} \right\}^2 + \frac{i}{2} \frac{\omega(\xi + n^2 q^2)}{n^4 q^4 f^2}} \quad (7)$$

$$\mu_i = \sqrt{\frac{\Lambda_0^2 \langle \tilde{\phi}^2 \rangle}{n^2 q^2} - \frac{1}{4} \left(\frac{\omega - \omega_{*i}}{n^2 q^2 f} \right)^2 + \frac{i}{2} \frac{\omega - \omega_{*i}}{n^2 q^2 f}} \quad (8)$$

$$\chi_e = \sqrt{\frac{\langle \tilde{\phi}^2 \rangle}{n^2 q^2} - \frac{1}{4} \left(\frac{\omega}{n^2 q^2 f} \right)^2 + \frac{i}{2} \frac{\omega}{n^2 q^2 f}} \quad (9)$$

$$\chi_i = \sqrt{\frac{\Lambda_0^2 \langle \tilde{\phi}^2 \rangle}{n^2 q^2} - \frac{1}{4} \left(\frac{\omega}{n^2 q^2 f} \right)^2 + \frac{i}{2} \frac{\omega}{n^2 q^2 f}} \quad (10)$$

where n is the toroidal mode number and $\xi = \omega_p^2 r^2 / c^2$. When $\langle \tilde{\phi}^2 \rangle$ is smaller than critical values, we take that the square roots have negative imaginary values so that the turbulent transport coefficients vanish for $\langle \tilde{\phi}^2 \rangle = 0$. A pure imaginary value for small $\langle \tilde{\phi}^2 \rangle$ implies nonlinear frequency shift rather than diffusion. We should note that the electron thermal diffusivity χ_e is less than χ_i by a factor of Λ_0 for large $\langle \tilde{\phi}^2 \rangle$. With the increase of the perpendicular wave number $k_\perp \sim nq$, the finite gyroradius effect reduces χ_i .

With the marginal stability condition ($\text{Im } \omega = 0$), we numerically solve the ballooning mode equation (6) to find the eigen modes and evaluate the real part of the mode frequency ω_r and the average amplitude of the fluctuation $\langle \tilde{\phi}^2 \rangle$. Figure 1 shows an example of the CDBM eigenmode for $s = 0.4$, $\alpha = 0.4$, $q = 20$, $\rho_i/r = 0.01$ and $nq = 134$. It has a small negative real frequency ($\omega_r = -0.05$) and propagates in the direction of ion drift motion. The saturation level of the fluctuation is $\langle \tilde{\phi}^2 \rangle = 6 \times 10^{-5}$ and this mode is unstable for smaller $\langle \tilde{\phi}^2 \rangle$. The eigen mode structure indicates that the dominant real part localizes in the outer region ($\eta = 0$), while the out-of-phase imaginary part in the inner region ($\eta = \pm\pi$).

The saturation level of the average fluctuation amplitude depends on the mode number nq . Figure 2 illustrates the nq dependence of the electron thermal diffusivity χ_e derived from the saturation fluctuation amplitude $\langle \tilde{\phi}^2 \rangle$. The maximum value of χ_e near $nq = 134$ gives the thermal diffusivity necessary for stabilizing the ballooning modes for all nq .

3 Parameter dependence

In order to study the parameter dependence of the mode frequency and the thermal diffusivity, we look for the maximum value of χ_e by numerically solving the ballooning mode equation (6) for wide range of nq and for various values of s and α . We have been much benefitted by parallel processing on a PC cluster of 8 CPUs with a numerical code using the Message Passing Interface (MPI) library.

The (s, α) dependence of the electron thermal diffusivity is shown in Figs. 3 and 4 for $\xi = 10^4$ and $q = 10^3$. Figure 3 depicts the case of small gyroradius, $\rho_i/r = 10^{-3}$, while Fig. 4 the case of larger gyroradius, $\rho_i/r = 10^{-2}$. The (s, α) dependence in Fig. 3 is almost same as that of $\rho_i/r = 0$ [2]. For larger ρ_i , χ_e is enhanced, typically, by a factor of 3 and the shear dependence is slightly reduced.

For small α ($\alpha < 0.4$), χ_e is a strongly increasing function of α ($\chi_e \propto \alpha^{3/2}$) and decreases for small and negative values of s . For larger value of α ($\alpha > 0.4$), χ_e is enhanced in the stronger shear region ($s > 0.5$) where the ideal ballooning mode is unstable. The reduction of χ_e in the weak and negative shear region is more prominent for larger α (larger pressure gradient). These behavior is similar to the result of the previous analyses without coupling to the drift motion, $\omega_{*i} = \omega_{*e} = 0$. One of the reasons is the large q value in the present calculation ($q = 10^3$). For realistic values of q , the effect of finite ω_{*i} and ω_{*e} affects the mode structure, but conclusive results have not been obtained yet.

The mode frequency is usually negative except in the region, $\chi_e > 10^{-4}$, where the ideal mode is unstable. The mode number nq decreases with the increase of χ_e . Typical value of nq is 300 for $\chi_e = 10^{-5}$ and 100 for $\chi_e = 10^{-4}$. As the mode number increases, the finite gyroradius effect enhances χ_e and reduces χ_i . Figure 4 indicates the contour of the ratio χ_i/χ_e on the (s, α) plane. In the region where $\alpha > 0.4$ and $s > 0.5$, χ_i approaches to χ_e , otherwise χ_i/χ_e is less than about 0.4.

4 Conclusion

Taking account of the gyroradius effects, we found that χ_e of the CDBM transport model is enhanced by a factor of 3 for $\rho/r = 10^{-2}$ and that the ratio χ_i/χ_e is considerably reduced for low α (L-mode) region. In the present calculation, the safety factor q is limited to a rather large value. The calculation with realistic values of q as well as the effect of the $E \times B$ rotation shear will be reported in future.

This work is partly supported by the Grant-in-Aid for Scientific Research of the Ministry of Education, Science, Sports and Culture Japan, by the collaboration programme of the Advanced Fusion Research Center of Kyushu University, and by the Collaboration programme of National Institute for Fusion Science.

References

- [1] K. Itoh, et al., *Plasma Phys. Control. Fusion.* **36**, 279 (1994).
- [2] A. Fukuyama, et al., *Plasma Phys. Control. Fusion.* **37**, 611 (1996).
- [3] M. Yagi, et al., *Phys. Fluids. B* **5**, 3702 (1993).
- [4] H. Strauss, *Phys. Fluids* **20**, 1354 (1977)
- [5] A. Fukuyama, et al., 17th IAEA Conf. on Fusion Energy IAEA-CN-69/TH1/6 (1998).
- [6] J. W. Connor and J. B. Taylor, *Nucl. Fusion* **17**, 1047 (1977).
- [7] K. Itoh, et al., *Plasma Phys. Control. Fusion.* **36**, 279 (1994)

Figure captions

- Fig. 1** Spatial structure of the eigen mode for the parameters $s = 0.4$, $\alpha = 0.4$, $q = 20$, $\rho/r = 0.01$ and $nq = 134$.
- Fig. 2** nq dependence of the thermal diffusivity χ_e derived from the marginal stability condition for the parameters $s = 0.4$, $\alpha = 0.4$, $q = 20$ and $\rho/r = 0.01$.
- Fig. 3** Contour of the electron thermal diffusivity χ_e on the $(s - \alpha)$ plane for $\rho/r = 10^{-3}$.
- Fig. 4** Contour of the electron thermal diffusivity χ_e on the $(s - \alpha)$ plane for $\rho/r = 0.01$.
- Fig. 5** Contour of the ratio χ_i/χ_e on the $(s - \alpha)$ plane for $\rho/r = 0.01$.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

Recent Issues of NIFS Series

- NIFS-552 T. Mutoh R. Kumazawa, T. Seki, F. Simpo, G. Nomura, T. Ido and T. Watan,
Steady State Tests of High Voltage Ceramic Feedthroughs and Co-Axial Transmission Line of ICRF Heating System for the Large Helical Device; June 1998
- NIFS-553 N. Noda, K. Tsuzuki, A. Sagara, N. Inoue, T. Muroga,
oronaization in Future Devices -Protecting Layer against Tritium and Energetic Neurals-; July 1998
- NIFS-554 S. Murakami and H. Saleem,
Electromagnetic Effects on Rippling Instability and Tokamak Edge Fluctuations, July 1998
- NIFS-555 H. Nakamura, K. Ikeda and S. Yamaguchi,
Physical Model of Nernst Element, Aug 1998
- NIFS-556 H. Okumura, S. Yamaguchi, H. Nakamura, K. Ikeda and K. Sawada,
Numerical Computation of Thermolectric and Thermomagnetic Effects, Aug. 1998
- NIFS-557 Y. Takeiri, M. Osakabe, K. Tsumori, Y. Oka, O. Kaneko, E. Asano, T. Kawamoto, R. Akiyama and M. Tanaka,
Development of a High-Current Hydrogen-Negative Ion Source for LHD-NBI System; Aug.1998
- NIFS-558 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Structure Organization of Polyampholytes; Sep 1998
- NIFS-559 R. Honuchi, K. Nishimura and T. Watanabe,
Kinetic Stabilization of Tilt Disruption in Field-Reversed Configurations, Sep 1998
(IAEA-CN-69/THP1/11)
- NIFS-560 S. Sudo, K. Khlopchenkov, K. Matsuoka, S. Okamura, C. Takahashi, R. Akiyama, A. Fujisawa, K. Ida, H. Ider, H. Iguchi, M. Isobe, S. Kado, K. Kondo, S. Kubo, H. Kuramoto, T. Minami, S. Morita, S. Nishimura, M. Osakabe, M. Sasao, B. Peterson, K. Tanaka, K. Toi and Y. Yoshimura,
Particle Transport Study with Tracer-Encapsulated Solid Pellet Injection. Oct 1998
(IAEA-CN-69/EXP1/18)
- NIFS-561 A. Fujisawa, H. Iguchi, S. Lee, K. Tanaka, T. Minami, Y. Yoshimura, M. Osakabe, K. Matsuoka, S. Okamura, H. Ider, S. Kubo, S. Ohdachi, S. Morita, R. Akiyama, K. Toi, H. Sanuki, K. Itoh, K. Ida, A. Shimizu, S. Takagi, C. Takahashi, M. Kojima, S. Hidekuma, S. Nishimura, M. Isobe, A. Ejiri, N. Inoue, R. Sakamoto, Y. Hamada and M. Fujiwara,
Dynamic Behavior Associated with Electric Field Transitions in CHS Heliotron/Torsatron, Oct 1998
(IAEA-CN-69/EX5/1)
- NIFS-562 S. Yoshikawa,
Next Generation Toroidal Devices; Oct. 1998
- NIFS-563 Y. Todo and T. Sato,
Kinetic-Magnetohydrodynamic Simulation Study of Fast Ions and Toroidal Alfvén Eigenmodes; Oct 1998
(IAEA-CN-69/THP2/22)
- NIFS-564 T. Watari, T. Shimozuma, Y. Takeiri, R. Kumazawa, T. Mutoh, M. Sato, O. Kaneko, K. Ohkubo, S. Kubo, H. Ider, Y. Oka, M. Osakabe, T. Seki, K. Tsumori, Y. Yoshimura, R. Akiyama, T. Kawamoto, S. Kobayashi, F. Shimpou, Y. Takita, E. Asano, S. Itoh, G. Nomura, T. Ido, M. Hamabe, M. Fujiwara, A. Iiyoshi, S. Monmoto, T. Bigelow and Y P Zhao,
Steady State Heating Technology Development for LHD, Oct. 1998
(IAEA-CN-69/FTP/21)
- NIFS-565 A. Sagara, K.Y. Watanabe, K. Yamazaki, O. Motojima, M. Fujiwara, O. Mitarai, S. Imagawa, H. Yamanishi, H. Chikaraishi, A. Kohyama, H. Matsui, T. Muroga, T. Noda, N. Ohyabu, T. Satow, A.A. Shishkin, S. Tanaka, T. Terai and T. Uda,
LHD-Type Compact Helical Reactors; Oct. 1998
(IAEA-CN-69/FTP/03(R))
- NIFS-566 N. Nakajima, J. Chen, K. Ichiguchi and M. Okamoto,
Global Mode Analysis of Ideal MHD Modes in L=2 Heliotron/Torsatron Systems, Oct. 1998
(IAEA-CN-69/THP1/08)
- NIFS-567 K. Ida, M. Osakabe, K. Tanaka, T. Minami, S. Nishimura, S. Okamura, A. Fujisawa, Y. Yoshimura, S. Kubo, R. Akiyama, D.S.Darrow, H. Ider, H. Iguchi, M. Isobe, S. Kado, T. Kondo, S. Lee, K. Matsuoka, S. Morita, I. Nomura, S. Ohdachi, M. Sasao, A. Shimizu, K. Tsumori, S. Takayama, M. Takechi, S. Takagi, C. Takahashi, K. Toi and T. Watan,
Transition from L Mode to High Ion Temperature Mode in CHS Heliotron/Torsatron Plasmas, Oct 1998
(IAEA-CN-69/EX2/2)

- NIFS-568 S. Okamura, K. Matsuoka, R. Akiyama, D.S. Darrow, A. Ejiri, A. Fujisawa, M. Fujiwara, M. Goto, K. Ida H. Iide, H. Iguchi, N. Inoue, M. Isobe, K. Itoh, S. Kado, K. Khlopchenov, T. Kondo, S. Kubo, A. Lazaros, S. Lee, G. Matsunaga, T. Minami, S. Morita, S. Murakami, N. Nakajima, N. Nikai, S. Nishimura, I. Nomura, S. Ohdachi, K. Ohkuni, M. Osakabe, R. Pavlichenko, B. Peterson, R. Sakamoto, H. Sanuki, M. Sasao, A. Shimizu, Y. Shirai, S. Sudo, S. Takagi, C. Takahashi, S. Takayama, M. Takechi, K. Tanaka, K. Toi, K. Yamazaki, Y. Yoshimura and T. Watan,
Confinement Physics Study in a Small Low-Aspect-Ratio Helical Device CHS; Oct. 1998
(IAEA-CN-69/OV4/5)
- NIFS-569 M.M. Skoric, T. Sato, A. Maluckov, M.S. Jovanovic,
Micro- and Macro-scale Self-organization in a Dissipative Plasma; Oct. 1998
- NIFS-570 T. Hayashi, N. Mizuguchi, T-H. Watanabe, T. Sato and the Complexity Simulation Group,
Nonlinear Simulations of Internal Reconnection Event in Spherical Tokamak; Oct. 1998
(IAEA-CN-69/TH3/3)
- NIFS-571 A. Iiyoshi, A. Komori, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Iide, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Monta, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyanbu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, H. Yamada, S. Yamaguchi, M. Yokoyama, K.Y.Watanabe, T. Watanabe, R. Akiyama, H. Chikaraishi, K. Haba, S. Hamaguchi, S. Iima, S. Imagawa, N. Inoue, K. Iwamoto, S. Kitagawa, Y. Kubota, J. Kodaira, R. Maekawa, T. Mito, T. Nagasaka, A. Nishimura, Y. Takita, C. Takahashi, K. Takahata, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Mura, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, M. Fujiwara and O. Motojima,
An Overview of the Large Helical Device Project; Oct. 1998
(IAEA-CN-69/OV1/4)
- NIFS-572 M. Fujiwara, H. Yamada, A. Ejiri, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Iide, S. Inagaki, S. Kado, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Monta, S. Murakami, S. Muto, T. Muto, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Narihara, K. Nishimura, N. Noda, T. Kobuchi, S. Ohdachi, N. Ohyanbu, Y. Oka, M. Osakabe, T. Ozaki, B.J. Peterson, A. Sagara, S. Sakakibara, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, H. Suzuki, Y. Takeiri, K. Tanaka, K. Toi, T. Tokuzawa, K. Tsumori, I. Yamada, S. Yamaguchi, M. Yokoyama, K. Yamauchi, H. Tamura, T. Tsuzuki, S. Yamada, N. Yanagi, H. Yonezu, Y. Hamada, K. Matsuoka, K. Mura, K. Ohkubo, I. Ohtake, M. Okamoto, S. Sato, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima and A. Iiyoshi,
Plasma Confinement Studies in LHD; Oct. 1998
(IAEA-CN-69/EX2/3)
- NIFS-573 O. Motojima, K. Akaishi, H. Chikaraishi, H. Funaba, S. Hamaguchi, S. Imagawa, S. Inagaki, N. Inoue, A. Iwamoto, S. Kitagawa, A. Komori, Y. Kubota, R. Maekawa, S. Masuzaki, T. Mito, J. Miyazawa, T. Morisaki, T. Muroga, T. Nagasaka, Y. Nakamura, A. Nishimura, K. Nishimura, N. Noda, N. Ohyanbu, S. Sagara, S. Sakakibara, R. Sakamoto, S. Satoh, T. Satow, M. Shoji, H. Suzuki, K. Takahata, H. Tamura, K. Watanabe, H. Yamada, S. Yamada, S. Yamaguchi, K. Yamazaki, N. Yanagi, T. Baba, H. Hayashi, M. Iima, T. Inoue, S. Kato, T. Kato, T. Kondo, S. Monuchi, H. Ogawa, I. Ohtake, K. Ooba, H. Sekiguchi, N. Suzuki, S. Takami, Y. Taniguchi, T. Tsuzuki, N. Yamamoto, K. Yasui, H. Yonezu, M. Fujiwara and A. Iiyoshi,
Progress Summary of LHD Engineering Design and Construction; Oct. 1998
(IAEA-CN-69/FT2/1)
- NIFS-574 K. Toi, M. Takechi, S. Takagi, G. Matsunaga, M. Isobe, T. Kondo, M. Sasao, D.S. Darrow, K. Ohkuni, S. Ohdachi, R. Akiyama, A. Fujisawa, M. Gotoh, H. Iide, K. Ida, H. Iguchi, S. Kado, M. Kojima, S. Kubo, S. Lee, K. Matsuoka, T. Minami, S. Morita, N. Nikai, S. Nishimura, S. Okamura, M. Osakabe, A. Shimizu, Y. Shirai, C. Takahashi, K. Tanaka, T. Watari and Y. Yoshimura,
Global MHD Modes Excited by Energetic Ions in Heliotron/Torsatron Plasmas; Oct. 1998
(IAEA-CN-69/EXP1/19)
- NIFS-575 Y. Hamada, A. Nishizawa, Y. Kawasumi, A. Fujisawa, M. Kojima, K. Narihara, K. Ida, A. Ejiri, S. Ohdachi, K. Kawahata, K. Toi, K. Sato, T. Seki, H. Iguchi, K. Adachi, S. Hidekuma, S. Hirokura, K. Iwasaki, T. Ido, R. Kumazawa, H. Kuramoto, T. Minami, I. Nomura, M. Sasao, K.N. Sato, T. Tsuzuki, I. Yamada and T. Watan,
Potential Turbulence in Tokamak Plasmas; Oct. 1998
(IAEA-CN-69/EXP2/14)
- NIFS-576 S. Murakami, U. Gasparino, H. Ider, S. Kubo, H. Maassberg, N. Marushchenko, N. Nakajima, M. Romé and M. Okamoto,
5D Simulation Study of Suprathermal Electron Transport in Non-Axisymmetric Plasmas; Oct. 1998
(IAEA-CN-69/THP1/01)
- NIFS-577 S. Fujiwara and T. Sato,
Molecular Dynamics Simulation of Structure Formation of Short Chain Molecules; Nov. 1998
- NIFS-578 T. Yamagishi,
Eigenfunctions for Vlasov Equation in Multi-species Plasmas Nov. 1998
- NIFS-579 M. Tanaka, A. Yu Grosberg and T. Tanaka,
Molecular Dynamics of Strongly-Coupled Multichain Coulomb Polymers in Pure and Salt Aqueous

Solutions, Nov. 1998

- NIFS-580 J Chen, N Nakajima and M Okamoto,
Global Mode Analysis of Ideal MHD Modes in a Heliotron/Torsatron System I. Mercier-unstable Equilibria. Dec 1998
- NIFS-581 M Tanaka, A Yu Grosberg and T Tanaka,
Comparison of Multichain Coulomb Polymers in Isolated and Periodic Systems. Molecular Dynamics Study, Jan. 1999
- NIFS-582 V S Chan and S Murakami,
Self-Consistent Electric Field Effect on Electron Transport of ECH Plasmas, Feb 1999
- NIFS-583 M Yokoyama, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani,
Roles of Bumpy Field on Collisionless Particle Confinement in Helical-Axis Heliotrons, Feb 1999
- NIFS-584 T H Watanabe, T. Hayashi, T. Sato, M Yamada and H. Ji,
Modeling of Magnetic Island Formation in Magnetic Reconnection Experiment, Feb 1999
- NIFS-585 R Kumazawa, T Mutoh, T Seki, F Shinpo, G Nomura, T Ido, T Watanabe, Jean-Marie Noterdaeme and Yangping Zhao,
Liquid Stub Tuner for Ion Cyclotron Heating, Mar 1999
- NIFS-586 A Sagara, M Iima, S Inagaki, N. Inoue, H Suzuki, K Tsuzuki, S Masuzaki, J Miyazawa, S Morita, Y Nakamura, N Noda, B Peterson, S Sakakibara, T Shimozuma, H Yamada, K Akaishi, H Chikaraishi, H Funaba, O Kaneko, K Kawahata, A Komori, N Ohyabu, O Motojima, LHD Exp Group 1, LHD Exp Group 2,
Wall Conditioning at the Starting Phase of LHD, Mar 1999
- NIFS-587 T Nakamura and T Yabe,
Cubic Interpolated Propagation Scheme for Solving the Hyper-Dimensional Vlasov-Poisson Equation in Phase Space, Mar 1999
- NIFS-588 W.X Wang, N. Nakajima, S Murakami and M Okamoto,
An Accurate Method for Neoclassical Transport Calculation, Mar 1999
- NIFS-589 K Kishida, K Araki, S Kishiba and K Suzuki,
Local or Nonlocal? Orthonormal Divergence-free Wavelet Analysis of Nonlinear Interactions in Turbulence, Mar. 1999
- NIFS-590 K Araki, K Suzuki, K. Kishida and S. Kishiba,
Multiresolution Approximation of the Vector Fields on T^3 , Mar 1999
- NIFS-591 K Yamazaki, H Yamada, K.Y Watanabe, K. Nishimura, S Yamaguchi, H Nakanishi, A Komori, H Suzuki, T Mito, H Chikaraishi, K. Murai, O. Motojima and the LHD Group,
Overview of the Large Helical Device (LHD) Control System and Its First Operation, Apr 1999
- NIFS-592 T. Takahashi and Y. Nakao,
Thermonuclear Reactivity of D-T Fusion Plasma with Spin-Polarized Fuel, Apr 1999
- NIFS-593 H. Sugama,
Damping of Toroidal Ion Temperature Gradient Modes, Apr 1999
- NIFS-594 Xiaodong Li ,
Analysis of Crowbar Action of High Voltage DC Power Supply in the LHD ICRF System, Apr 1999
- NIFS-595 K Nishimura, R Honuchi and T. Sato,
Drift-kink Instability Induced by Beam Ions in Field-reversed Configurations, Apr 1999
- NIFS-596 Y Suzuki, T-H Watanabe, T Sato and T Hayashi,
Three-dimensional Simulation Study of Compact Toroid Plasmoid Injection into Magnetized Plasmas, Apr. 1999
- NIFS-597 H. Sanuki, K Itoh, M Yokoyama, A Fujisawa, K Ida, S Toda, S-I Itoh, M Yagi and A Fukuyama
Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma, May 1999

- NIFS-598 S. Nakazawa, N. Nakajima, M. Okamoto and N. Ohyabu,
One Dimensional Simulation on Stability of Detached Plasma in a Tokamak Divertor; June 1999
- NIFS-599 S. Murakami, N. Nakajima, M. Okamoto and J. Nhrenberg,
Effect of Energetic Ion Loss on ICRF Heating Efficiency and Energy Confinement Time in Heliotrons;
June 1999
- NIFS-600 R. Horiuchi and T. Sato,
Three-Dimensional Particle Simulation of Plasma Instabilities and Collisionless Reconnection in a Current Sheet;; June 1999
- NIFS-601 W. Wang, M. Okamoto, N. Nakajima and S. Murakami,
Collisional Transport in a Plasma with Steep Gradients; June 1999
- NIFS-602 T. Mutoh, R. Kumazawa, T. Saku, K. Saito, F. Simpo, G. Nomura, T. Watanabe, X. Jikang, G. Cattanei, H. Okada, K. Ohkubo, M. Sato, S. Kubo, T. Shimozuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B.J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Narihara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohyabu, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Sato, T. Monsaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K. Ida, N. Noda, K. Yamazaki, K. Akaiishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iyoshi and LHD Exp. Group 1 and 2,
First ICRF Heating Experiment in the Large Helical Device; July 1999
- NIFS-603 P.C. de Vries, Y. Nagayama, K. Kawahata, S. Inagaki, H. Sasao and K. Nagasaki,
Polarization of Electron Cyclotron Emission Spectra in LHD; July 1999
- NIFS-604 W. Wang, N. Nakajima, M. Okamoto and S. Murakami,
 δf Simulation of Ion Neoclassical Transport; July 1999
- NIFS-605 T. Hayashi, N. Mizuguchi, T. Sato and the Complexity Simulation Group,
Numerical Simulation of Internal Reconnection Event in Spherical Tokamak, July 1999
- NIFS-606 M. Okamoto, N. Nakajima and W. Wang,
On the Two Weighting Scheme for δf Collisional Transport Simulation; Aug. 1999
- NIFS-607 O. Motojima, A.A. Shishkin, S. Inagaki, K. Y. Watanabe,
Possible Control Scenario of Radial Electric Field by Loss-Cone-Particle Injection into Helical Device; Aug. 1999
- NIFS-608 R. Tanaka, T. Nakamura and T. Yabe,
Constructing Exactly Conservative Scheme in Non-conservative Form; Aug. 1999
- NIFS-609 H. Sugama,
Gyrokinetic Field Theory; Aug. 1999
- NIFS-610 M. Takechi, G. Matsunaga, S. Takagi, K. Ohkuni, K. Toi, M. Osakabe, M. Isobe, S. Okamura, K. Matsuoka, A. Fujisawa, H. Iguchi, S. Lee, T. Minami, K. Tanaka, Y. Yoshimura and CHS Group,
Core Localized Toroidal Alfvén Eigenmodes Destabilized By Energetic Ions in the CHS Heliotron/Torsatron, Sep. 1999
- NIFS-611 K. Ichiguchi,
MHD Equilibrium and Stability in Heliotron Plasmas; Sep. 1999
- NIFS-612 Y. Sato, M. Yokoyama, M. Wakatani and V. D. Pusovitov,
Complete Suppression of Pfirsch-Schluter Current in a Toroidal $l=3$ Stellarator; Oct. 1999
- NIFS-613 S. Wang, H. Sanuki and H. Sugama,
Reduced Drift Kinetic Equation for Neoclassical Transport of Helical Plasmas in Ultra-low Collisionality Regime; Oct. 1999
- NIFS-614 J. Miyazawa, H. Yamada, K. Yasui, S. Kato, N. Fukumoto, M. Nagata and T. Uyama,
Design of Spheromak Injector Using Conical Accelerator for Large Helical Device; Nov. 1999
- NIFS-615 M. Uchida, A. Fukuyama, K. Itoh, S.-I. Itoh and M. Yagi,
Analysis of Current Diffusive Ballooning Mode in Tokamaks; Dec. 1999