



18th IAEA Fusion Energy Conference
Sorrento, Italy, 4 to 10 October 2000

IAEA-CN-77/ EXP3/12

NATIONAL INSTITUTE FOR FUSION SCIENCE

MHD Characteristics in High- β Regime of the Large Helical Device

S. Sakakibara, H. Yamada, K. Y. Watanabe, Y. Narushima, K. Toi, S. Ohdachi, M. Takichi,
S. Yamamoto, K. Narihara, K. Tanaka, N. Ashikawa, P. Devaries, M. Emoto, H. Funaba,
M. Goto, K. Ida, H. Idei, K. Ikeda, S. Inagaki, N. Inoue, M. Isobe, S. Kado, O. Kaneko,
K. Kawahata, K. Khlopenkov, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami,
J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Mutoh, Y. Nagayama,
Y. Nakamura, H. Nakanishi, K. Nishimura, N. Noda, T. Notake, T. Kobuchi, Y. Liang, N. Ohyabu,
Y. Oka, M. Osakabe, T. Ozaki, R.O. Pavlichenko, B.J. Peterson, A. Sagara, K. Saito, R. Sakamoto,
H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimoizuma, M. Shoji, H. Suzuki, Y. Takeiri,
N. Tamura, T. Tokuzawa, Y. Torii, K. Tsumori, I. Yamada, S. Yamaguchi, M. Yokoyama,
Y. Yoshimura, T. Watari, N. Nakajima, K. Ichiguchi, H. Takahashi, A.W. Cooper, K. Yamazaki,
O. Motojima, Y. Hamada, M. Fujiwara

(Received - Sep. 14, 2000)

NIFS-652

Sep. 2000

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

RESEARCH REPORT NIFS Series

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designating Member State(s) or of the designating organization(s). In particular, neither the IAEA nor any other organization or body sponsoring this meeting can be held responsible for any material reproduced in this preprint.

TOKI, JAPAN

MHD Characteristics in High- β Regime of the Large Helical Device

S.Sakakibara 1), H.Yamada 1), K.Y.Watanabe 1), Y.Narushima 1), K.Toi 1), S.Ohdachi 1), M.Takechi 1), S.Yamamoto 3), K.Narihara 1), K.Tanaka 1), N.Ashikawa 2), P.Devaries 1), M.Emoto 1), H.Funaba 1), M.Goto 1), K.Ida 1), H.Idei 1), K.Ikeda 1), S.Inagaki 1), N.Inoue 1), M. Isobe 1), S.Kado 1), O.Kaneko 1), K.Kawahata 1), K.Khlopenkov 1), A.Komori 1), S.Kubo 1), R.Kumazawa 1), S.Masuzaki 1), T.Minami 1), J.Miyazawa 1), T.Morisaki 1), S.Morita 1), S.Murakami 1), S.Muto 1), T.Mutoh 1), Y.Nagayama 1), Y.Nakamura 1), H.Nakanishi 1), K.Nishimura 1), N.Noda 1), T.Notake 3), T.Kobuchi 2), Y.Liang 2), N.Ohyabu 1), Y.Oka 1), M.Osakabe 1), T.Ozaki 1), R.O.Pavlichenko 1), B.J.Peterson 1), A.Sagara 1), K.Saito 3), R.Sakamoto 1), H. Sasao 2), M.Sasao 1), K.Sato 1), M.Sato 1), T. Seki 1), T.Shimozuma 1), M.Shoji 1), H.Suzuki 1), Y.Takeiri 1), N.Tamura 2), T.Tokuzawa 1), Y. Torii 3), K.Tsumori 1), I.Yamada 1), S.Yamaguchi 1), M.Yokoyama 1), Y.Yoshimura 1), T.Watari 1), N.Nakajima 1), K.Ichiguchi 1), H.Takahashi 4), W.A.Cooper 5), K.Yamazaki 1), O.Motojima 1), Y.Hamada 1), M.Fujiwara 1)

1) National Institute for Fusion Science, Toki, Japan

2) Department of Fusion Science, Graduate Univ. for Advanced Studies, Hayama, Japan

3) Department of Energy Engineering and Science, Nagoya Univ., Nagoya, Japan

4) Princeton Plasma Physics Laboratory, Princeton University, Princeton, USA

5) Centre de recherches en physique des plasmas, Ecole Polytechnique Federale de Lausanne, Lausanne, Switzerland

e-mail contact of main author: sakakis@lhd.nifs.ac.jp

Abstract: The highest volume averaged beta value β_i of 2.2% at $B_t = 0.75$ T (gas puff) and 2.4 % at $B_t = 1.3$ T (pellet) in all of helical devices have been achieved in the Large Helical Device (LHD). The β_i dependence of MHD activities has been investigated in NBI plasmas. The $n/m = 1/2$ mode excited in the core region and $\iota = 1$ resonant modes in peripheral region have been observed. Both of the fluctuation amplitudes increase with β_i and the pressure gradient. The strong $n/m = 1/2$ mode which affects plasma profile have been observed in high- β_i discharges and the abrupt disappearance of the mode leads to the restoration of T_e profile. Violent instabilities which terminate the plasma and degradation of global energy confinement have not been observed so far.

1. Introduction

In helical plasmas with no net currents, an understanding of pressure driven modes such as resistive/ideal interchange modes and ballooning modes have been a major key issue for realization of high- β plasmas. In a heliotron configuration, resistive/ideal interchange modes are expected to be unstable easily because there are magnetic hill even in high- β plasmas although magnetic well is formed from center region by the Shafranov shift. The characteristics of excited instabilities and critical beta value strongly depend on the feature of magnetic configuration. Heliotron-E has the strong magnetic shear for stabilization of ideal interchange modes, while there is strong magnetic hill in peripheral region. The internal disruption with particle and energy loss have been triggered by $n/m = 1/1$ mode in high- β plasmas with the large pressure gradient [1]. The control of the pressure profile with a strong gas puff has flattened the pressure gradient around the $\iota = 1$ surface, and the volume averaged beta value of 2 % was realized by the suppression of the internal disruption [2]. In Compact Helical System (CHS) with large finite beta effects related to low-aspect-ratio [3], the volume averaged equilibrium beta value of 2.1 % was achieved by confinement

improvement given by means of turning off gas puff [4]. Then strong global instability was not observed [5].

Since LHD experiments were started in 1998, plasma parameters have been improved with the progress of heating power systems during every experimental campaign [6]. Recent experiments based on high power neutral beam heating enable us to obtain a volume-averaged beta value β_t of over 2 %, where β_t is defined as $2\mu_0\langle p \rangle / B_{av}^2$ and B_{av} is averaged toroidal magnetic field in plasma region. The highest volume averaged beta value β_t of 2.2% at $B_t = 0.75$ T (gas puff) and 2.4 % at $B_t = 1.3$ T (pellet) in all of helical devices has been achieved in the *unstable* configuration with magnetic hill. In previous experiments, the pedestal structure has been observed and has made a great contribution to global plasma confinement [7,8]. However, the structure has formed large pressure gradient in peripheral region with magnetic hill. The $r=1$ resonant modes has been observed in high-field (low- β) plasmas and the relationship between pressure gradient and mode activity has been investigated [9]. On the other hand, $n/m = 1/2$ mode located in the core region also appears as threshold with β_t of about 0.3 % [10]. Here, observation results of MHD activity in NBI plasmas with higher β_t range with up to 2.2 % are reported.

2. High- β Experiments

Figure 1 shows the time behavior of high- β_t discharge of over 2 %. The magnetic axis position R_{ax} and toroidal magnetic field B_t are set at 3.6 m and 0.75 T, respectively. The target plasma was produced by a neutral beam of tangentially co-injection (NBI#2) and maintained by counter neutral beam injection (NBI#1) from 1.32 s addition to NBI#2. Hydrogen gas was supplied by gas puff from 0.6 s to the end of discharge, and line averaged electron density \bar{n}_e gradually increase with time and approaches $3 \times 10^{19} \text{ m}^{-3}$. The β_t^{dia} estimated by the diamagnetic flux measurement approaches about 2.2 % at 2.75 s. Net plasma current I_p flows in the co- direction and reaches about 20 kA at the end of discharge. It has been expected that this I_p is composed of Ohkawa currents produced by unbalanced NBI and bootstrap currents [11]. The β_t^{dia} is saturated at 2.45 s and increases again at 2.66 s. While \bar{n}_e monotonically increase with time, the central electron temperature T_{e0} and the central plasma pressure decrease from 1.72 s and starts to rise at 2.68 s. The magnetic fluctuation with $f \leq 50$ kHz was measured in the time range from 1.7 s to 3.0 s, and the coherent modes with $n/m = 1/2$, 2/2 and 2/3 were observed. These modes have rotational frequency of 1 kHz, 3 kHz and 8 kHz, respectively. The amplitude of $n/m = 1/2$ mode increases and saturate when $\beta_t^{dia} = 2.1$ %, and this mode abruptly disappears when β_t^{dia} starts to increase. The time at which the $n/m = 2/3$ mode starts to grow up is the same as the $n/m = 1/2$ mode




FIG. 1 Time behavior of high- β discharge with gas puff and the extended view in the time window from 1.7 s to 3.0 s.

case, and there is no change of the amplitude when the $n/m = 1/2$ mode disappears.

Figure 2 shows the rotational transform and T_e profiles before and after the disappearance of the $n/m = 1/2$ mode. The local flattening of T_e profile around $r = 1/2$ surface has been observed when the $n/m = 1/2$ mode exists. Although this structure is intermittently observed, it appears conspicuously with the increase in the fluctuation amplitude of the $n/m = 1/2$ mode. The width of local flattening Δ/a is about 0.3, and this is consistent with an island width estimated by using measured radial component of the $n/m = 1/2$ mode amplitude. The T_e flattening disappears with the $n/m = 1/2$ mode after 2.66 s. Then T_e profile has recovered a peaked one and keeps the shape to the end of discharge. The \bar{n}_e increases in this phase, with keeping the hollow profile.

One of the possibility of this phenomenon is the disappearance of the $r = 1/2$ surface caused by plasma currents in the co- direction and finite- β effects. The similar phenomenon was also observed in Ohmic current experiments of CHS [12]. The rotational transform profiles shown in Fig.2 are calculated by using 3-D MHD equilibrium code VMEC [13] for the currentless case of $I_p = 0$ kA and the cases of $I_p = 18$ kA with broad current profile ($j = j_0(1 - \rho^2)$) and peaked one ($j = j_0(1 - \rho^2)^3$). Where $\rho = (\Psi/\Psi^*)^{1/2}$ and Ψ is the toroidal flux function which is normalized by the value of the last closed flux surface. At 2.52 s, the T_e flattening is observed at $R = 3.25 \sim 3.50$ m and $3.85 \sim 4.10$ m. The relation between these positions and predicted $r = 1/2$ surfaces suggest that I_p has broader profile than that with $j = j_0(1 - \rho^2)$. Also, the fact that I_p has a little change of 1 kA before and after an extinction of the $n/m = 1/2$ mode means that the disappearance of the $r = 1/2$ surface may be caused by the change in the current profile rather than absolute value of I_p . The R_{ax} derived from T_e profile at 2.52 s is about 3.65 m and the Shafranov shift is relatively small because the decrease in central β value. After disappearance of the $n/m = 1/2$ mode, R_{ax} shifts to about 3.75m (at 2.8 s), which causes a rise of central rotational transform through Shafranov shift addition to the increase in I_p .




FIG. 2 Rotational transform and T_e profile at (a) 2.52 s and (b) 2.80 s.

3. Beta Dependence of MHD modes

Figure 3(a) shows the changes in fluctuation amplitudes of observed global modes with $m \leq 3$ with β_t^{dia} in 16 NBI plasmas with $R_{ax} = 3.6$ m and $B_t \leq 1.5$ T. Here, discharges with I_p of less than 5kA (at $B_t = 0.75$ T) are selected except that with $\beta_t^{dia} > 1.8$ % in the case of Fig.1. This means that the changes in location of resonant surface due to I_p is relative small when $\beta_t^{dia} \leq 1.8$ % in this figure. The fluctuation amplitude is a root mean square of the mode amplitude in the time window of 20 ms and normalized by the vacuum magnetic field at probe position. Fluctuation amplitudes of $r = 1$ resonant modes with $n/m = 1/1, 2/2$ and $3/3$ increase with β_t^{dia} and reach about 10^{-4} when β_t^{dia} approaches about 2 %. The $n/m = 1/2$ mode appears when β_t^{dia} exceeds about 0.3 %. The fluctuation amplitudes keep to increase with β_t^{dia} and approaches 8×10^{-4} when β_t^{dia} is about 1.4 %.

The changes in the β gradients at the specific surfaces with β_t^{dia} are shown in Fig.3 (b).

The open circle and square are β gradients at $\rho = 0.5$ and 0.9 , respectively, and they approximately correspond to the major rational $\iota = 1/2$ and 1 surfaces. The data are selected at the conditions that the plasmas reach quasi-steady state and gas puff is still turned on. The increase of β gradients at $\rho = 0.5$ in $\beta_t^{dia} \geq 0.5\%$ is larger than in $\beta_t^{dia} < 0.5\%$ and this corresponds to the appearance of $n/m = 1/2$ mode. The β gradient at $\rho = 0.9$ increases linearly with β_t^{dia} .

The time development of β gradients at $\rho = 0.5$ and 0.9 surfaces in the discharge of Fig.1 (Shot#16964) are shown in Fig.3 (b). The time range is from 1.2 s to 3.3 s and divided into three phases, that is, A: before and B: during the growth of $n/m = 1/2$ mode, and C: after the disappearance of this mode. The β gradient at $\rho = 0.9$ increases and saturates during the phase A and keeps the constant to the end of discharge. This saturation depends on NBI power, and the same behavior of β gradient has been also observed in low-power discharges. In phase A (open triangle), the β gradient at $\rho = 0.5$ is much smaller than that in quasi-steady state plasmas and the amplitude of $n/m = 1/2$ mode is relatively small. In the phase B (closed triangle), while the β gradient abruptly rise to that in quasi-steady state plasmas with $\beta_t^{dia} = 1.4\%$, the $n/m = 1/2$ mode amplitude starts to increase. The amplitude approaches 7×10^{-4} at the end of phase B, and it is the same or small compared with the maximum value of $n/m = 1/2$ mode amplitude in $\beta_t^{dia} \geq 1.4\%$. This fact suggests the possibility that $n/m = 1/2$ mode observed in $\beta_t^{dia} \geq 1.4\%$ also affect the plasma profile and restricts local β gradient. The restoration of pressure profile after the disappearance of the $n/m = 1/2$ mode in phase C is consistent with this suggestion.

4. Discussions and Summary

In high- β_t experiments, a magnetic configuration with an inward-shifted R_{ax} of 3.6 m has been selected because neoclassical transport and particle confinement of high-energy ions are superior to the outward-shifted case. However, this configuration is unfavorable from the viewpoint of MHD stability. Theoretical prediction suggests that the high- n ballooning mode is unstable when the central beta is 8 % (peaked profile) [14], and destabilization of the interchange mode becomes a major concern. The stability beta limits estimated by the low- n ideal mode analysis and the Mercier criterion are higher for the outward shifted plasma than the inward shifted case because of magnetic well formation. The good energy confinement in the inward-shifted configuration has been obtained in experiments [15].

As experimental results, $\iota = 1$ resonant modes in peripheral region and the $n/m = 1/2$ mode in core region have been dominantly observed in the β_t^{dia} range of up to 2.2 %. The $\iota = 1$ resonant modes have been observed even in low- β_t region and the amplitude keep to increase with β_t^{dia} and pressure gradient. The $n/m = 1/2$ mode appears when β_t^{dia} reaches to




FIG 3 (a) Changes in the amplitudes of observed coherent modes and (b) $d\beta/d\rho$ at $\rho = 0.5$ and 0.9 surfaces as a function of β_t^{dia} .

0.3% and the amplitude increases with β_t^{dia} . In the discharge with over 2 %, the $n/m = 1/2$ mode which affects the local β gradient has been observed. The reason for the growth of $n/m = 1/2$ mode amplitude in a discharge is the abrupt increase in local β gradient, which reaches about the same value as that in quasi-steady state plasmas with $\beta_t^{dia} = 1.4$ %. The maximum value of the $n/m = 1/2$ modes observed in $\beta_t^{dia} \geq 1.4$ % have the same or higher amplitudes compared with the $n/m = 1/2$ mode which affects plasma profile. There is a possibility that the local β gradient is limited by the $n/m = 1/2$ mode in $\beta_t^{dia} \geq 1.4$ %. The restoration of the pressure profile due to the disappearance of $n/m = 1/2$ mode is consistent with the possibility.

Figure 4 shows the unstable region of ideal $n/m = 1/2$ modes in the plane of β_t^{dia} and β gradient at $\rho = 0.5$. MHD stability has been calculated by the 3-D MHD stability analysis code TERPSICHORE [16] for different pressure profile. The measured β gradient at $\rho = 0.5$, which approximately correspond to $r = 1/2$ surface, fits that in the $P = P_0(1 - \rho^2)^{0.5-1}$ cases. In this case, the present operational region is located in marginal against the low- n ideal instability. If observed $n/m = 1/2$ mode is related to ideal interchange mode, the discharge with keeping low β gradient in the unstable- β_t^{dia} region may realize the production of stable higher- β_t plasmas without destabilization of this mode (dotted line with an arrow in Fig. 4).

The authors would like to thank all members of the device engineering group for their operational support.

Reference

- [1] HARRIS, J.H., et al., Phys. Rev. Lett. **53** (1984) 2242.
- [2] MOTOJIMA, O., et al., Nucl. Fusion **25** (1985) 1783.
- [3] YAMADA, H., et al., in the Proc. 14th Int. Conf. on Plasma Physics and Controlled Nuclear Fusion Res. 1992 Vol.2, IAEA, Vienna (1993) 493.
- [4] OKAMURA, S., et al., Nucl. Fusion **35** (1995) 283.
- [5] SAKAKIBARA, S., et al., J. Phys. Soc. Jpn **63** (1994) 4406.
- [6] MOTOJIMA, O., et al. Phys. Plasmas **6** (1999) 1843.
- [7] YAMADA, H., et al., Phys. Rev. Lett. **84** (2000) 1216.
- [8] OHYABU, N., et al., Phys. Rev. Lett. **84** (2000) 103.
- [9] TOI, K., et al., in the Proc. of 27th EPS Conf. on Controlled Fusion and Plasma Physics, Budapest, 2000
- [10] TAKECHI, M., et al., in the Proc. of 10th Int. Toki Conf, 1999
- [11] WATANABE, K.Y., et al., in the Proc of 27th EPS Conf. on Controlled Fusion and Plasma Physics., Budapest, 2000
- [12] SAKAKIBARA, S., et al., Jpn. J.Appl. Phys. **63** (1994) 4406.
- [13] HIRSHMAN, S.P., et al., Comput. Phys. Commun. **43** (1986) 143.
- [14] NAKAJIMA, N., et al., Phys. Plasmas **3** (1996) 4556.
- [15] FUJIWARA, M., et al., OV1/4 in this conference.
- [16] COOPER, W.A., et al., Plasma Pys. Control. Fusion **34** (1992) 1011.




FIG 4 Unstable region of ideal $n/m = 1/2$ modes in the plane of β_t^{dia} and β gradient at $\rho = 0.5$.

Recent Issues of NIFS Series

- NIFS-600 R Honuchi and T Sato,
Three-Dimensional Particle Simulation of Plasma Instabilities and Collisionless Reconnection in a Current Sheet, June 1999
- NIFS-601 W Wang, M Okamoto, N Nakajima and S Murakami
Collisional Transport in a Plasma with Steep Gradients; June 1999
- NIFS-602 T Mutoh, R Kumazawa, T Saki, K Sato, F Simpo, G Nomura, T Watan, X Jikang, G Cattanei, H Okada, K Ohkubo, M Sato, S Kubo, T Shimozuma, H Idei, Y Yoshimura, O Kaneko, Y Takeiri, M Osakabe, Y Oka, K Tsumori, A Komori, H Yamada, K Watanabe, S Sakakibara, M Shoji, R Sakamoto, S Inagaki, J Miyazawa, S Monta, K Tanaka, BJ Peterson, S Murakami, T Minami, S Ohdachi, S Kado, K Narihara, H Sasao, H Suzuki, K Kawahata, N Ohyabu, Y Nakamura, H Funaba, S Masuzaki, S Muto, K Sato, T Monsaki, S Sudo, Y Nagayama, T Watanabe, M Sasao, K Ida, N Noda, K Yamazaki, K Akaishi, A Sagara, K Nishimura, T Ozaki, K Toi, O Motojima, M Fujiwara, A Iiyoshi and LHD Exp. Group 1 and 2,
First ICRF Heating Experiment in the Large Helical Device; July 1999
- NIFS-603 P.C de Vries, Y Nagayama, K Kawahata, S Inagaki, H Sasao and K Nagasaki,
Polarization of Electron Cyclotron Emission Spectra in LHD, July 1999
- NIFS-604 W Wang, N Nakajima, M Okamoto and S Murakami,
 δf Simulation of Ion Neoclassical Transport; July 1999
- NIFS-605 T Hayashi, N Mizuguchi, T Sato and the Complexity Simulation Group,
Numerical Simulation of Internal Reconnection Event in Spherical Tokamak; July 1999
- NIFS-606 M Okamoto, N Nakajima and W. Wang,
On the Two Weighting Scheme for δf Collisional Transport Simulation; Aug 1999
- NIFS-607 O. Motojima, AA Shishkin, S. Inagaki, K.Y. Watanabe,
Possible Control Scenario of Radial Electric Field by Loss-Cone-Particle Injection into Helical Device; Aug. 1999
- NIFS-608 R. Tanaka, T. Nakamura and T. Yabe,
Constructing Exactly Conservative Scheme in Non-conservative Form; Aug. 1999
- NIFS-609 H. Sugama,
Gyrokinetic Field Theory, Aug 1999
- NIFS-610 M. Takechi, G. Matsunaga, S. Takagi, K. Ohkuni, K. Toi, M. Osakabe, M. Isobe, S. Okamura, K. Matsuoka, A. Fujisawa, H. Iguchi, S. Lee, T. Minami, K. Tanaka, Y. Yoshimura and CHS Group,
Core Localized Toroidal Alfvén Eigenmodes Destabilized By Energetic Ions in the CHS Heliotron/Torsatron; Sep. 1999
- NIFS-611 K. Ichiguchi,
MHD Equilibrium and Stability in Heliotron Plasmas; Sep 1999
- NIFS-612 Y Sato, M Yokoyama, M. Wakatani and V D. Puvotitov,
Complete Suppression of Pfirsch-Schluter Current in a Toroidal $I=3$ Stellarator; Oct. 1999
- NIFS-613 S. Wang, H. Sanuki and H. Sugama,
Reduced Drift Kinetic Equation for Neoclassical Transport of Helical Plasmas in Ultra-low Collisionality Regime; Oct 1999
- NIFS-614 J. Miyazawa, H Yamada, K Yasui, S Kato, N. Fukumoto, M Nagata and T Uyama,
Design of Spheromak Injector Using Conical Accelerator for Large Helical Device; Nov 1999
- NIFS-615 M. Uchida, A Fukuyama, K Itoh, S-I Itoh and M. Yagi,
Analysis of Current Diffusive Ballooning Mode in Tokamaks, Dec 1999

- NIFS-616 M. Tanaka, A.Yu Grosberg and T. Tanaka,
Condensation and Swelling Behavior of Randomly Charged Multichain Polymers by Molecular Dynamics Simulations; Dec. 1999
- NIFS-617 S. Goto and S. Kida,
Sparseness of Nonlinear Coupling; Dec. 1999
- NIFS-618 M.M. Skoric, T. Sato, A. Maluckov and M.S. Jovanovic,
Complexity in Laser Plasma Instabilities Dec 1999
- NIFS-619 T.-H. Watanabe, H. Sugama and T. Sato,
Non-dissipative Kinetic Simulation and Analytical Solution of Three-mode Equations of Ion Temperature Gradient Instability; Dec 1999
- NIFS-620 Y. Oka, Y. Takeiri, Yu.I. Belchenko, M. Hamabe, O. Kaneko, K. Tsumori, M. Osakabe, E. Asano, T. Kawamoto, R. Akiyama,
Optimization of Cs Deposition in the 1/3 Scale Hydrogen Negative Ion Source for LHD-NBI System ;Dec. 1999
- NIFS-621 Yu.I. Belchenko, Y. Oka, O. Kaneko, Y. Takeiri, A. Krivenko, M. Osakabe, K. Tsumori, E. Asano, T. Kawamoto, R. Akiyama,
Recovery of Cesium in the Hydrogen Negative Ion Sources;Dec 1999
- NIFS-622 Y. Oka, O. Kaneko, K. Tsumori, Y. Takeiri, M. Osakabe, T. Kawamoto, E. Asano, and R. Akiyama,
H- Ion Source Using a Localized Virtual Magnetic Filter in the Plasma Electrode: Type I LV Magnetic Filter; Dec. 1999
- NIFS-623 M. Tanaka, S. Kida, S. Yanase and G. Kawahara,
Zero-absolute-vorticity State in a Rotating Turbulent Shear Flow;Jan 2000
- NIFS-624 F. Leuterer, S. Kubo,
Electron Cyclotron Current Drive at $\omega \approx \omega_c$ with X-mode Launched from the Low Field Side; Feb. 2000
- NIFS-625 K. Nishimura,
Wakefield of a Charged Particulate Influenced by Emission Process of Secondary Electrons; Mar. 2000
- NIFS-626 K. Itoh, M. Yagi, S.-I. Itoh, A. Fukuyama,
On Turbulent Transport in Burning Plasmas;Mar. 2000
- NIFS-627 K. Itoh, S.-I. Itoh, L. Giannone,
Modelling of Density Limit Phenomena in Toroidal Helical Plasmas; Mar. 2000
- NIFS-628 K. Akaishi, M. Nakasuga and Y. Funato,
True and Measured Outgassing Rates of a Vacuum Chamber with a Reversibly Absorbed Phase;Mar 2000
- NIFS-629 T. Yamagishi,
Effect of Weak Dissipation on a Drift Orbit Mapping; Mar. 2000
- NIFS-630 S. Toda, S.-I. Itoh, M. Yagi, A. Fukuyama and K. Itoh,
Spatial Structure of Compound Dither in L/H Transition;Mar. 2000
- NIFS-631 N. Ishihara and S. Kida,
Axial and Equatorial Magnetic Dipoles Generated in a Rotating Spherical Shell; Mar. 2000
- NIFS-632 T. Kuroda, H. Sugama, R. Kanno and M. Okamoto,
Ion Temperature Gradient Modes in Toroidal Helical Systems; Apr. 2000

- NIFS-633 VD Pustovitov,
Magnetic Diagnostics: General Principles and the Problem of Reconstruction of Plasma Current and Pressure Profiles in Toroidal Systems. Apr 2000
- NIFS-634 AB Mikhailovskii, SV Kononov, VD Pustovitov and VS Tsypin,
Mechanism of Viscosity Effect on Magnetic Island Rotation. Apr 2000
- NIFS-635 H Naitou, T Kuramoto, T. Kobayashi, M Yagi, S Tokuda and T Matsumoto,
Stabilization of Kinetic Internal Kink Mode by Ion Diamagnetic Effects. Apr 2000
- NIFS-636 A Kageyama and S.Kida,
A Spectral Method in Spherical Coordinates with Coordinate Singularity at the Origin. Apr. 2000
- NIFS-637 R. Honuchi, W Pei and T Sato,
Collisionless Driven Reconnection in an Open System. June 2000
- NIFS-638 K Nagaoka, A Okamoto, S Yoshimura and MY Tanaka,
Plasma Flow Measurement Using Directional Langmuir Probe under Weakly Ion-Magnetized Conditions; July 2000
- NIFS-639 Alexei Ivanov,
Scaling of the Distribution Function and the Critical Exponents near the Point of a Marginal Stability under the Vlasov-Poisson Equations; Aug 2000
- NIFS-640 K. Ohi, H Naitou, Y Tauchi, O. Fukumasa,
Observation of the Limit Cycle in the Asymmetric Plasma Divided by the Magnetic Filter; Aug 2000
- NIFS-641 H. Momota, G.H Miley and J. Nadler,
Direct Energy Conversion for IEC Propulsions; Aug. 2000
- NIFS-642 Y Kondoh, T Takahashi and H. Momota,
Revisit to the Helicity and the Generalized Self-organization Theory; Sep. 2000
- NIFS-643 H. Soltwisch, K. Tanaka,
Changes of the Electron Density Distribution during MHD Activity in CHS; Sep 2000
- NIFS-644 Fujisawa, A., Iguchi, H., Minami, T., Yoshimura, Y., Sanuki, H., Itoh, K., Isobe, M., Nishimura, S., Tanaka, K., Osakabe, M., Nomura, I., Ida, K., Okamura, S., Toi, K., Kado, S., Akiyama, R., Shimizu, A., Takahashi, C., Kojima, M., Matsuoka, K., Hamada, Y., Fujiwara, M.,
Observation of Bifurcation Property of Radial Electric Field Using a Heavy Ion Beam Probe; Sep 2000
 (IAEA-CN-77/EX6/6)
- NIFS-645 Todo, Y., Watanabe, T.-H., Park, H.-B., Sato, T.,
Fokker-Planck Simulation Study of Alfvén Eigenmode Burst Sep 2000
 (IAEA-CN-77/TH6/2)
- NIFS-646 Muroga, T., Nagasaka, T.,
Development of Manufacturing Technology for High Purity Low Activation Vanadium Alloys; Sep 2000
 (IAEA-CN-77/FTP1/09)
- NIFS-647 Itoh, K., Sanuki, H., Toda, S., Itoh, S.-I., Yagi, M., Fukuyama, A., Yokoyama, M.,
Theory of Dynamics in Long Pulse Helical Plasmas; Sep. 2000
 (IAEA-CN-77/THP1/19)
- NIFS-648 R. Sakamoto, H. Yamada, K. Tanaka, K. Narihara, S. Morita, S. Sakakibara, S. Masuzaki, L.R. Baylor,

P.W. Fisher, S.K. Combs, M.J. Gouge, S. Kato, A. Komori, O. Kaneko, N. Ashikawa, P. de Varies, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, K. Ikeda, S. Inagaki, M. Isobe, S. Kado, K. Kawahata, K. Khlopenkov, S. Kubo, R. Kumazawa, T. Minami, J. Miyazawa, T. Morisaki, S. Murakami, S. Muto, T. Mutoh, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Nishimura, N. Noda, T. Notake, T. Kobuchi, Y. Liang, S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, R.O. Pavlichenko, B.J. Peterson, A. Sagara, K. Saito, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimoizuma, M. Shoji, S. Sudo, H. Suzuki, M. Takechi, Y. Takeiri, N. Tamura, K. Toi, T. Tokuzawa, Y. Torii, K. Tsumori, I. Yamada, S. Yamaguchi, S. Yamamoto, Y. Yoshimura, K.Y. Watanabe, T. Watari, K. Yamazaki, Y. Hamada, O. Motojima and M. Fujiwara,

Impact of Pellet Injection on Extension of Operational Region in LHD: Sep. 2000
(IAEA-CN-77/ EXP4/18)/

NIFS-649 K. Ichiguchi, M. Wakatani, T. Umemura, T. Tatsuno and B.A. Carreras

Improved Stability due to Local Pressure Flattening in Stellarators: Sep 2000
(IAEA-CN-77/ THP2/08)

NIFS-650

N. Noda, Y. Nakamura, Y. Takeiri, T. Mutoh, R. Kumazawa, M. Sato, K. Kawahata, S. Yamada, T. Shimoizuma, Y. Oka, A. Iiyoshi, R. Sakamoto, Y. Kubota, S. Masuzaki, S. Inagaki, T. Morisaki, H. Suzuki, N. Ohyabu, K. Adachi, K. Akaishi, N. Ashikawa, H. Chikaraishi, P. C. de Vries, M. Emoto, H. Funaba, M. Goto, S. Hamaguchi, K. Ida, H. Idei, K. Ikeda, Imagawa, N. Inoue, M. Isobe, A. Iwamoto, S. Kado, O. Kaneko, S. Kitagawa, K. Khlopenkov, T. Kobuchi, A. Komori, S. Kubo, Y. Liang, R. Maekawa, T. Minami, T. Mito, J. Miyazawa, S. Morita, K. Murai, S. Murakami, S. Muto, Y. Nagayama, H. Nakanishi, K. Narihara, A. Nishimura, K. Nishimura, A. Nishizawa, T. Notake, S. Ohdachi, M. Okamoto, M. Osakabe, T. Ozaki, R. O. Pavlichenko, B. J. Peterson, A. Sagara, K. Saito, S. Sakakibara, H. Sasao, M. Sasao, K. Sato, T. Seki, M. Shoji, H. Sugama, K. Takahata, M. Takechi, H. Tamura, N. Tamura, K. Tanaka, K. Toi, T. Tokuzawa, Y. Torii, K. Tsumori, K. Y. Watanabe, T. Watanabe, T. Watari, N. Yanagi, I. Yamada, H. Yamada, S. Yamaguchi, S. Yamamoto, T. Yamamoto, M. Yokoyama, Y. Yoshimura, I. Ohtake, R. Akiyama, K. Haba, M. Iima, J. Kodaira, K. Tsuzuki, K. Itoh, K. Matsuoka, K. Ohkubo, S. Satoh, T. Satow, S. Sudo, S. Tanahashi, K. Yamazaki, O. Motojima, Y. Hamada, M. Fujiwara,

Experimental Studies toward Long-Pulse Steady-State Operations in LHD: Sep. 2000
(IAEA-CN-77/ EX4/5)

NIFS-651

H. Yamada, K.Y. Watanabe, K. Yamazaki, S. Murakami, S. Sakakibara, K. Narihara, K. Tanaka, M. Osakabe, K. Ida, G. Rewoldt, N. Ashikawa, P. de Varies, M. Emoto, H. Funaba, M. Goto, H. Idei, K. Ikeda, S. Inagaki, N. Inoue, M. Isobe, S. Kado, O. Kaneko, K. Kawahata, K. Khlopenkov, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Muto, T. Mutoh, Y. Nagayama, N. Nakajima, Y. Nakamura, H. Nakanishi, K. Nishimura, N. Noda, T. Notake, T. Kobuchi, Y. Liang, S. Ohdachi, N. Ohyabu, Y. Oka, T. Ozaki, R.O. Pavlichenko, B.J. Peterson, A. Sagara, K. Saito, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimoizuma, M. Shoji, H. Sugama, H. Suzuki, M. Takechi, Y. Takeiri, N. Tamura, K. Toi, T. Tokuzawa, Y. Torii, K. Tsumori, I. Yamada, S. Yamaguchi, S. Yamamoto, M. Yokoyama, Y. Yoshimura, T. Watari, K. Itoh, K. Matsuoka, K. Ohkubo, I. Ohtake, S. Satoh, T. Satow, S. Sudo, S. Tanahashi, T. Uda, Y. Hamada, O. Motojima, M. Fujiwara,

Energy Confinement and Thermal Transport Characteristics of Net-Current Free Plasmas in Large Helical Device: Sep. 2000
(IAEA-CN-77/ EX6/7)

NIFS-652

S. Sakibara, H. Yamada, K. Y. Watanabe, Y. Narushima, K. Toi, S. Ohdachi, M. Takichi, S. Yamamoto, K. Narihara, K. Tanaka, N. Ashikawa, P. Devaries, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, K. Ikeda, S. Inagaki, N. Inoue, M. Isobe, S. Kado, O. Kaneko, K. Kawahata, K. Khlopenkov, A. Komori, S. Kubo, R. Kumazawa, S. Masuzaki, T. Minami, J. Miyazawa, T. Morisaki, S. Morita, S. Murakami, S. Muto, T. Mutoh, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Nishimura, N. Noda, T. Notake, T. Kobuchi, Y. Liang, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, R.O. Pavlichenko, B.J. Peterson, A. Sagara, K. Saito, R. Sakamoto, H. Sasao, M. Sasao, K. Sato, M. Sato, T. Seki, T. Shimoizuma, M. Shoji, H. Suzuki, Y. Takeiri, N. Tamura, T. Tokuzawa, Y. Torii, K. Tsumori, I. Yamada, S. Yamaguchi, M. Yokoyama, Y. Yoshimura, T. Watari, N. Nakajima, K. Ichiguchi, H. Takahashi, A.W. Cooper, K. Yamazaki, O. Motojima, Y. Hamada, M. Fujiwara

MHD Characteristics in High- β Regime of the Large Helical Device: Sep. 2000
(IAEA-CN-77/ EXP3/12)