

# NATIONAL INSTITUTE FOR FUSION SCIENCE

## Comparison of Recombination Rate Coefficients Given by Empirical Formulas for Ions from Hydrogen through Nickel

T. Kato and E. Asano

(Received - Apr. 22, 1999 )

NIFS-DATA-54

June 1999

### RESEARCH REPORT NIFS-DATA Series

This report was prepared as a preprint of compilation of evaluated atomic, molecular, plasma-wall interaction, or nuclear data for fusion research, performed as a collaboration research of the Data and Planning Center, the National Institute for Fusion Science (NIFS) of Japan. This document is intended for future publication in a journal or data book after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

NAGOYA, JAPAN

# **Comparison of Recombination Rate Coefficients given by Empirical Formulas for Ions from Hydrogen through Nickel**

**T. Kato and E. Asano**

**National Institute for Fusion Science  
Toki, Gifu 509 - 5292, Japan**

## **Abstract**

Recombination rate coefficients based on several empirical formulae are compared for ions of H, He, C, N, O, Ne, Na, Mg, Al, Si, S, P, Cl, Ar, Ca, Fe and Ni. The total rate coefficients including radiative recombination and dielectronic recombination are shown in graphs.

## **Key words**

**recombination rate coefficients, radiative recombination, dielectronic recombination, plasma, ions, empirical formulae**

## 1. Introduction

Recombination rate coefficients as well as ionization rate coefficients are important in understanding ionization balance in both laboratory and astrophysical plasmas. We compared ionization rate coefficients by several empirical formulas for various ions in Ref.1. From both experimental and theoretical viewpoints, recombination is less well-studied than ionization. Experimental data is not plentiful. In this report we compare recombination rate coefficients given by empirical formulas which are commonly used for laboratory and astrophysical plasmas. We compare values for the total recombination rate coefficient which is the sum of radiative recombination and dielectric recombination at low density limit. We do not compare three body recombination rate coefficients since they are not important at low density plasmas and they can be easily derived from ionization rate coefficients. As we will see in the graphs of this paper, the dielectronic recombination rate coefficients are confused for L and M shell ions. The empirical formulas examined here are not new. However we hope our comparison will stimulate an improvement of this data in the future.

Recently detailed calculations for dielectronic recombination have been published as well as the experimental data<sup>2</sup>. We do not include these individual data in our comparison but we hope new empirical formulas based on these new data can be made in the near future.

## 2. Empirical formula

We have compared the following six empirical formulas. The rate coefficients given here are the sum over all the excited states of a final ion from the ground state of an initial ion.

### 2.1. Aldrovandi and Pequingot <sup>3</sup>

Radiative and dielectronic recombination rate coefficients are calculated for all non-hydrogenic ions of He, C, N, O, Ne, Na, Mg, Al, Si, P, S, Cl, Ar and Ca.

The results are given for the Maxwell averaged radiative recombination rate coefficient:

$$\alpha_r = A_r (T_e / 10^4)^{-\eta} \quad (1)$$

where  $T_e$  is the electron temperature in K. The dielectronic recombination rate coefficient is given by

$$\alpha_{di} = A_{di} T_e^{-3/2} \exp(-T_0/T_e) (1 + B_{di} \exp(-T_i/T_e)) \quad (2)$$

$A_r$ ,  $\eta$ ,  $A_{di}$ ,  $T_0$ ,  $B_{di}$ ,  $T_i$  are fit parameters in Eqs.(1) and (2). The fit values are given in Tables in Ref.3 for each ion.

### 2.2 Mewe

Atomic data by Mewe<sup>4, 5</sup> are widely used especially for astrophysical studies, solar flares and X-ray astronomy. The radiative recombination rate coefficient is approximated by

$$\alpha_{Z,z}^r = 10^{-10} \alpha z^{2\eta+1} \Delta T_e^{-\eta} \quad \text{cm}^3 \text{ s}^{-1} \quad (3)$$

where  $\alpha$ ,  $\eta$  and  $\Delta$  are given in table II in Ref.4. Z and z are the nuclear charge and ionic charge of the ion respectively.

The dielectronic recombination rate coefficient for  $z < Z - 4$  is approximated by

$$\alpha_{Z,z}^d = 10^{-3} T_e^{-3/2} (z^2 + 13.4)^{-1/2} \\ \times \sum \alpha_i z^{\beta_i} \exp(-10^4 g_i z \delta_i / T_e) \quad \text{cm}^3 \text{ s}^{-1} \quad (4)$$

where the parameters  $\alpha_i$  etc. are given in table III in Ref. 4. For higher z the results are presented in Ref.5.

The elements included are H, He, C, N, O, Ne, Na, Mg, Al, Si, S, Ar, Ca, Fe and Ni. For other elements, P and Cl, the parameters are scaled (K. Masai<sup>6</sup>) and used for the graphs in this report.

### 2.3 Hulse

The atomic transport code called MIST

by R. Hulse<sup>7</sup> is often used in the modelling of magnetic confined fusion plasmas. The atomic data from MIST were kindly given to us by Prof. T. Amano in National Institute for Fusion Science in 1980's and we use that data in our comparison.

#### 2.4 Arnaud et al.

Arnaud and Rothenflug<sup>8</sup> evaluated cross sections for ions of H, He, C, N, O, Ne, Na, Mg, Al, Si, S, Ar, Ca, Fe, Ni which are major species in astrophysical plasmas. Later Arnaud and Raymond<sup>9</sup> updated the cross section for iron ions. They used parametric formulas and the values for parameters are given in tables in their papers. The parametric fits are valid only in a given temperature range.

Radiative recombination to H-like ions is calculated using the formula given by Seaton<sup>10</sup>,

$$\alpha_r = 5.197 \times 10^{-14} Z\lambda^{1/2} \{ 0.4288 + 0.5\ln(\lambda) + 0.469\lambda^{-1/3} \} \text{ cm}^3 \text{ s}^{-1} \quad (5)$$

where  $\lambda = 157890 Z^2/T(K)$  and Z is the nuclear charge. For non hydrogenic ions Eq. (1) from Ref. 3 is used.

The dielectronic recombination rate coefficients are fitted by Eq. (2) except for recombination of He-like ions. For the dielectronic recombination of He-like ions to Li-like ions the formula given by Younger<sup>11</sup> is used,

$$\alpha_d = 7.6 \times 10^{-11} A(z) \exp(-D(z)/x) x^{3/2} \text{ cm}^3 \text{ s}^{-1} \quad (6)$$

where  $x = kT/I(Li)$  and  $z = Z-2$

$$A(z) = (z+1)^3 / [z^2(z^2+13.4)^{1/2}(1+0.16(z+1) + 0.017(z+1)^2)]$$

$$D(z) = 3.0 \{(z+1)^2/z^2\} / \{1+0.015z^3/(z+1)^2\}$$

$I(Li)$  is the ionization potential of the Li-like ion.

For Fe ions they fit the rate coefficients with the formula,

$$\alpha_d = T^{-2/3} \sum_i c_i \exp(-E_i/kT) \text{ cm}^3 \text{ s}^{-1} \quad (7)$$

The parameters  $c_i$  and  $E_i$  are given in Table 3 in Ref.9.

#### 2.5 Itikawa et al

Itikawa, Kato and Sakimoto<sup>12</sup> evaluated the rate coefficients for electron impact excitation, ionization and recombination for the analysis of X-ray spectra of H-like and He-like S, Ca and Fe ions from solar flares. For the radiative recombination processes they used formulas by Arnaud et al. For dielectronic recombination rate coefficients, several data are compared. They found large discrepancies between different data sources for L-shell ions such as Li -> Be and Be -> B like ions. The recommended data are fitted by Eq. (7) and the fit parameters are given for H-like, He-like, Li-like and Be-like ions as the initial ions. They adopted the data by Romanik<sup>13</sup> for the dielectronic recombination of Li- and Be-like ions.

Figures 1, 2 and 3 show the recommended data by Itikawa et al. for S, Ca and Fe ions, respectively. The radiative recombination (RR) and dielectronic recombination (DR) are separately indicated in the figures. It is seen that DR process is important for low charged ions.

#### 3. Graphs

The total recombination rate coefficients (radiative recombination and dielectronic recombination) by Aldrovandi and Pequignot<sup>3</sup>, Arnaud et al<sup>8,9</sup>, Mewe et al<sup>5</sup>, Hulse<sup>7</sup> and Itikawa et al<sup>12</sup>. are compared by the element from H through Ni (H, He, C, N, O, Ne, Na, Mg, Al, Si, P, S, Cl, Ar, Ca, Fe, Ni ) as a function of electron temperature.

#### 4. Summary

From the comparison of the graphs large discrepancies are found especially for low charged ions. In order to discuss the ionization balance in plasma we need more reliable data for recombination. The discrepancies are mainly due to dielectronic recombination rate coefficients. Recently more accurate data are becoming available from both theoretical calculation and experimental measurements. We are making a database for these individual data (<http://dbshino.nifs.ac.jp>). It will be necessary to develop recommended data for recombination processes including the new data in the near future.

### Acknowledgement

We would like to thank Dr. M. Arnaud for the useful information and discussions. We would like to thank Prof. K. Masai, Dr. M. Arnaud, Prof. T. Amano for providing us the codes to calculate the recombination rate coefficients. We are also grateful for Dr. R. More for reading the manuscript and helpful suggestions.

### References

1. T. Kato, K. Masai and M. Arnaud, NIFS-DATA-14 (1991)
2. e.g. "Recombination of Atomic Ions", NATO ASI series B 296 (Plenum Press, New York, 1992),  
e.g. I. Murakami and T. Kato, Dielectronic recombination, Journal of Plasma and Fusion Research, 74, 549, (1998), The Japan Society of Plasma Science and Nuclear Science (in Japanese)
3. S.M.V. Aldrovandi and D. Pequignot, Astron. & Astrophys. 25, 137 (1973)  
S.M.V. Aldrovandi and D. Pequignot, Revista Brasileira de Fisica, 4, 491 (1974)
4. R. Mewe, J. Schrijver and J. Sylwester, Astron. & Astrophys. Suppl. 40, 323 (1980)
5. R. Mewe and J. Schrijver, Astron. & Astrophys. Suppl. 40, 335 (1980), R. Mewe, J. Schrijver and J. Sylwester, Astron. & Astrophys. 87, 55 (1980)
6. K. Masai, private communication (1990)

7. R.A. Hulse, Nucl. Technol./Fusion, 3, 259 (1983)
8. M. Arnaud and R. Rothenflug, Astron. & Astrophys. Suppl. 60, 425 (1985)
9. M. Arnaud and J. Raymond, Astrophys. Journal 398, 394 (1992)
10. M.J. Seaton, Monthly Notices Roy. Astron. Soc. 119, 81 (1959)
11. S.M. Younger, J. Quant. Spectrosc. Radiat. Transfer 29, 67 (1983)
12. Y. Itikawa, T. Kato and K. Sakimoto, The Institute of Space and Astronautical Science Report No. 657, ISSN 0285-6808 (1995)
13. C.J. Romanik, Astrophys. J. 330, 1022 (1988)

### Figure captions

Fig.1 The recombination rate coefficients by Itikawa et al<sup>10</sup> for S ions of  $S^{12+}$  (Be-like -> B-like),  $S^{13+}$  (Li-like -> Be-like),  $S^{14+}$  (He-like -> Li-like) and  $S^{15+}$  (H-like -> He-like). The dotted line, dashed line and solid line indicate the radiative recombination, the dielectronic recombination and total recombination rate coefficients, respectively.

Fig.2 The recombination rate coefficients by Itikawa et al<sup>10</sup> for Ca ions of  $Ca^{16+}$  (Be-like -> B-like),  $Ca^{17+}$  (Li-like -> Be-like),  $Ca^{18+}$  (He-like -> Li-like) and  $Ca^{19+}$  (H-like -> He-like). The dotted line, dashed line and solid line indicate the radiative recombination, the dielectronic recombination and total recombination rate coefficients, respectively.

Fig.3 The recombination rate coefficients by Itikawa et al<sup>10</sup> for Ca ions of  $Fe^{22+}$  (Be-like -> B-like),  $Fe^{23+}$  (Li-like -> Be-like),  $Fe^{24+}$  (He-like -> Li-like) and  $Fe^{25+}$  (H-like -> He-like). The dotted line, dashed line and solid line indicate the radiative recombination, the dielectronic recombination and total recombination rate coefficients, respectively.




Fig.1











Fig.2










Fig.3



## **GRAPHS**































































































































































































































































































































































































































































## Publication List of NIFS-DATA Series

- NIFS-DATA-1 Y. Yamamura, T. Takiguchi and H. Tawara,  
*Data Compilation of Angular Distributions of Sputtered Atoms*; Jan 1990
- NIFS-DATA-2 T. Kato, J. Lang and K. E. Berrington,  
*Intensity Ratios of Emission Lines from OV Ions for Temperature and Density Diagnostics*, Mar 1990 [At Data and Nucl Data Tables 44(1990)133]
- NIFS-DATA-3 T. Kaneko,  
*Partial Electronic Straggling Cross Sections of Atoms for Protons*; Mar. 1990
- NIFS-DATA-4 T. Fujimoto, K. Sawada and K. Takahata,  
*Cross Section for Production of Excited Hydrogen Atoms Following Dissociative Excitation of Molecular Hydrogen by Electron Impact*; Mar. 1990
- NIFS-DATA-5 H. Tawara,  
*Some Electron Detachment Data for H<sup>-</sup> Ions in Collisions with Electrons, Ions, Atoms and Molecules – an Alternative Approach to High Energy Neutral Beam Production for Plasma Heating*; Apr 1990
- NIFS-DATA-6 H. Tawara, Y. Itikawa, H. Nishimura, H. Tanaka and Y. Nakamura,  
*Collision Data Involving Hydro-Carbon Molecules*; July 1990 [Supplement to Nucl Fusion 2(1992)25. Atomic and Molecular Processes in Magnetic Fusion Edge Plasmas (Plenum, 1995) p461]
- NIFS-DATA-7 H. Tawara,  
*Bibliography on Electron Transfer Processes in Ion-Ion/Atom/Molecule Collisions –Updated 1990*; Aug. 1990
- NIFS-DATA-8 U.I.Safranova, T.Kato, K.Masai, L.A.Vainshtein and A.S.Shlyapzeva,  
*Excitation Collision Strengths, Cross Sections and Rate Coefficients for OV, SiXI, FeXXIII, MoXXXIX by Electron Impact (1s<sup>2</sup>2s<sup>2</sup>-1s<sup>2</sup>2s2p-1s<sup>2</sup>2p<sup>2</sup> Transitions)* Dec.1990
- NIFS-DATA-9 T. Kaneko,  
*Partial and Total Electronic Stopping Cross Sections of Atoms and Solids for Protons*; Dec. 1990
- NIFS-DATA-10 K. Shima, N. Kuno, M. Yamanouchi and H. Tawara,  
*Equilibrium Charge Fraction of Ions of Z=4-92 (0.02-6 MeV/u) and Z=4-20 (Up to 40 MeV/u) Emerging from a Carbon Foil*; Jan.1991 [AT.Data and Nucl. Data Tables 51(1992)173]
- NIFS-DATA-11 T. Kaneko, T. Nishihara, T. Taguchi, K. Nakagawa, M. Murakami, M. Hosono, S. Matsushita, K. Hayase, M. Moriya, Y. Matsukuma, K. Miura and Hiro Tawara,  
*Partial and Total Electronic Stopping Cross Sections of Atoms for a Singly Charged Helium Ion: Part I*; Mar. 1991
- NIFS-DATA-12 Hiro Tawara,  
*Total and Partial Cross Sections of Electron Transfer Processes for Be<sup>q+</sup> and B<sup>q+</sup> Ions in Collisions with H, H<sub>2</sub> and He Gas Targets -Status in 1991-*; June 1991
- NIFS-DATA-13 T. Kaneko, M. Nishikori, N. Yamato, T. Fukushima, T. Fujikawa, S. Fujita, K. Miki, Y. Mitsunobu, K. Yasuhara, H. Yoshida and Hiro Tawara,  
*Partial and Total Electronic Stopping Cross Sections of Atoms for a Singly Charged Helium Ion : Part II*; Aug. 1991
- NIFS-DATA-14 T. Kato, K. Masai and M. Amaud,  
*Comparison of Ionization Rate Coefficients of Ions from Hydrogen through Nickel* : Sep. 1991
- NIFS-DATA-15 T. Kato, Y. Itikawa and K. Sakimoto,  
*Compilation of Excitation Cross Sections for He Atoms by Electron Impact*, Mar. 1992
- NIFS-DATA-16 T. Fujimoto, F. Koike, K. Sakimoto, R. Okasaka, K. Kawasaki, K. Takiyama, T. Oda and T. Kato,  
*Atomic Processes Relevant to Polarization Plasma Spectroscopy* ; Apr. 1992
- NIFS-DATA-17 H. Tawara,

*Electron Stripping Cross Sections for Light Impurity Ions in Colliding with Atomic Hydrogens Relevant to Fusion Research; Apr. 1992*

- NIFS-DATA-18 T. Kato,  
*Electron Impact Excitation Cross Sections and Effective Collision Strengths of N Atom and N-Like Ions -A Review of Available Data and Recommendations- ; Sep. 1992 [Atomic Data and Nuclear Data Tables, 57, 181-214 (1994)]*
- NIFS-DATA-19 Hiro Tawara,  
*Atomic and Molecular Data for H<sub>2</sub>O, CO & CO<sub>2</sub> Relevant to Edge Plasma Impurities, Oct. 1992*
- NIFS-DATA-20 Hiro. Tawara,  
*Bibliography on Electron Transfer Processes in Ion-Ion/Atom/Molecule Collisions -Updated 1993-; Apr. 1993*
- NIFS-DATA-21 J. Dubau and T. Kato,  
*Dielectronic Recombination Rate Coefficients to the Excited States of C I from C II; Aug. 1994*
- NIFS-DATA-22 T. Kawamura, T. Ono, Y. Yamamura,  
*Simulation Calculations of Physical Sputtering and Reflection Coefficient of Plasma-Irradiated Carbon Surface; Aug. 1994 [J. Nucl. Mater., 220 (1995) 1010]*
- NIFS-DATA-23 Y. Yamamura and H. Tawara,  
*Energy Dependence of Ion-Induced Sputtering Yields from Monoatomic Solids at Normal Incidence; Mar. 1995 [At. Data and Nud. Data Tables, 62 (1996) 149]*
- NIFS-DATA-24 T. Kato, U. Safranova, A. Shlyaptseva, M. Comille, J. Dubau,  
*Comparison of the Satellite Lines of H-like and He-like Spectra; Apr 1995 [Atomic Data and Nuclear Data Tables, 67 , 225 (1997)]*
- NIFS-DATA-25 H. Tawara,  
*Roles of Atomic and Molecular Processes in Fusion Plasma Researches - from the cradle (plasma production) to the grave (after-burning) -; May 1995*
- NIFS-DATA-26 N. Toshima and H. Tawara  
*Excitation, Ionization, and Electron Capture Cross Sections of Atomic Hydrogen in Collisions with Multiply Charged Ions; July 1995*
- NIFS-DATA-27 V.P. Shevelko, H. Tawara and E.Salzborn,  
*Multiple-Ionization Cross Sections of Atoms and Positive Ions by Electron Impact; July 1995 [Suppl. Nucl. Fusion, 6 (1996) 101]*
- NIFS-DATA-28 V.P. Shevelko and H. Tawara,  
*Cross Sections for Electron-Impact Induced Transitions Between Excited States in He: n, n'=2,3 and 4; Aug. 1995 [Suppl. Nucl. Fusion, 6 (1996) 27]*
- NIFS-DATA-29 U.I. Safranova, M.S. Safranova and T. Kato,  
*Cross Sections and Rate Coefficients for Excitation of  $\Delta n = 1$  Transitions in Li-like Ions with 6<Z<42; Sep. 1995 [Physica Scripta, 54, 68-84 (1996)]*
- NIFS-DATA-30 T. Nishikawa, T Kawachi, K. Nishihara and T. Fujimoto,  
*Recommended Atomic Data for Collisional-Radiative Model of Li-like Ions and Gain Calculation for Li-like Al Ions in the Recombining Plasma; Sep. 1995*
- NIFS-DATA-31 Y. Yamamura, K. Sakaoka and H. Tawara,  
*Computer Simulation and Data Compilation of Sputtering Yield by Hydrogen Isotopes (<sup>1</sup>H<sup>+</sup>, <sup>2</sup>D<sup>+</sup>, <sup>3</sup>T<sup>+</sup>) and Helium (<sup>4</sup>He<sup>+</sup>) Ion Impact from Monatomic Solids at Normal Incidence; Oct. 1995*
- NIFS-DATA-32 T. Kato, U. Safranova and M. Ohira,  
*Dielectronic Recombination Rate Coefficients to the Excited States of CII from CIII; Feb. 1996 [Physica Scripta, 53, 461-472 (1996), Physica Scripta, 55, 185-199 (1997)]*
- NIFS-DATA-33 K.J. Snowdon and H. Tawara,  
*Low Energy Molecule-Surface Interaction Processes of Relevance to Next-Generation Fusion Devices;*

- NIFS-DATA-34 T. Ono, T. Kawamura, K. Ishii and Y. Yamamura,  
*Sputtering Yield Formula for B<sub>4</sub>C Irradiated with Monoenergetic Ions at Normal Incidence*; Apr. 1996 [J Nucl. Mater., 232 (1996) 52]
- NIFS-DATA-35 I. Murakami, T. Kato and J. Dubau,  
*UV and X-Ray Spectral Lines of Be-Like Fe Ion for Plasma Diagnostics*; Apr. 1996 [Physica Scripta, 54, 463-470 (1996)]
- NIFS-DATA-36 K. Moribayashi and T. Kato,  
*Dielectronic Recombination of Be-like Fe Ion*; Apr. 1996 [Physca Scripta Vol 55, 286-297 (1997)]
- NIFS-DATA-37 U. Safronova, T. Kato and M. Ohira,  
*Dielectronic Recombination Rate Coefficients to the Excited States of CII from CIV*; July 1996 [J. Quant. Spectrosc. Radiat. Transfer, 58, 193 - 215, (1997)]
- NIFS-DATA-38 T. Fujimoto, H. Sahara, G. Csanak and S. Grabbe,  
*Atomic States and Collisional Relaxation in Plasma Polarization Spectroscopy: Axially Symmetric Case*; Oct. 1996
- NIFS-DATA-39 H. Tawara (Ed)  
*Present Status on Atomic and Molecular Data Relevant to Fusion Plasma Diagnostics and Modeling*; Jan. 1997
- NIFS-DATA-40 Inga Yu. Tolstikhina,  
*LS-Averaged I/Z Method as a Tool of Studying the Interactions of Highly Charged Ions with a Metal Surface*; Jan. 1997
- NIFS-DATA-41 K. Moribayashi and T. Kato,  
*Atomic Nuclear Charge Scaling for Dielectronic Recombination to Be-like Ions*; Apr. 1997
- NIFS-DATA-42 H. Tawara,  
*Bibliography on Electron Transfer Processes in Ion-ion / Atom / Molecule Collisions -Updated 1997-*; May 1997
- NIFS-DATA-43 M. Goto and T. Fujimoto,  
*Collisional-radiative Model for Neutral Helium in Plasma: Excitation Cross Section and Singlet-triplet Wavefunction Mixing*; Oct. 1997
- NIFS-DATA-44 J. Dubau, T. Kato and U. I. Safronova,  
*Dielectronic Recombination Rate Coefficients to the Excited States of CI From CII*; Jan. 1998
- NIFS-DATA-45 Y. Yamamura, W. Takeuchi and T. Kawamura,  
*The Screening Length of Interatomic Potential in Atomic Collisions*; Mar. 1998
- NIFS-DATA-46 T. Kenmotsu, T. Kawamura, T. Ono and Y. Yamamura,  
*Dynamical Simulation for Sputtering of B4C*; Mar. 1998
- NIFS-DATA-47 I. Murakami, K. Moribayashi and T. Kato,  
*Effect of Recombination Processes on FeXXIII Line Intensities*, May 1998
- NIFS-DATA-48 Zhijie Li, T. Kenmotsu, T. Kawamura, T. Ono and Y. Yamamura,  
*Sputtering Yield Calculations Using an Interatomic Potential with the Shell Effect and a New Local Model*; Oct. 1998
- NIFS-DATA-49 S. Sasaki, M. Goto, T. Kato and S. Takamura,  
*Line Intensity Ratios of Helium Atom in an Ionizing Plasma*; Oct. 1998
- NIFS-DATA-50 I. Murakami, T. Kato and U. Safronova,  
*Spectral Line Intensities of NeVII for Non-equilibrium Ionization Plasma Including Dielectronic Recombination Processes*; Jan. 1999
- NIFS-DATA-51 Hiro Tawara and Masa Kato,  
*Electron Impact Ionization Data for Atoms and Ions -up-dated in 1998-*, Feb. 1999

NIFS-DATA-52 J.G. Wang, T. Kato and I. Murakami,  
*Validity of  $n^{-3}$  Scaling Law in Dielectronic Recombination Processes;* Apr. 1999

NIFS-DATA-53 J.G. Wang, T. Kato and I. Murakami,  
*Dielectronic Recombination Rate Coefficients to Excited States of He from He<sup>+</sup>;* Apr. 1999

NIFS-DATA-54 T. Kato and E. Asano,  
*Comparison of Recombination Rate Coefficients Given by Empirical Formulas for Ions from Hydrogen through Nickel;* June 1999