

§3. Particle Acceleration in Large-amplitude MHD Waves Created by Strong Disturbances

Ohsawa, Y., Toida, M. (Dept. Physics, Nagoya Univ.), Ishiguro, S.

We have studied the evolution of magneto-hydrodynamic waves created by a strong disturbance in a collisionless plasma and particle acceleration caused by these waves: specifically, 1) evolution of large-amplitude Alfvén waves produced behind a magnetosonic shock front and ultrarelativistic electron acceleration occurring there, 2) effects of the inhomogeneity of the external magnetic field on positron acceleration in a shock wave in an electron-positron-ion plasma, 3) effects of ion composition on nonlinear evolution of oblique magnetosonic waves in two-ion-species plasmas, and 4) multi-dimensional effects on electron trapping and acceleration in an oblique shock wave.

1) Evolution of large-amplitude Alfvén waves behind a magnetosonic shock front and associated ultrarelativistic electron acceleration

A strong disturbance in a magnetized plasma produces a magnetosonic shock wave. Moreover, large-amplitude Alfvén waves are generated behind the shock front. We have observed the evolution of these waves with one-dimensional, fully kinetic, relativistic, electromagnetic simulations. Furthermore, we have recognized three types of ultrarelativistic electron acceleration caused by these Alfvén waves [1]. These energization processes can take place in both weak and strong external magnetic fields.




Fig. 1. Wave trajectories.

Figure 1 shows the trajectories of the waves created by a strong disturbance. The two thick lines indicate forward (right going) and backward (left going) shock fronts, while the thin lines represent the trajectories of Alfvén waves; the dotted line is the trajectory of a strong-magnetic-field (SMF) pulse, which is thought to be developing into the Alfvén wave.

Figure 2 displays magnetic-field profiles (the top and




Fig. 2. Snap shots of magnetic and density profiles and electron phase space (x, γ).

second panels), electron density (third panel), and electron phase space (x, γ) (bottom panel), where γ is the Lorentz factor. In this figure, the forward and backward shock fronts are near $x/(c/\omega_{pe})=2370$ and 2020 , respectively. The SMF pulse is near $x/(c/\omega_{pe})=2230$, and Alfvén waves are on its left. In the Alfvén wave region including the SMF pulse, we find ultrarelativistic electrons with $\gamma \sim 300$.

2) Positron acceleration

Positron acceleration to $\gamma \sim 10^4$ by a shock wave has been demonstrated with particle simulations with real ion-to-electron mass ratio, $m_i/m_e=1836$. Furthermore, effects of the spatial gradient of the external magnetic field on the acceleration has been investigated in detail [2].

3) Oblique magnetosonic waves in two-ion-species plasmas

The study of the effects of ion composition on perpendicular magnetosonic waves [3] has been extended to oblique waves [4].

4) Multi-dimensional simulation of electron trapping

Multi-dimensional effects on electrons in an oblique shock wave have been investigated in detail by means of two-dimensional electromagnetic particle simulations and test particle calculations. It has been shown that whistler wave instabilities can cause detrapping of energetic electrons from the shock wave [5].

[1] Y. Takeyama, S. Nakayama, and Y. Ohsawa, Phys. Plasmas **18**, 092307 (2011).

[2] T. Iwata, S. Takahashi, and Y. Ohsawa, Phys. Plasmas **19**, 022302 (2012).

[3] M. Toida, Y. Ohsawa, and T. Jyounouchi, Phys. Plasmas **2**, 3329 (1995); M. Toida, H. Higashino, and Y. Ohsawa, J. Phys. Soc. Jpn **76**, 104052 (2007).

[4] M. Toida and Y. Kondo, Phys. Plasmas **18**, 062303 (2011).

[5] M. Toida and J. Joho, submitted to J. Phys. Soc. Jpn.